



Hungarian University of Agriculture and Life Sciences

PhD DISSERTATION

Adnan Eser

Gödöllő

2022



Hungarian University of Agriculture and Life Sciences

**IMPACT OF NITROGEN TOPDRESSING ON THE QUALITY PARAMETERS OF
WINTER WHEAT (*TRITICUM AESTIVUM L.*) YIELD**

DOI: 10.54598/003390

PhD DISSERTATION

Adnan Eser

Gödöllő

2022

The PhD School

Name: Doctoral School of Plant Sciences

Discipline: Crop Production and Horticultural Sciences

Head: Prof. Lajos Helyes, DSc
Faculty of Agricultural and Environmental Sciences
Department of Horticulture and Technology
Hungarian University of Agriculture and Life Sciences

Supervisor(s): Prof. emeritus Márton Jolánkai, DSc
Faculty of Agricultural and Environmental Sciences
Institute of Crop Production

.....

Dr Lajos Helyes

Head of Doctoral School

.....

Dr Márton Jolánkai

Supervisor

1. TABLE OF CONTENTS

| | |
|---|------|
| 1. TABLE OF CONTENTS | iii |
| 1.1 Table of figures | v |
| 1.2. Table of tables | vii |
| 2. INTRODUCTION | x |
| 2.1. ACRONYMS | xiii |
| 3. OBJECTIVES TO ARCHIVE | 14 |
| 4. LITERATURE OVERVIEW | 16 |
| 4.1. Importance of the wheat | 16 |
| 4.2. Quality criteria of the wheat | 19 |
| 4.2.1. Factors that affect the quality of wheat | 19 |
| 4.2.1.1. Genotype | 19 |
| 4.2.1.2. Seed quality | 20 |
| 4.2.1.3. Environmental factors | 22 |
| 4.3. Ecological requirements of the winter wheat | 23 |
| 4.4. The grain protein | 24 |
| 4.3.1. Quality of the protein | 26 |
| 4.5. The nitrogen fertilizer | 30 |
| 4.5.1. Effect of nitrogen on quality, yield and development | 30 |
| 4.5.2. Other effects of nitrogen fertilizer | 31 |
| 4.5.3. Negative effects of nitrogen fertilizer | 32 |
| 4.5.4. Efficiency of the split dose application of nitrogen fertilizer | 33 |
| 4.5.5. Amount of the nitrogen application | 34 |
| 4.6. The grain moisture | 36 |
| 5. MATERIALS AND METHODS | 37 |
| 5.1. Field features of experiments | 37 |
| 5.2. Meteorological properties of the experimental fields | 39 |
| 5.2.1 Hatvan-Nagyombos meteorological properties | 39 |
| 5.2.2. Gödöllő meteorological properties | 43 |
| 5.3. Studied winter wheat varieties | 45 |
| 5.4. Treatments | 46 |
| 5.5. Investigations | 47 |
| 5.6. Statistical analyses | 47 |
| 6. RESULTS | 48 |
| 6.1. Test Weight Results | 48 |

| | |
|--|-----------|
| 6.2. Thousand Kernel Weight Values..... | 50 |
| 6.3. The grain moisture..... | 53 |
| 6.4. The grain protein | 55 |
| 6.5. The gluten content..... | 61 |
| 6.6 The Zeleny number (ml) | 66 |
| 7. CONCLUSION AND RECOMMENDATIONS | 71 |
| 8. NEW SCIENTIFIC RESULTS | 73 |
| 9. THE PUBLICATIONS OF THE AUTHOR IN THE RESEARCH FIELD | 76 |
| 10. BIBLIOGRAPHY..... | 80 |
| 11. ACKNOWLEDGEMENT | 97 |
| 12. APPENDIX 1 – TABLES AND FIGURES | 98 |
| 12.1. Test Weight | 98 |
| 12.2. Thousand Kernel Weight..... | 100 |
| 12.3. The Grain Moisture | 103 |
| 12.4. Protein Content..... | 104 |
| 12.5. Gluten Content..... | 107 |
| 12.6. Anova-Spss outputs..... | 112 |
| 12.6.1. Examination of Parameters According to Categorical Variables in Different Years..... | 112 |
| 12.6.2. Examination of Parameters According to Categorical Variables in Different Sites-Hatvan | 112 |
| 12.6.3. Examination of Parameters According to Categorical Variables in Different Sites-Gödöllő | 115 |
| 12.6.4. Examination of Parameters According to Categorical Variables in Different Genotypes..... | 116 |

1.1 Table of figures

| | |
|---|----|
| 1. Figure Nitrogen uptake of wheat during the growth period (Brown et al., 2005; Orlof et al., 2012)..... | 15 |
| 2. Figure Wheat growth stages for Zadoks and Feekes scale (Large, 1954; Zadoks et al., 1974)..... | 16 |
| 3. Figure Effect of seed quality on grain yield, Kumar. D, and Seth. R, (2004)..... | 21 |
| 4. Figure Effect of seed size on seedling vigor in wheat Simmone et. al., (2000)..... | 22 |
| 5. Figure Effect of seed size on germination percentage Hossein et. al., (2011)..... | 22 |
| 6. Figure Wheat gluten protein and its impacts on wheat processing quality. Ma et al., (2019). Frontiers of Agricultural Science and Engineering..... | 25 |
| 7. Figure Total protein yields in the non-favorable crop year. Nagygompos 2014. Eser et. al. (2017)..... | 26 |
| 8. Figure Remix loaf volume results for flour samples of two Canadian genotypes (Manitou and the unnamed line 11-463A), with a range of protein contents. Baking quality is indicated as loaf volume (L.V.) and protein content is expressed on the basis of 14% moisture basis (m.b.). Bushuk et al. (1969). | 28 |
| 9. Figure Baking quality of wheat is determined by wheat cultivar, the interplay of natural grain constituents' proteins, carbohydrates and lipids, growing conditions of wheat as well as milling and baking technology. Langenkämper et al.,(2019)..... | 29 |
| 10. Figure Effect of applied nitrogen on winter wheat grain protein. Yara, (2011)..... | 35 |
| 11. Figure The experimental field view by satellite at 2016-2017, Nagygompos, Hungary..... | 38 |
| 12. Figure The experimental field view by satellite at 2019-2020, Gödöllő, Hungary..... | 39 |
| 13. Figure Max, Min and Average Weather Temperature 2015-2016 (Nagygompos, Hungary). worldweatheronline.com..... | 40 |
| 14. Figure Max, Min and Average Weather Temperature 2016-2017 (Nagygompos, Hungary). worldweatheronline.com..... | 41 |
| 15. Figure Rainfall and Rain Days 2015-2016 (Nagygompos, Hungary). worldweatheronline.com..... | 42 |
| 16. Figure Rainfall and Rain Days 2016-2017 (Nagygompos, Hungary). worldweatheronline.com..... | 43 |
| 17. Figure Max, Min and Average Weather Temperature 2018-2019 (Gödöllő, Hungary). worldweatheronline.com..... | 44 |
| 18. Figure Rainfall and Rain Days 2018-2019 (Gödöllő, Hungary). worldweatheronline.com... | 45 |
| 19. Figure Impact of N topdressing applications on moisture content of wheat varieties. 2015- 2016 and 2016–2017 (Nagygompos, Hungary)..... | 53 |
| 20. Figure Impact of N topdressing applications on moisture content of wheat varieties. 2018- 2019 (Gödöllő, Hungary)..... | 54 |
| 21. Figure Impact of N topdressing applications on protein content of wheat varieties. 2015-2016 and 2016–2017 (Nagygompos, Hungary) and 2018-2019 (Gödöllő, Hungary) combined..... | 57 |
| 22. Figure Impact of N topdressing applications on protein content of wheat varieties untreated plots. 2015-2016 and 2016–2017 (Nagygompos, Hungary)..... | 58 |
| 23. Figure Impact of N topdressing applications on Zeleny Nr. of wheat varieties. 2015-2016 and 2016–2017 (Nagygompos, Hungary) 1..... | 66 |
| 24. Figure Impact of N topdressing applications on wheat grain test weight. 2015-2016 (Nagygompos, Hungary)..... | 98 |
| 25. Figure Impact of N topdressing applications on wheat grain test weight. 2016-2017 (Nagygompos, Hungary)..... | 99 |

| | |
|---|-----|
| 26. Figure Impact of N topdressing applications on wheat grain test weight. 2018-2019 (Gödöllő, Hungary) 2 | 99 |
| 27. Figure Impact of N topdressing applications on wheat grain test weight. 2018–2019 (Gödöllő, Hungary) Box Plot Alföld and MV Karéj..... | 100 |
| 28. Figure Impact of N topdressing applications on thousand kernel weight of wheat varieties. 2015-2016 (Nagygyombos, Hungary)..... | 101 |
| 29. Figure Impact of N topdressing applications on thousand kernel weight of wheat varieties. 2016-2017 (Nagygyombos, Hungary)..... | 101 |
| 30. Figure Impact of N topdressing applications on thousand kernel weight of wheat varieties. 2018-2019 (Gödöllő, Hungary) 2..... | 102 |
| 31. Figure Impact of N topdressing applications on thousand kernel weight of wheat varieties. 2018-2019 (Gödöllő, Hungary) 3..... | 103 |
| 32. Figure Impact of N topdressing applications on moisture content of wheat varieties. 2018-2019 (Gödöllő, Hungary) 3..... | 104 |
| 33. Figure Impact of N topdressing applications on protein content of wheat varieties. 2018–2019 (Gödöllő, Hungary) 2..... | 105 |
| 34. Figure Impact of N topdressing applications on protein content in Alföld. 2015-2016 and 2016–2017 (Nagygyombos, Hungary) and 2018-2019 (Gödöllő, Hungary) combined | 106 |
| 35. Figure Impact of N topdressing applications on protein content of MV Karéj. 2015-2016 and 2016–2017 (Nagygyombos, Hungary) and 2018-2019 (Gödöllő, Hungary) combined | 107 |
| 36. Figure Impact of N topdressing applications on gluten content of wheat varieties. 2015-2016 (Nagygyombos, Hungary) | 108 |
| 37. Figure Impact of N topdressing applications on gluten content of wheat varieties. 2016-2017 (Nagygyombos, Hungary) | 108 |
| 38. Figure Impact of N topdressing applications on gluten content of wheat varieties. 2018-2019 (Gödöllő, Hungary) 2 | 109 |
| 39. Figure Impact of N topdressing applications on gluten content of wheat varieties. 2018–2019 (Gödöllő, Hungary) 3 | 110 |
| 40. Figure Impact of N topdressing applications on gluten content of wheat varieties. 2015-2016 and 2016–2017 (Nagygyombos, Hungary) and 2018-2019 (Gödöllő, Hungary) combined..... | 111 |

1.2. Table of tables

| | |
|--|----|
| 1. Table GIEWS Crop Prospects And Food Situation: World cereal production, FAO 2021 Note: Includes rice in milled terms. Totals and percentage change computed from unrounded data. ¹ Data for the European Union from the year 2020 (including the 2020/21 marketing year) excludes the United Kingdom of Great Britain and Northern Ireland. | 17 |
| 2. Table GIEWS Crop Prospects and Food Situation: Wheat production: Leading producers, FAO 2021 (¹ Data for the European Union prior the year 2020 includes the United Kingdom of Great Britain and Northern Ireland.)..... | 18 |
| 3. Table Soil type of the experimental field at Hungarian University of Agriculture and Life Sciences, Crop Production Institute, Nagygyombos, Hungary..... | 38 |
| 4. Table Soil type of the experimental field at Hungarian University of Agriculture and Life Sciences, Crop Production Institute, Gödöllő, Hungary | 39 |
| 5. Table Martonvásári Fajtakatalógus 2020..... | 46 |
| 6. Table Impact of N topdressing applications on wheat grain test weight. 2015-2016 and 2016–2017 (Nagygyombos, Hungary)..... | 48 |
| 7. Table Impact of N topdressing applications on wheat grain test weight. 2018–2019 (Gödöllő, Hungary) | 49 |
| 8. Table Comparison of the Means of Test Weight Parameters Measured in the Sample of All Genotypes Based on Categorical Variables, 2016, 2017 Hatvan and 2019 Gödöllő crop years. . | 50 |
| 9. Table Impact of N topdressing applications on thousand kernel weight of wheat varieties. 2015-2016 and 2016–2017 (Nagygyombos, Hungary) | 51 |
| 10. Table Impact of N topdressing applications on thousand kernel weight of wheat varieties. 2018-2019 (Gödöllő, Hungary) 1 | 52 |
| 11. Table Comparison of the Means of Thousand Kernel Weight Parameters Measured in the Sample of All Genotypes Based on Categorical Variables, 2016,2017 Hatvan and 2019 Gödöllő crop years. | 53 |
| 12. Table Comparison of the Means of Moisture Content Parameters Measured in the Sample of All Genotypes Based on Categorical Variables, 2016 and 2017 Hatvan and 2019 Gödöllő crop years. | 55 |
| 13. Table Impact of N topdressing applications on protein content of wheat varieties. 2015-2016 and 2016–2017 (Nagygyombos, Hungary)..... | 56 |
| 14. Table Impact of N topdressing applications on protein content of wheat varieties. 2018–2019 (Gödöllő, Hungary) | 56 |
| 15. Table Comparison of the Means of Protein Content Parameters Measured in the Sample of All Genotypes Based on Categorical Variables, 2016,2017 Hatvan and 2019 Gödöllő crop years. | 59 |
| 16. Table Comparison of the Means of Protein Content Parameters Measured in the Sample of All Genotypes in 2016, Based on Categorical Variables, Hatvan site. | 60 |
| 17. Table Comparison of the Means of Protein Content Parameters Measured in the Sample of Alföld and Karéj Genotypes in Gödöllő in 2019, Based on Categorical Variables, Gödöllő site. | 60 |
| 18. Table Comparison of the Means of Protein Content Parameters Measured in the Sample of Karej Genotype Based on Categorical Variables | 61 |
| 19. Table Impact of N topdressing applications on gluten content of wheat varieties. 2015-2016 and 2016–2017 (Nagygyombos, Hungary)..... | 62 |
| 20. Table Impact of N topdressing applications on gluten content of wheat varieties. 2018–2019 (Gödöllő, Hungary) 1 | 62 |
| 21. Table Comparison of the Means of Gluten Content Parameters Measured in the Sample of All Genotypes Based on Categorical Variables | 63 |

| | |
|---|-----|
| 22. Table Comparison of the Means of Gluten Content Parameters Measured in the Sample of All Genotypes in 2016 Based on Categorical Variables | 64 |
| 23. Table Comparison of the Means of Gluten Content Parameters Measured in the Sample of Toldi Genotype Based on Categorical Variables | 64 |
| 24. Table Comparison of the Means of Gluten Content Parameters Measured in the Sample of Alföld and Karéj Genotypes in Gödöllő in 2019 Based on Categorical Variables | 65 |
| 25. Table Impact of N topdressing applications on Zeleny Nr. of wheat varieties. 2015-2016 and 2016–2017 (Nagygyombos, Hungary) 2 | 67 |
| 26. Table Impact of N topdressing applications on Zeleny Nr. of wheat varieties. 2015-2016 and 2018–2019 (Gödöllő, Hungary) | 67 |
| 27. Table Comparison of the Means of Zeleny Nr. Parameters Measured in the Sample of All Genotypes Based on Categorical Variables | 68 |
| 28. Table Comparison of the Means of Zeleny Nr. Parameters Measured in the Sample of Alföld and Karéj Genotypes in Gödöllő in 2019 Based on Categorical Variables | 68 |
| 29. Table Comparison of the Means of Zeleny Nr. Parameters Measured in the Sample of Alföld and Karéj Genotypes in 2016 Based on Categorical Variables | 69 |
| 30. Table Comparison of the Means of Zeleny Nr. Parameters Measured in the Sample of Karéj Genotype Based on Categorical Variables | 70 |
| 31. Table Impact of N topdressing applications on moisture content of wheat varieties. 2015-2016 (Nagygyombos, Hungary) | 103 |
| 32. Table Impact of N topdressing applications on moisture content of wheat varieties. 2016-2017 (Nagygyombos, Hungary) | 103 |
| 33. Table Impact of N topdressing applications on moisture content of wheat varieties. 2018-2019 (Gödöllő, Hungary) 2 | 104 |
| 34. Table Impact of N topdressing applications on protein content of wheat varieties. 2015-2016 (Nagygyombos, Hungary) | 105 |
| 35. Table Impact of N topdressing applications on protein content of wheat varieties. 2016-2017 (Nagygyombos, Hungary) | 105 |
| 36. Table Impact of undivided/split N topdressing applications on wheat grain based on years, 2015-2016, 2016-2017 and 2018-2019 crop seasons, Spss-One Way Anova | 112 |
| 37. Table Impact of undivided/split N topdressing applications on wheat grain based on Hatvan-Nagygyombos site, 2015-2016 and 2016-2017 crop seasons, Spss-One Way Anova | 113 |
| 38. Table Impact of undivided/split N topdressing applications on wheat grain based on Hatvan-Nagygyombos site by Genotype, 2015-2016 and 2016-2017 crop seasons, Spss-One Way Anova | 113 |
| 39. Table Impact of undivided/split N topdressing applications on wheat grain, based on Hatvan-Nagygyombos site by Treatments, 2015-2016 and 2016-2017 crop seasons, Spss-One Way Anova | 114 |
| 40. Table Impact of undivided/split N topdressing applications on wheat grain based on Gödöllő site by Genotype, 2018-2019 crop seasons, Spss-One Way Anova | 115 |
| 41. Table Impact of undivided/split N topdressing applications on wheat grain based on Gödöllő site by Treatment, 2018-2019 crop seasons, Spss-One Way Anova | 116 |
| 42. Table Impact of undivided/split N topdressing applications on wheat grain based on Alföld genotype by years, 2015-2016, 2016-2017 and 2018-2019 crop seasons, Spss-One Way Anova | 117 |
| 43. Table Impact of undivided/split N topdressing applications on wheat grain based on Alföld genotype by Site, 2015-2016, 2016-2017 and 2018-2019 crop seasons, Spss-One Way Anova | 117 |

| | |
|---|-----|
| 44. Table Impact of undivided/split N topdressing applications on wheat grain based on Alföld genotype by Treatment, 2015-2016, 2016-2017 and 2018-2019 crop seasons, Spss-One Way Anova..... | 119 |
| 45. Table Impact of undivided/split N topdressing applications on wheat grain based on Karéj genotype by Year, 2015-2016, 2016-2017 and 2018-2019 crop seasons, Spss-One Way Anova | 119 |
| 46. Table Impact of undivided/split N topdressing applications on wheat grain based on Karéj genotype by Site, 2015-2016, 2016-2017 and 2018-2019 crop seasons, Spss-One Way Anova | 120 |
| 47. Table Impact of undivided/split N topdressing applications on wheat grain based on Karéj genotype by Treatment, 2015-2016, 2016-2017 and 2018-2019 crop seasons, Spss-One Way Anova..... | 121 |
| 48. Table Impact of undivided/split N topdressing applications on wheat grain, 2015-2016 and 2016-2017 crop seasons, Spss-One Way Anova..... | 126 |
| 49. Table Impact of undivided/split N topdressing applications on wheat grain, 2018-2019 crop seasons 0kg/ha, Spss-One Way Anova..... | 127 |
| 50. Table Impact of undivided/split N topdressing applications on wheat grain, 2018-2019 crop seasons 80kg/ha, Spss-One Way Anova..... | 128 |
| 51. Table Impact of undivided/split N topdressing applications on wheat grain, 2018-2019 crop seasons 80+40 kg/ha, Spss-One Way Anova..... | 129 |
| 52. Table Impact of undivided/split N topdressing applications on wheat grain, 2018-2019 crop seasons 120 kg/ha, Spss-One Way Anova..... | 130 |
| 53. Table Impact of undivided/split N topdressing applications on wheat grain, 2018-2019 crop seasons 120+40 kg/ha, Spss-One Way Anova..... | 131 |
| 54. Table Impact of undivided/split N topdressing applications on wheat grain, 2018-2019 crop seasons 160 kg/ha, Spss-One Way Anova..... | 133 |

2. INTRODUCTION

The Earth's rising population growth is contributing to increasing hunger, as well as creating insufficient and unbalanced nutrition, all of which continue to be major problems for human survival. Although different opinions are put forward for the solution, the most notable consensus among experts has been to engineer an increase in plant and animal products. Wheat, the most widely grown plant species in the world, is a plant of high strategic importance due to the fact that it has been grown since the earliest times; its agricultural process is easier than other plants; the product is more adaptable to transportation and storage conditions; and its inherent high nutrition solves one of the most pressing economic problems. Wheat is also one of the most predominant cereal grains in Hungary and Turkey, and it retains high economic value. The goal of wheat production is two-fold: to provide both quantity and quality.

The milling and baking qualities of wheat are mainly determined by its genetic basis; however, it can be influenced by management techniques as well (Pollhamer, 1981; Grimwade et al., 1996; Vida et al., 1996; Pepó, 2010). The determination of wheat milling quality is complex because the quality measurements depend on the kernel hardness, protein, starch, internal insect infestation, color, disease, size, and moisture parameters (Posner, 2003). One of the most important parameters, in terms of nutritional content, is the protein ratio in wheat. Although it is determined genetically, the protein ratio is a feature that can also be increased by Nitrogen fertilizer applications. The desired ratio is 11-14% for red hard winter wheat. For optimum grain yield, the critical protein ratio was specified as 11.5% (Goos et al., 1982). However, it is required that this rate should not be less than 12.5% in order to increase the quality of bread. Baking quality of wheat flour is determined by grain protein concentration (GPC) and its composition, and is highly influenced by environmental factors such as Nitrogen (N) fertilization management (Xue et al., 2019). The protein content of wheat crops has an important impact on the nutritional quality for humans and livestock and on the crops' functional properties in food processing (Shewry and Halford, 2002). The economic value of winter wheat is affected by the genotype, cropping year, agro-climatic parameters, as well as the agronomic applications and coordination (Gyóri, 2006; Várallyay, 2008).

Varieties that differ greatly in terms of yield cause less variation in grain protein yields. Since the grain protein concentration is strongly related to the relationship between protein yield and grain yield, there is a somewhat significant inverse relationship between yield and protein

concentration from this combination. The protein yield of wheat has increased significantly throughout its breeding history but could not keep up with the genetic yield increase in protein yield. The result was low protein concentrations (Van Lill and Purchase, 1995). Nitrogen (N) is one of the macronutrients required for plant growth, with high effect on quality and quantity values of winter wheat. The quality of wheat varieties is strongly influenced by crop year, season, genotype effects, and the effects of the management systems are also determinative of some physical and gluten quality characteristics of the grain (Rakszegi et al., 2016). Horváth and co-workers (2014) also established that increasing levels of N topdressing and increased number of applications had beneficial effects on the protein content as well as on wet gluten values of wheat grain. Szentpétery and co-workers (2005) proved that increasing fertilizer dose applications had a beneficial effect on the protein and gluten contents, as well as improvements in quality. Kismányoky and Tóth (2010) explained that the increasing rate of N fertilization application, as well as the additional organic fertilizers, has influenced the biomass production and N uptake of winter wheat. Additionally, as a result of intensive chemical fertilization, the amounts of organic matter and humus in the soil will decrease, and the fertilizers will be washed away as they cannot hold in the soil. Plant nutrients (fertilizer) will not be transformed into a chemical form in the soil that the plant could uptake; thus, the physical and chemical properties of the soil will deteriorate. As a result, the upper parts of the soil will be sandy while the lower parts will be stony. One of the biggest factors that cause environmental pollution and deterioration of the natural balance is agricultural activity where chemicals are used extensively. Moreover, the agricultural methods in which chemicals are used not only cause environmental pollution and deterioration of the natural balance, but also threaten the lives of all living things via the food chain. The aim of this study is to investigate the changes in qualitative parameters of the winter wheat varieties sown in four crop seasons, each of which had different levels of split and/or undivided dose applications of Nitrogen fertilizers. This practice would determine the optimum Nitrogen ratio required in order to reach the optimum protein rate in Nitrogen fertilizer applications with variable ratios. With the help of the equations obtained within the research, the aim is to determine the most appropriate Nitrogen dosage, to reach the highest level of efficiency and quality, and to apply the Nitrogen use at the best possible levels. Due to the fact that the Nitrogen requirement of the plant is not fully met, the results would include low yield, excessive fertilization, pollution of the environment, and economic losses due to the cost of fertilizers. A study conducted in France stated that Nitrogen fertilizer costs constitute 28% of wheat production (Quievreux, 1997). In temperate regions, wheat is the main crop grown for both human and animal consumption. Therefore, all research that focuses on raising wheat production efficiency, crop quantity, and quality

characteristics is of utmost significance. The aim of this study is to find the quality changes on tested winter wheat varieties with the application of Nitrogen fertilizer. As Nitrogen fertilization has a considerably high economic and environmental effect on wheat farming and wheat quality, we therefore should find the most appropriate formula for supplying fertilizer to get the best results to be able to improve the quality of the winter wheat products, as well as consider the economic and environmental side effects of the this kind of farming approach.


2.1. ACRONYMS

| | | |
|------|---|---|
| M | : | Arithmetic Mean |
| SPSS | : | Statistical Program for Social Sciences |
| SD | : | Standard Deviation |
| df | : | Degrees of Freedom |
| N | : | Frequency |
| % | : | Percent |
| p | : | Significance Value |
| t | : | t statistic |
| F | : | F statistic |
| r | : | Correlation Coefficient |


3. OBJECTIVES TO ARCHIVE

It is possible to divide the development period of wheat into three periods depending on the nitrogen intake amount. The first period is from emergence to tillering, when nitrogen uptake is low (5-20%). Research has shown that the nitrogen of wheat is taken most intensively in the (Figure 1) period of tillering and earing (Zadoks' 25-58) according to the Figure 2 growth period in Zadoks (Brown et al., 2005; Orlof et al., 2012). Nitrogen in this period contributes positively to the amount of tillering and the number of grains per ear, which directly affects the yield. As seen in Figure 1, the third period is the period from spiking to harvest, and nitrogen uptake is slow in this period as well. The main purpose of fertilization is to keep the nitrogen needed by the wheat plant in the soil ready in time during its growth and development periods. For this reason, making fertilization according to growth periods increases the efficiency of nitrogen use. On the negative side, applying more nitrogen than the appropriate nitrogen dose can

encourage vegetative growth, which causes a decrease in yield.



1. Figure Nitrogen uptake of wheat during the growth period (Brown et al., 2005; Orlof et al., 2012)



2. Figure Wheat growth stages for Zadoks and Feekes scale (Large, 1954; Zadoks et al., 1974)

4. LITERATURE OVERVIEW

4.1. Importance of the wheat

There are more lands planted with wheat in the world than with any other crop. It provides 20 percent of the world's caloric consumption; even 50 percent for the world's poorest, for whom it also provides 20 percent of their protein consumption (Washington Wheat Facts 2016/2017).

| (Million tonnes) | 2018 | 2019 | 2020 estimate | Change: 2020 over 2019 (%) |
|------------------|---------|---------|---------------|----------------------------|
| Asia | 1 185.1 | 1 196.0 | 1 220.6 | 2.1 |
| Far East | 1 085.7 | 1 089.3 | 1 107.4 | 1.7 |
| Near East | 65.2 | 73.5 | 78.0 | 6.2 |
| CIS in Asia | 34.2 | 33.2 | 35.2 | 6.1 |
| Africa | 198.6 | 192.4 | 198.9 | 3.4 |

| | | | | |
|-----------------------------------|---------|---------|---------|-------|
| North Africa | 37.5 | 36.2 | 32.7 | -9.5 |
| West Africa | 66.0 | 65.5 | 66.9 | 2.1 |
| Central Africa | 6.8 | 6.7 | 6.6 | -2.7 |
| East Africa | 56.6 | 55.4 | 56.4 | 1.8 |
| Southern Africa | 31.8 | 28.6 | 36.3 | 27.0 |
| Central America and the Caribbean | 42.5 | 42.4 | 42.7 | 0.6 |
| South America | 197.8 | 228.8 | 233.4 | 2.0 |
| North America | 495.4 | 479.3 | 496.8 | 3.7 |
| Europe | 497.2 | 542.0 | 518.9 | -4.3 |
| European Union ¹ | 294.2 | 324.1 | 281.5 | -13.1 |
| CIS in Europe | 188.0 | 202.7 | 202.2 | -0.2 |
| Oceania | 30.9 | 27.9 | 50.0 | 79.0 |
| World | 2 647.4 | 2 708.8 | 2 761.3 | 1.9 |
| Developing countries | 1 615.3 | 1 650.7 | 1 686.8 | 2.2 |
| Developed countries | 1 032.1 | 1 058.1 | 1 074.5 | 1.6 |
| - wheat | 732.1 | 760.7 | 774.0 | 1.7 |
| - coarse grains | 1 408.0 | 1 445.3 | 1 474.1 | 2.0 |
| - rice (milled) | 507.3 | 502.8 | 513.2 | 2.1 |

1. Table GIEWS Crop Prospects And Food Situation: World cereal production, FAO 2021

Note: Includes rice in milled terms. Totals and percentage change computed from unrounded data. ¹Data for the European Union from the year 2020 (including the 2020/21 marketing year) excludes the United Kingdom of Great Britain and Northern Ireland.

The total global wheat output exceeded 776.7 million tons in 2021, according to FAOSTAT data (FAO, 2022). All over the world, 18.3% of daily calories and 19.8% of protein taken from foods are provided from wheat products (FAO, 2018). Wheat is the dominant crop in temperate countries being used for human food and livestock feed. Its success depends partly on its adaptability and high yield potential but also on the gluten protein fraction which confers the viscoelastic properties that allow dough to be processed into bread, pasta, noodles, and other food products. Globally, protein deficiency is a significant problem that takes second place to calorie deficiency in human nutrition (Robinson et al., 1986) – Table 1 shows that wheat is one of most widely produced crop products. Wheat also contributes essential amino acids, minerals, and vitamins, and beneficial phytochemicals and dietary fiber components to the human diet,

and these are particularly abundant in whole-grain products (Shewry, 2009). Improving global food security is critical, given that over 800 million people remain food insecure (FAO, 2017) and its association with conflict and civil unrest (Kalkuhl et al., 2016; Bellemare, 2015). The leading wheat producers in the world are the EU, China, USA, Russia, India and Turkey, as shown in Table 2; however, wheat is not suitable for cultivation in certain parts of the world for climatic reasons. Some countries are completely foreign-dependent for this reason. Chief among these are Egypt, Indonesia, Algeria, Brazil, Japan and S. Korea, which import the most (Atar, 2017).

| (Million tonnes) | Average 5yrs | 2019 | 2020 estimate | 2021 forecast |
|--|--------------|-------|---------------|---------------|
| European Union ¹ | 143.1 | 155.7 | 125.2 | 137.0 |
| China (mainland) | 133.4 | 133.6 | 134.2 | 135.5 |
| India | 100.4 | 103.6 | 107.6 | 109.0 |
| Russian Federation | 78.4 | 74.5 | 85.9 | 79.0 |
| USA | 52.8 | 52.6 | 49.7 | 50.0 |
| Canada | 32.5 | 32.3 | 35.2 | 33.0 |
| Ukraine | 26.1 | 28.3 | 25.1 | 26.0 |
| Pakistan | 25.4 | 24.4 | 25.3 | 26.0 |
| Australia | 23.8 | 15.2 | 33.3 | 26.0 |
| Turkey | 20.3 | 19.0 | 20.5 | 19.5 |
| Argentina | 18.7 | 19.8 | 17.6 | 19.0 |
| Iran | 14.3 | 14.5 | 14.0 | 14.5 |
| Kazakhstan | 13.9 | 11.5 | 14.3 | 14.0 |
| United Kingdom of Great Britain and Northern Ireland | - | - | 9.7 | 14.0 |
| Egypt | 8.8 | 9.0 | 9.0 | 9.0 |
| Other countries | 66.6 | 66.9 | 67.5 | 68.5 |
| World | 758.4 | 760.7 | 774.0 | 780.0 |

2. Table *GIEWS Crop Prospects and Food Situation: Wheat production: Leading producers, FAO 2021* (¹Data for the European Union prior the year 2020 includes the United Kingdom of Great Britain and Northern Ireland.)

4.2. Quality criteria of the wheat

- Variety
- Test weight
- Thousand grain weight
- Vitreousness ratio
- Ash amount
- Protein and gluten content
- Pigment ratio
- Lipoxidase activity
- Gluten quality
- Milling efficiency

Güleç et al., (2010) stated the factors that determine the quality of wheat as per this list.

4.2.1. Factors that affect the quality of wheat

The three main factors that affect the quality of the wheat production described by (Denčić et al., 2012; Drezner et al., 2007; Gooding and Davies, 1997; Peterson et al., 1992) are listed below. However, the list might be extended depending on producer, manufacturer or end-users like transporters, storage facilities, millers, et al.

- Genotype
- Seed quality
- Environmental factors

4.2.1.1. Genotype

Wheat (*Triticum aestivum* L.) quality may be improved by breeding elite varieties, improving crop/farming management practices and exploiting the synergism between genotype and the environment (Awulachew, 2019; Gaju et al., 2011; Peterson et al., 1992). Nass et al. (1976) tried different nitrogen doses in six bread wheat cultivars. They reported that the most important


factor in providing yield increase with nitrogen application is the variety feature and while a significant increase in grain yield can be achieved with nitrogen fertilizer application in high-yielding varieties, there is little or no yield increase in medium and low-yielding varieties. It is known that certain quality parameters are highly influenced by genetic factors (i.e., hardness is clearly genetically determined) (Vázquez et al. 2012; Carson and Edwards, 2009; Wrigley, 2007). Although genotype is an important factor, the determining factor in the effectiveness of genotype is Genotype x environment interaction (Šramková et al., 2009; Johansson et al., 2020). Varieties with genotypically high yield and quality characteristics can show their yield and quality characteristics if they are grown in suitable ecological conditions and appropriate cultural practices are performed (Anonymous, 2012). A good genotype takes high amounts of nitrogen from soil and fertilizers and uses it to increase grain yield rather than stems. The efficiency of nitrogen use for crop production in low-input sustainable agriculture can increase yields, reduce costs, and reduce the harmful effects of excess fertilizer on the environment (Arpacioğlu., 2018). Šekularac et. al., (2018) research shows that genotype was the most significant factor on the gluten index variation.

4.2.1.2. Seed quality

The major seed quality characteristics are physical quality, genetic purity, physiological quality and seed health. The health status of seeds is judged by the presence or the absence of insect infestation and seed-borne diseases. Seed-borne refers to the particular plant diseases that are transmitted by seeds. Using seeds which are free from disease inoculum is a primary means of reducing the introduction of pathogens into fields; it also reduces inoculum production and the speed of secondary spread into the inoculum threshold and disease severity after it's diagnosed. The prime factor in guaranteeing an increase in productivity of at least 10–15% is seed quality. To achieve this quality level, all the production factors that will affect seed viability, guarantee genetic purity and lack of disease should be taken into account. The success of modern agriculture depends on using pathogen-free seeds with high yielding character, which, in turn, improves disease management (Gaur et al., 2020).

Seed quality is defined via standards of excellence in certain characteristics and/or attributes that will determine the performance of the seed when sown or stored (Banu et. al. (2004) and


effect on the performance of the quality and quantity of the wheat production as well, as shown in Figure 3.




3. *Figure Effect of seed quality on grain yield, Kumar. D, and Seth. R, (2004)*

High quality seed is important to ensure maximum seed germination and seedling vigor, which in turn is instrumental in achieving maximum yield. Poorer quality seeds show low viability, reduced germination and emergence rates, poor tolerance to sub-optimal conditions and low seedling growth rates as shown in Figure 4 and Figure 5. The seed quality is also reflected in the final growth, maturity of plants, their uniformity and stability of yield. Seed size is an important physical indicator of seed related to yield, market grade factors and harvest efficiency (Gadisa, 2019).

The concentrations of P and K in the seed were significantly associated with grain yield where source was a significant factor and it affected the yield up to 8% in the research of McDonald and Hussein (2017); however, researchers have found that using larger size seeds resulted in a 4-5% yield increase as well.



4. Figure Effect of seed size on seedling vigour in wheat *Simmons et. al., (2000)*



5. Figure Effect of seed size on germination percentage *Hossein et. al., (2011)*

4.2.1.3. Environmental factors

The expression of the genetically determined good properties are influenced by environmental factors (Horváth, 2016). A study done with 12 varieties at 10 different locations by Miezán et al., (1977) reveal that protein content was affected by both the experiment site and nitrogen fertilization; however, Bayfield (1936) found that protein content in wheat increased as soil texture became heavier and as the soil fertility increased. Some ecological factors, including soil physiological and chemical properties and geographic latitude, can also affect wheat quality (Awulachew, 2019). Another experiment by Terman (1979) held in pots shows that nitrogen and moisture supply, light, temperature, and other growth factors greatly affect yield-protein relationships among cultivars. Naturally, weather impacts may have direct or indirect influence

on the performance of agricultural production and the food industry. Weather and climate remain the major uncontrollable driving forces in agricultural production; despite advances in technology and crop varieties, those factors still present more constraints on yield and affect production management practices from seedbed preparation to harvest than do soil management and agronomic practices (Decker, 1994; Olesen et al., 2000). An experiment in northeast Austria showed that a warming of 2°C in the air temperature would shorten the crop-growing period by up to 20 days and would decrease the potential winter wheat yield on nearly all of the soil types in the region (Thaler et al., 2012), as well as Lv, Z. et al. (2013) found that under rain-fed conditions, wheat yield is reduced in the north regions of China in three future periods, while wheat yield increases in the south regions of China.

Future climate scenarios may be beneficial for wheat in some regions, but could reduce productivity in zones where optimal temperatures already exist (Ortiz et al., 2008).

Water availability can be considered a basic factor related to yield quality and quantity performance of grain crops (Jolánkai et al., 2016). If cultivars that provide genotypically high quality products are not grown in suitable environments and care is not taken with the cultivation techniques, they do not reflect the desired quality characteristics.

It should be taken into consideration that every single environmental unit has effects on crop production -- e.g., hedgerows which reduce the wind speed. Those have been predicted to have particularly positive effects on winter wheat production (Thaler et al., 2012); however, results showed that N supply was the topmost environmental factor affecting protein content and composition (Triboi et al., 2000; Johansson et al., 2001).

4.3. Ecological requirements of the winter wheat

Wheat is an annual plant and is grown almost all over the world because it has many varieties that can be grown in all climates and soil conditions.


Wheat is generally grown in warm and cool climate conditions. Wheat does not like high temperatures in the early stages of development (germination, tillering). If the average humidity is above 60%, the plant continues to develop normally. It does not require much heat in the advanced stage of vegetative growth. A temperature of 10-15°C with a relative humidity of 66% and little light are suitable ingredients for decent growth. Wheat grows best in temperatures between 21°C/70°F and 24°C/75° F. The minimum temperature that wheat can handle during its growth cycle is about 4°C/ 40°F. Wheat does not grow well if temperatures exceed 35°C/95°

F. Wheat will grow optimally in a deep, fertile, well-irrigated and well-aerated soil at a pH between 5.5 and 7.5 (Nasa (n.d.)). Wheat can be grown in climatic regions with an annual precipitation of 350-1150 mm. A good quality and abundant product can still be produced in places with annual precipitation of 500-600 mm or in irrigations that will provide this moisture in the soil.

Sandy loam soils with deep, clayey, loamy-clay and sufficient organic matter, phosphorus and lime, guarantee the best wheat. The higher the organic matter in the soil, the higher the wheat yield. (T.C. Tarım ve Orman Bakanlığı (n.d.)).


4.4. The grain protein

Wheat germ is well-known high-quality nutritional product. Wheat germ has 10.80% water, 26.50% crude protein, 8.56% crude fat and 4.18% ash content on a dry weight basis. Wheat germ contains lipids (10%-15%), proteins (26%-35%), sugars (17%), fiber (1.5%-4.5%) and minerals (4%). It also contains substantial amount of biologically active compounds like tocopherols, phytosterols, policosanols, carotenoids, thiamin and riboflavin as 300-740 mg / kg, 24-50 mg / kg, 10 mg / kg, 4-18 mg / kg, 15-23 mg / kg and 6-10 mg / kg, respectively (Mughal., 2019). Protein is the primary quality component of cereal grains. Protein concentration is influenced by both environmental and genotypic factors that are difficult to separate (Fowler et al., 1990). Wheat grain proteins are classically categorized and separated according to solubility in albumin, globulin, gliadin and glutenin (Osborne, 1907). Proteins make up 7–22% of wheat's dry weight (Godfrey et al., 2010); according to some sources, the protein content in wheat grain can be up to 25% (Gooding and Davies, 1997).



6. Figure Wheat gluten protein and its impacts on wheat processing quality. Ma et al., (2019). *Frontiers of Agricultural Science and Engineering*.

One study stated that the protein ratio is the most important quality criterion determining the quality of wheat, and it determines the water absorption, stability, resistance and elasticity values of the flour (Bilgin and Korkut, 2005). In more recent times, gliadin and glutenin make up between 60-80% of total grain protein (Langenkämper and Zörb, 2019). Gluten is a large family of proteins, and accounts for up to 80% of the total protein content. It is responsible for the unique elasticity and stickiness of wheat dough -- the properties that make it so useful in bread making (Arnarson, 2019). Eser et al., (2017) mention (and show in Figure 7) that the results obtained suggest that a strong correlation was detected between the total amount of protein and the experimental treatments, regardless of the impact of crop years.



7. Figure Total protein yields in the non-favorable crop year. Nagygyombos 2014. Eser et. al. (2017)

In comparative studies with varieties developed in different breeding periods, it is stated that modern varieties provide a yield advantage of at least 50% compared to the improved varieties, but it is quite difficult to improve the yield and grain protein percentages together with the variety breeding methods due to the negative relationship between these two (Slafer et al., 1994). Similarly, Van Lill and Purchase (1995) mention that while high-yielding new varieties were obtained with the breeding studies for yield, the progress in terms of grain protein ratio and bread quality in the obtained varieties remained at a certain level. Research has shown that one of the most important cultivation techniques -- fertilization applied at the right time and in the appropriate amount -- can increase the quality of products that can satisfy the demands of the producers, consumers and industrialists.

4.3.1. Quality of the protein


Protein has two components associated with it, namely, the quantity or protein content and the quality of this protein. Both are affected by the levels of nitrogen and sulphur in the plant as it develops through the season (Yara, n.d.). More valuable nutrients for the nutrition of humans and animals are found in the outer parts of the wheat grain, especially in the aleurone layer. Proteins in the aleurone layer (albumin and globulin) constitute approximately 20% of the total protein in the grain and have different structural properties from gluten proteins. The aleurone layer's proteins do not have viscoelastic properties and thus do not take part in dough formation, but are important for physiological nutrition (Ereku et. al., 2016).

Gluten material forms the skeleton of the dough and holds in the gases created by the yeasts, allowing the bread to retain a shape. Protein quality is related to gluten quantity and characteristics -- although the quality of the protein of wheat is affected by agronomical applications, and it depends on its inherent variety as well. The elongation and its resistance to it, shape, elasticity, gas holding power, and capacity of the dough all depend on these properties. When it comes to protein quality, it is there to provide the essential amino acids that are important in human and animal nutrition; but the suitability of the physical properties of the protein is for bread making, rather than the nutritional and biological value of the protein. Whether the protein quality is high or not is determined by the sedimentation test performed on the flour. A high sedimentation value indicates that the quality is high and the volume of breads made from such flour is also high. In this respect, we should take into account not only the protein amount but also the protein quality in the flour to be baked; in other words, the quality difference in the amount of protein in breads made from the same flour arises from the protein quality (Ertugay, 1982). Apart from the amount of protein for wheat, the desired quality of these proteins is also a necessary criterion (Edwards et. al., 2007). Some researchers explain that the low sedimentation value is an important quality parameter, as high temperatures (>30°C) in the grain filling period increase the amount of gliadin (Panozzo and Eagles 2000). As a result, we can say the following: the amount and quality of protein in the flour are the two most important factors that determine the highest usefulness value in the flour.

4.4. The baking quality

The effects of nitrogen fertilization are clearly seen on gluten and gluten index values, which are important criteria of bread quality in sedimentation measurements made on the protein amounts. The proper arrangement of protein webs is effective for a good flour yield as well as warding off pest damage (Todab, n.d.).

Baking quality of wheat flour is determined by grain protein concentration (GPC) and its composition. It's also highly influenced by environmental factors, such as nitrogen (N) fertilization management (Xue et al., 2019). Increasing the level of protein content will raise the baking quality (Gabriel, et al., 2017; Wrigley and Batey, 2003).



8. Figure Remix loaf volume results for flour samples of two Canadian genotypes (Manitou and the unnamed line 11-463A), with a range of protein contents. Baking quality is indicated as loaf volume (L.V.) and protein content is expressed on the basis of 14% moisture basis (m.b.). Bushuk et al. (1969).

Studies in the Bushuk et al., (1969) (Figure 8) indicate the strong correlation between protein content of the grain and the baking quality.

Besides bread making quality, the flour milling performance of wheat is also of considerable technological importance (Kent, 1984). Much less is known about the factors that affect intracultivar variation in milling quality than those that affect bread-making quality, but starch damage level which, in turn, affects water absorption and dough rheology, is another aspect. A minor protein fraction associated with the starch granule surface has been implicated as having a role in controlling grain endosperm texture (hardness), an important component of milling quality (Schofield and Greenwell, 1987).



9. Figure Baking quality of wheat is determined by wheat cultivar, the interplay of natural grain constituents' proteins, carbohydrates and lipids, growing conditions of wheat as well as milling and baking technology. Langenkämper et al.(2019)

Quality bread production depends mainly on the raw material from which the bread is made, as well as the processing capacity of factories and bakeries, the methods used in the enterprise, the equipment, and the state of the equipment. A quality bread requires a quality wheat that provides quality flour. In this respect, the wheat and flour quality and the factors affecting the quality should be examined separately (Ertugay, 1982).

4.5. The nitrogen fertilizer

The nitrogenous fertilizer industry also produces synthetic ammonia, nitric acid, ammonium nitrate, and urea. Synthetic ammonia and nitric acid are used primarily as intermediates in the production of ammonium nitrate and urea fertilizers. The following is a list of primary specific products classified as nitrogenous fertilizers (Khan et al., 2018):

- ammonia liquor
- ammonium nitrate
- ammonium sulfate
- anhydrous ammonia
- aqua ammonia
- fertilizers, mixed, produced in nitrogenous fertilizer plants
- fertilizers, natural
- nitric acid

Nitrogenous fertilizer application is one of the most critical factors that can increase the quality of wheat. The dose and time of application of nitrogenous fertilizers are also important in terms of positively affecting the quality characteristics of wheat. The nitrogen fertilizer, as applied during the growing period and especially during tillering, plays an important role in the quality of the grain (Hoeser and Schafer, 1969).

4.5.1. Effect of nitrogen on quality, yield and development

Among the nutrients used in wheat nutrition, nitrogen is the greatest nutritional element that affects the yield and quality of wheat (Başar et al., 1998; Arpacioğlu., 2018), and its importance is constantly emphasized (Gallagher et al., 1983; Gauer et al., 1992). Nitrogen is often the most limiting factor in crop production. Hence, application of fertilizer nitrogen results in higher biomass yields and protein yield; additionally, concentration in plant tissue is commonly increased. Nitrogen often affects amino acid composition of protein and, in turn, its nutritional quality, says Blumenthal et al., (2008); moreover, the increasing nitrogen supply generally

improves kernel integrity and strength, resulting in better milling properties of the grain. In Sohail et al., (2018), a study to determine the effect of different nitrogen fertilizer application methods on yield and quality in wheat, nitrogen fertilization (12 kg/da of pure nitrogen) was applied separately in wheat sowing, tillering, stem lift and grain filling development periods and reached the highest yield with 506.0 kg/da and 14.9% protein value. It stated that the grain protein ratio was lower in the nitrogen fertilizer application applied when the planting had been done. The study concluded that the nitrogen fertilizer applied in the early periods increased the vegetative development and tillering potential, the nitrogen fertilizer applied in the tillering and booting-heading periods increased the grain protein and gluten content, and the importance of nitrogen in terms of quality was emphasized. Reddcliffe et al., (2000) showed results in their study in which they used the divided N application as 0 - 175 kg N/ha at the stage of tillering and 0-40-80 kg N/ha at the flag leaf stage in New Zealand, showing that grain yield and protein ratio increased with increasing nitrogen fertilizer amount.

4.5.2. Other effects of nitrogen fertilizer

Nitrogenous fertilizer application increased the protein amount in the flour in parallel with the increase in the protein amount in the grain, and this increase was found to be statistically significant (Horváth et al., 2014; Ercan and Bildik, 1993). Studies have also determined that protein ratio, gluten amount and sedimentation value increase linearly with the increase of nitrogen fertilizer application (Paredes et al., 1985).

Nitrogenous fertilizer application in dry farming not only increases the yield in winter wheat but also increases the use of water (Xu et al., 2020). Fertilized plants can also use water from the lower layers of the soil. While nitrogen-free winter wheat achieves its water intake from the 90 cm profile, it uses 180 cm moisture profile when nitrogen is given (Hagin and Tucker, 1982). By promoting plant growth with application of nitrogenous fertilizer, a faster and deeper root system is developed, and plants can benefit from deeper water. As the nitrogen dose increases, the efficiency increases and in parallel with this, the efficiency of using water -- which then increases up to 35%. In the following studies, the difference between the fertilizers in terms of the protein content of the grain was found to be significant, and ammonium nitrate fertilizer gave better results than others in this respect (Yılmaz and Şimşek 2013; Başar et al., 1998; Akkaya A. 1994). Soil or foliar Fe fertilizers had little or no effect on the green parts and grain Fe content; however, increasing N applied in any Fe fertilization strongly increased the green

portion and grain Fe content. The results show that improving the N nutritional status of plants through fertilization is an important agricultural practice in increasing the grain Fe content and resulting improvement in human health (Özden, 2012). The effects of different nitrogen (N) applications on the Zn uptake and accumulation of corn and wheat were investigated by Rehman (2019). As N applied to wheat and maize increased, more Zn accumulated in green portions and the uptake of foliar applied Zn increased significantly.

4.5.3. Negative effects of nitrogen fertilizer

The negative effects of nitrogen fertilizer have great importance -- in terms of the input costs of production. Thus, the high costs of the production of nitrogen fertilizers, the high energy requirement, the excess nitrogen fertilizer used, and the economic as well as environmental impacts should be considered (Atar et al., (2017); Wang et al., (2017); Gündoğmuş, (1998); Abad et al., (1996). However, it has been shown that increasing the nitrogen doses after a certain level (i.e., overdosing) does not provide the desired increases in yield (Johnson et al., 1973); Styk ve Dziamba, (1984); Eser et al., (1999)). In summation, the nitrogen fertilization is effective and good for winter wheat production, but due to side effects such as cost and environmental harms, it should not be overdosed.

Burkart and Stoner, (2008) and Hamlin, (2006) mention that once a shallow aquifer has been contaminated by nitrate, it may take decades for the groundwater quality to improve even after pollution controls have been implemented. Therefore, in the quest for increasing agricultural production, public health should also be considered a critical factor.

It's generally accepted that fertilization increases soil microbial biomass in agricultural soils on the short and long term (Nguyen et al., 2018), however, application of solely continuous N fertilizer can decrease soil microbial biomass (Wallenstein et al., 2006; Treseder, 2008). Additionally, it may even alter the soil microbial stoichiometry and diversity via the nitrogen (Wang et al., 2018a, b). Osmotic stress conditions and C substrate deficiency caused by the extreme amounts of applied nitrogen also suggest that nitrogen fertilizer should never be overdosed (Yevdokimov et al., 2008).

4.5.4. Efficiency of the split dose application of nitrogen fertilizer

Split fertilizer N application has been proposed for improving N uptake efficiency in wheat (*Triticum aestivum L.*) production systems (Belete et al., 2018; Rahman et al., 2011; Jia et al., 1996; Alcoz et al., 1993). Studies have demonstrated that the split N application under unfavorable conditions for cultivation result in higher N use by the plant, reducing its losses to the environment with a yield increase (Brezolin et al., 2016; Ferrari et al., 2016). The timing of its application is one of the most important factors on what effects are produced (Gooding and Davies, 1992). The availability of nitrogen for the wheat during various phases of its growth and development is another important factor influencing the yield and quality of grain (Zende et al., 2005). Application of N fertilizer to wheat, preferably as a top dressing between tillering and stem elongation, is a strategy to be recommended from the standpoint of both the environment and farmer returns, says Lopez-Bellido et al., (2005). Their research affirms the mean wheat use of N fertilizer ranging from 14.1% when applied at sowing, to 54.8% as applied as a top dressing at the beginning of stem elongation. Delayed nitrogen fertilization had no significant increase on the percentage of the protein (Lloveras et al., 2001). Split dose application of the Nitrogen fertilizer has positive effects on most of the crops such as rice production (Sathiya and Ramesh, 2009; Shaiful et al., 2012; Djaman et al., 2018), cotton production (Hallikeri et al., 2010), Brassica species (Ahmad et al., 1999), maize production (Joshi et al., 2014; Hammad et al., 2011), potato production (Ayyub et al., 2019; Al-Moshileh et al., 2005), poppy production (Lošák and Richter, 2004), soybean production (Singh and Singh, 2013a), sunflower production (Khazada et al., 2016), barley production (Singh et al., 2013b), oats production (Alipatra et al., 2013), tomato production (Singandhupe et al., 2003), sugarcane production (Koochekzadeh et al., 2009), and others.


The results of Madan and Munjal's (2009) research showed that splitting the recommended dose of nitrogen recorded higher total protein content (12.68%) as compared to control (10.23%), and splitting of recommended dose of nitrogen increased the grain yield by 4.1%. The investigation by Xue et al., (2016) demonstrates the same thing with an application of nitrate and the urea form of the nitrogen fertilizer. Nitrogen applied at the late growth stages (boot stage or heading) had considerable increase on baking quality and nitrate, showing a greater effect than the urea form. The results of Rahman et al., (2011) research shows that plant N uptake was significantly influenced by N rate and N splitting, and also the interaction of N rate and N splitting. Total N uptake was maximum under N rate of 120 kg/ha applied as three

equal splits as 1/3rd basal with 1/3rd as top dress at CR1 plus 1/3rd as top dress at 1st node stage. Late-stage growth with N fertilization increased the loaf volume of wheat flour by raising grain protein concentration and altering its composition. However, splitting the N changed grain protein composition by enhancing the percentages of gliadin and glutenin as well as certain high molecular weight glutenin subunits (HMW-GS), which led to an improved baking quality of wheat flour. Thus, splitting nitrogen application had greater results on baking quality. What should be considered as well is the wheat's N dose and time of application, which is determined by the amount of organic matter in the soil, the previous crop, and the anticipated yield.

4.5.5. Amount of the nitrogen application

Numerous studies stipulate that overdose application, rather than the appropriate amount of nitrogen fertilizer application, had no significant positive effect on wheat quality parameters (Abedi et al., 2011; Cui et al., 2010; Chen et al., 2004; Lloveras et al., 2001) and even dropped the yield performances (Jolánkai, 1985). Nitrogen use efficiency is closely related to fertilization, the genetic and morphological characteristics of the plant, as well as the climate and soil characteristics.

Effect of applied nitrogen on winter wheat grain protein



10. Figure Effect of applied nitrogen on winter wheat grain protein. Yara, (2011)

The research of Khalilzadeh et al., (2012) found that there was high variation among genotypes in terms of nitrogen utilization efficiency.

Following the research of Mandic et al., (2015) they advised that the use of large amounts of N increases production costs and reduces the economic benefits. The findings of Liu et al., (2016) reveal that the recommended N rates reduced the residual inorganic N, nitrate leaching, and direct nitrous oxide emissions by 8–27%, 29–52%, and 19–36% in the three regions, respectively. These findings suggest that this N recommendation method provides an option to balance the yield, grain quality, income, nitrogen use efficiency, and environmental impacts of winter wheat production. It should be noted that the optimal nitrogen requirements of each species differ in different environmental conditions and different agricultural applications. Gelinas et al., (2009) and Öztürk and Gökkuş (2008) expressed the effect of the environmental factors on the quality parameters of winter wheat. Arpacioğlu (2018) research ran five different nitrogen dose applications (0-50-100-150-200 kg N/ha) on wheat; the trials showed that plant height, spike length, grain yield, thousand grain weight, gluten index and nitrogen uptake efficiency increased to about 150 kg/ha nitrogen application and decreased after this dose.

While the maximum unit area yield in grain yield was obtained from the application of nitrogen dose of 150 kg/ha, this value was determined as 153 kg N/ha in the economic analysis. Çetin et al., (1999) suggested 140 kg N/ha dose of nitrogen fertilizer base on their study where the 0-60-120-240 kg N/ha doses were tested. Researchers found positive correlation between protein content of the grain and the increasing level of N application; however, protein content of the tested grains varied between 10.1-19.4%. The trials of Zecevic et al., (2010) with 60-90-120 kg N/ha results showed that 120 kg N/ha application gave the best sedimentation value, while Hussein et al., (2006) trials with 0-50-100-150-200 kg N/ha application showed that even the grain yield and biological yield were statistically similar at doses of 150 kg N/ha and 200 kg N/ha. However, a dose of 200 kg N/ha, compared to a dose of 150 Kg N/ha, significantly increased the protein content. Similarly, the results of Yousaf et al., (2014) with 0-80-100-120-150 kg N/ha applications showed that 120 and 150 kg N/ha achieved the highest plant height, a greater number of tillers, the maximum number of fertile tillers, and highest yield respectively, as compared to the control and other treatments. However, 120 kg N/ha is still recommended for maximum growth and wheat yield.

Similar to above, Mantai et al., (2015) with 0-30-60-120 kg N/ha application on oats; Grabowska and Kozlara (2006) with 40-80-120 kg N/ha application on hemp; Biesiada and Kus (2010) with 50-150-250 kg N/ha application on sweet basil; Djaman et al., (2018) with 0-60-90-120-150 kg N/ha application on rice; Mut et al., (2005) with 0-60-120-180 kg N/ha application on triticale; Szulc et al., (2022) with 150 and 150-soil content kg N/ha application on maize; Singh et al., (2019) with 70-100-130 kg N/ha application on canola, cereals and many more other doses and plants, showed similar results.

4.6. The grain moisture

The moisture content of wheat is a relevant issue for harvest, transport, storage and even the milling properties and the required energy for milling. Importantly, lowering the grain moisture led to about a twofold decrease in the required milling energy (Hassoon et. al., (2021) & Jung et. al., (2018)). Excess water in wheat reduces the commercial value as it causes a decrease in dry matter, and it makes storage difficult as it encourages germination as a result of bacterial and fungal activity. Factors such as growing needs, storage conditions, and harvest time affect the water rate in wheat. Specifically, the grain moisture content is higher in the product years

that receive more precipitation during the maturity period of wheat (Bulut, 2012). Worldwide, most grain farming is carried out in climatic regions that are not suitable for storage because of temperature and humidity values. Grains stored under high temperature and relatively high humidity conditions also deteriorate. In addition to the losses in the quantity and quality of the grain due to these deteriorations, risks are posed to consumer health. In addition, the use of spoiled grains in animal nutrition indirectly poses risks to human health (Posner and Hibbs, 2005; Brooker et al., 1992). The upper limit for moisture content in wheat is 14.6% for storing the wheat (Ünal, 2002). Moisture content is also used for determining the most suitable harvest time (Ankara, 2015). What is the impact of moisture on end-product quality? High-moisture grain risks becoming infested with insects, mold and other unwanted organisms. In extreme cases, this can lead to toxins that end up in flour to the detriment of end-product quality. Conversely, too low a moisture level is also a problem: grain with less than nine per cent moisture requires additional processing to achieve the required conditioning level. High moisture content may be due to early harvesting, before the grain has dried below 12.5 per cent, or wet conditions around harvest time. Often, early harvesting is practiced avoiding wet weather. There are steps that can be taken to reduce the moisture content of harvested grain, including drying before storage, blending, or drying in storage using silos with aeration systems. It is important not to store grain with high moisture content and, especially, to avoid storage of high-moisture grain in silo bags (Qail 2018).

5. MATERIALS AND METHODS

5.1. Field features of experiments

In these three-year sets: 2015-2016, 2016-2017 and 2018-2019, growing seasons for a field trial of high milling and baking quality winter wheat (*Triticum aestivum* L.) varieties were set up under identical agronomic conditions using split-plot design (10 m²/plot). The trials were established at the experimental fields of the Hungarian University of Agriculture and Life Sciences, Crop Production Institute, Hungary in two different sites. The 2015-2016 and 2016 - 2017 crop seasons at Nagygyombos-Hungary was the experimental site of Hungarian University of Agriculture and Life Sciences, Crop Production Institute. This site in Nagygyombos lies between latitude 47°40'53.00 "N and 47°41'90.00", longitude 19°40'30.00"E and 19°40'20.00"E, shown in Figure 11 and soil parameters of the experimental field on Table 3.



11. Figure The experimental field view by satellite at 2016-2017, Nagygyombos, Hungary

Soil type of the trial site at Nagygyombos was chernozem (calciustoll).

| | Humus % | pH (H ₂ O) | K _A | Sand % | Silt % | Clay % | CaCO 3 |
|--------|---------|--------------------------|----------------|--------|-----------|-----------|-----------|
| Medium | 2.65 | 7.30 | 45 | 49 | 25 | 26 | 1.86 |

3. Table Soil type of the experimental field at Hungarian University of Agriculture and Life Sciences, Crop Production Institute, Nagygyombos, Hungary

- Organic matter content %: 2,65
- CaCO₃ %: 1,86
- pH (KCl): 7,30
- K_A: 45
- P₂O₅ (mg/kg): 643
- K₂O (mg/kg): 293

The is the Gödöllő-Hungary experimental site of the Hungarian University of Agriculture and Life Sciences, Crop Production Institute, used for 2018–2019 and 2019–2020 crop seasons. The experimental site in Gödöllő lies between latitude 47°59'39.88"N and 47°59'60.54"N, longitude 19°37'13.26" and 19°36'82.81", shown in Figure 12.



12. Figure The experimental field view by satellite at 2019-2020, Gödöllő, Hungary

The soil type of the experimental field was sand-based brown forest soil (Chromic Luvisol). The textural classification of the soil was sandy loam with parameters shown in Table 4. The agronomic characteristic of the soil was neutral sandy soil with variable clay content. The soil structure was susceptible to compaction issues. The water retention characteristics were poor due to the high sand content. The soil was exposed to the impacts of drought.

| | Humus % | pH (H ₂ O) | KA | Sand % | Silt % | Clay % | CaCO ₃ |
|--------|---------|-----------------------|----|--------|--------|--------|-------------------|
| Medium | 1.32 | 7.08 | 40 | 49 | 25 | 26 | 0 |

4. Table Soil type of the experimental field at Hungarian University of Agriculture and Life Sciences, Crop Production Institute, Gödöllő, Hungary

5.2. Meteorological properties of the experimental fields


5.2.1 Hatvan-Nagygyombos meteorological properties

Figure 13 and Figure 14 provide information about the weather temperature, and Figure 15 and Figure 16 show rainfall amounts and rainy days of experimental site #1.

Hatvan

Max, Min and Average Temperature (°C)


Zoom 1m 3m 6m YTD 1y All



13. Figure Max, Min and Average Weather Temperature 2015-2016 (Nagygyombos, Hungary).
worldweatheronline.com

Hatvan

Max, Min and Average Temperature (°C)




14. Figure Max, Min and Average Weather Temperature 2016-2017 (Nagygyombos, Hungary). worldweatheronline.com

The experimental field #1's weather data showed differences between trial years: in 2016 summer the average temperature recorded approximately 3°C lower than the 2015 and 2017 weather average temperatures; however, in 2016 winter also was 6 °C colder than the 2015 and 2017 winter average.

Hatvan

Average Rainfall Amount (mm) and Rainy Days


Zoom 1m 3m 6m YTD 1y All



15. Figure Rainfall and Rain Days 2015-2016 (Nagygyombos, Hungary).
worldweatheronline.com

Hatvan

Average Rainfall Amount (mm) and Rainy Days



16. Figure Rainfall and Rain Days 2016-2017 (Nagygyombos, Hungary).
worldweatheronline.com

Recorded rainfall amounts show that experimental site #1 had 994,54 mm rainfall in 2015, 1.236,68 mm in 2016, and 1.176,47 mm. in 2015 -- which was drier than 2016 and 2017 crop years.


5.2.2. Gödöllő meteorological properties

Figure 17 provides information about the weather temperature, and Figure 18 shows the rainfall amount and rainy days of experimental site #2.

Godollo

Max, Min and Average Temperature (°C)

Zoom 1m 3m 6m YTD 1y All




17. Figure Max, Min and Average Weather Temperature 2018-2019 (Gödöllő, Hungary).
worldweatheronline.com

Summer and winter in site #2 was approximately 2°C warmer compared to site #1 on average.

Godollo

Average Rainfall Amount (mm) and Rainy Days



18. Figure Rainfall and Rain Days 2018-2019 (Gödöllő, Hungary). worldweatheronline.com

Site #2 received 906,64 mm rainfall in 2018 and 920 mm in 2019

5.3. Studied winter wheat varieties

The present study examined the performance of five high baking quality winter wheat varieties:

2015-2016 and 2016-2017 crop seasons in Nagygyombos - Hungary

-Alföld

-Mv Karéj

-Mv Nádor

-Mv Toldi

-Mv Toborzó

-Alföld

-Mv Karéj

| Variety | Alföld | Mv Karéj | Mv Nádor | Mv Toldi | Mv Toborzó |
|---|---------------|-----------------|-----------------|-----------------|-------------------|
| Ripening time | Early | Medium | Medium | Early | Early |
| Kernel type | Hard | Hard | Hard | Hard | Hard |
| Height (cm) | 75-95 | 80-90 | 60-80 | 80-90 | 80-90 |
| Yield (t/ha/) | 5,5-7,0 | 7,0-8,0 | 7,5-9,5 | 6,0-7,5 | 6,0-7,5 |
| TKW (gr) | 39-44 | 45-50 | 45-50 | 40-45 | 48-54 |
| Resistance (1-9) | 8 | 7 | 9 | 8 | 7 |
| Test weight (kg hl⁻¹) | 78-82 | 82-85 | 77-82 | 78-82 | 79-82 |
| Protein % | 14-17 | 11-13 | 12-14 | 13-15,5 | 13-15 |
| Gluten % | 34-40 | 29-32 | 28-32 | 30-36 | 32-36 |

5. Table Martonvásári Fajtakatalógus 2020

5.4. Treatments

The three-year experiment was set up in split-plot design with nine plot replications regarding each experimental factor such as variety and N application (time and dose) in each investigated year. The plots were sown and harvested with plot machines. Apart from N topdressing, all other agronomic treatments as well as sowing and harvesting were identically applied to all plots to study the impact of N treatments independently. N fertilizer topdressing was applied in single or split doses. N was applied in the form of ammonium nitrate (NH₄NO₃); the amounts indicate the N content in this paper, not the molecule. The applied fertilizer was granular ammonium nitrate with 34% content of the active ingredient. N was investigated in 6 different variants: 4 levels single and 2 levels split-dose treatments. Single application: 0, 80, 120, and

160 kg ha⁻¹ N, split-dose application: 80+40 kg ha⁻¹ and 120+40 kg ha⁻¹ in two applications. Single applications were done at the tillering stage, while split-dose treatment was applied at the stage of tillering and heading. There was no N application in autumn in any of the crop years.

5.5. Investigations

Grain yields of the winter wheat varieties were sampled and measured from each harvested plot. The protein, test weight, thousand grain weight, and baking quality parameters were measured from harvested wheat grain. Analyses were done at the research laboratory of the Hungarian University of Agriculture and Life Sciences, Crop Production Institute.

Dickey-john® Instalab® 600 Analyser, Near-infrared (NIR) spectroscopic equipment Mininfra Scan-T Plus 2.02 version (Arana, 2016) were used to measure gluten, protein, and Zeleny sedimentation values of whole grains. Falling Number was also studied to determine amylase enzyme activity in the flour. The Hagberg Falling Number (HFN) Perten Type:1400 system, which meets the requirements of the AACC (American Association of Cereal Chemists) No.56-81.04, ICC (International Cereal Chemists) No. 107/1 (2010), and PN EN ISO 3093:2010 standards, was used to determine the Falling Number. The OS 1 type equipment by the ISO 7971-3:2019 standard was used to measure test weight. Thousand grain weight and test weight were measured with the KERN EMS and the Sartorius MA-30 precision scales. Farinograph (Valorigraph) instrument had been used to describe baking quality of the dough.

5.6. Statistical analyses

For the statistical evaluation of the results, we used the Explore and ANOVA modules of the IBM SPSS V.23 software. For data analysis, part Independent Sample T Test, ANOVA, and Pearson Correlation Analysis were used to test the hypotheses of the study. The effect of the different treatments on seed germination was analyzed using one-way ANOVA at a 0,05 level of significance. Values are given as the mean ± standard deviation of four measurements. LSD (least significant difference) tests were used to determine the significant difference among the data. The statistical significance level was $p < 0,05$. That enabled us to determine the differences between the studied doses -- whether or not the obtained results had significant variations.

6. RESULTS

6.1. Test Weight Results

Table 6-7 and 8 as well as Figure 24-27 (Appendix 1) provide information on nitrogen application effects on test weight of the tested winter wheat varieties with the impact of undivided/split dose of N supply on two experimental sites during three years. The results obtained from experiment site #1 during first two years of the experiment shown on Table 6, test weight (kg hl^{-1}) values slightly decreased in some of the tested varieties by the increasing level of undivided N application, but the changes found were not significant. In addition, a positive effect of split-dose treatment had been detected, except in case of Mv Nádor 80+40 kg ha^{-1} to 120 kg ha^{-1} N application. The highest result had been recorded for the Alföld 80 kg ha^{-1} single dose application with 81.5 kg hl^{-1} at 2016-2017th experimental year and the lowest for MV Toborzó 80 kg ha^{-1} single dose application with 70.95 kg hl^{-1} at 2016-2017th experimental year. However, split dose N application did not present significant changes among the tested winter wheat varieties, similar results were reported by Pollhamer (1981) and Horváth and co-workers (2014).

| Test weight (kg hl^{-1}) | | | | | |
|-------------------------------------|--------|----------|----------|------------|----------|
| N topdressing | Alföld | MV Nádor | MV Karéj | MV Toborzó | MV Toldi |
| 0+ | 79.15 | 75.79 | 75.57 | 76.47 | 77.97 |
| 80+ | 79.17 | 74.95 | 75.68 | 76.19 | 77.29 |
| 80+40 | 78.86 | 75.13 | 77.03 | 76.49 | 77.50 |
| 120+ | 78.54 | 75.13 | 75.39 | 76.37 | 76.81 |
| 120+40 | 79.20 | 75.36 | 75.84 | 76.59 | 76.83 |
| 160+ | 78.26 | 74.90 | 75.57 | 76.26 | 76.79 |

6. Table Impact of N topdressing applications on wheat grain test weight. 2015-2016 and 2016-2017 (Nagyombos, Hungary)

The results obtained from experiment site #2 in the experiment shown on Table 7, were that the test weight (kg hl^{-1}) values slightly decreased in some of the tested varieties via the increasing level of undivided N application, but the changes found were not significant. In addition, a positive effect of split dose treatment was detected. The highest result recorded was for the

Alföld 120 kg ha⁻¹ single dose application and MV Karéj 80 kg ha⁻¹ single dose application with 74.55 kg hl⁻¹ at 2018-2019th experimental year, and the lowest was for MV Karéj 40 kg ha⁻¹ single dose application as 67.80 kg hl⁻¹ at 2018-2019th experimental year.

| Test weight (kg hl⁻¹) | | |
|---|---------------|-----------------|
| N topdressing | Alföld | MV Karéj |
| 0+ | 70,73 | 69,32 |
| 80+ | 72,88 | 71,10 |
| 80+40 | 73,12 | 71,55 |
| 120+ | 73,35 | 71,62 |
| 120+40 | 72,45 | 69,75 |
| 160+ | 70,80 | 69,23 |

7. Table Impact of N topdressing applications on wheat grain test weight. 2018–2019 (Gödöllő, Hungary)

Regarding the study employed during three crop years in two different experimental fields showed that maximum test weight was recorded in both sites was in the Alföld variety with single dose application of the Nitrogen fertilizer. Minimum test weights varied on different sites with different varieties, and the minimum test weights were recorded on single dose applications of Nitrogen fertilizer. Alföld and MV Karéj had higher numbers at Site 1 than Site 2: Alföld 9.19% and MV Karéj 7.70% were the higher values on site 1. Test weight showed significant differences between site, genotype and year; however, level and type of the treatment did not show significant effect from the one-way ANOVA statistical analysis shown on Table 36-54.

The significant level of the difference based on year, site, genotypes and treatment in means of test weight measured in the sample of all genotypes was examined through ANOVA and outcomes were reflected in Table 8.

| | Test Weight (hl) | N | M | SD | F/t | df | p | Difference |
|------|-------------------------|----------|----------|-----------|------------|-----------|----------|-------------------|
| Year | 2016 | 90 | 76.80 | 1.75 | 84.353 | 2-213 | .000 | 1, 2 > 3 |
| | 2017 | 90 | 76.52 | 2.57 | | | | |
| | 2019 | 36 | 71.33 | 2.52 | | | | |
| Site | Hatvan | 180 | 76.66 | 2.20 | 12.971 | 214 | .000 | 1 > 2 |

| | | | | | | | | |
|-----------|--------------|----|-------|------|--------|-------|------|-----------|
| | Gödöllő | 36 | 71.33 | 2.52 | | | | |
| Genotypes | Alföld | 54 | 76.63 | 3.39 | 10.596 | 4-211 | .000 | 1 > 2 > 3 |
| | Karéj | 54 | 73.91 | 3.42 | | | | |
| | Nador | 36 | 75.21 | 1.00 | | | | |
| | Toborzo | 36 | 76.39 | 2.09 | | | | |
| | Toldi | 36 | 77.19 | 2.36 | | | | |
| Treatment | 0 kg/ha | 36 | 75.83 | 3.32 | .242 | 5-210 | .943 | |
| | 80 kg/ha | 36 | 75.88 | 3.00 | | | | |
| | 80+40 kg/ha | 36 | 76.03 | 2.82 | | | | |
| | 120 kg/ha | 36 | 75.79 | 2.69 | | | | |
| | 120+40 kg/ha | 36 | 75.79 | 3.07 | | | | |
| | 160 kg/ha | 36 | 75.30 | 3.23 | | | | |

8. Table Comparison of the Means of Test Weight Parameters Measured in the Sample of All Genotypes Based on Categorical Variables, 2016, 2017 Hatvan and 2019 Gödöllő crop years.

As seen in Table 8, according to the outcomes it is found out that there is no significant difference in statistical level in test weight based on treatment ($F(5-210) = .242, p > .05$), however, there is a significant difference in statistical level in test weight based on year ($F(2-213) = 84.353, p < .001$), site ($t(214) = 12.971, p < .001$) and genotypes ($F(4-211) = 10.596, p < .001$) in the sample of all genotypes. Accordingly, the mean values of test weight in 2016 and 2017 tend to be higher than in 2019; Hatvan tends to be higher than in Gödöllő. In addition; Alföld and Karej tend to be higher than in Nador; Alföld tends to be higher than in Karej.

6.2. Thousand Kernel Weight Values

Table 9-10-11 as well as Figure 28-31 (Appendix 1) provide information on nitrogen application effects on thousand kernel weight varieties of the tested winter wheat, with the additional impact of an undivided/split-dose of N supply. The results obtained from experiment at site #1 during first two years of the experiment shown on Table 9, the thousand kernel weight value decreased slightly in most of the cases for the increasing undivided/split level of N applications: however, increasing number of N treatments had better effect in the comparison of 80+40 kg ha⁻¹ to 120 kg ha⁻¹ and 120+40 kg ha⁻¹ to 160 kg ha⁻¹, except for Mv Toborzó comparison of 120+40 kg ha⁻¹ to 160 kg ha⁻¹. Mv Nádor and Mv Toldi showed significant

differences via a one-way ANOVA test of thousand kernel weight. Similar results were reported by Szentpétery and co-workers (2005) and Horváth and co-workers (2014). The highest thousand grain weight was recorded on the untreated ($0 \text{ kg kg ha}^{-1} \text{ N}$) MV Toborzó plot with $51,90 \text{ g/}$ thousand kernel weight, and the lowest was detected on an Alföld plot treated with undivided $160 \text{ kg ha}^{-1} \text{ N}$ application resulting 31.40 g/ thousand kernel weight.

| Thousand kernel weight (g/1000 kernel) | | | | | |
|---|---------------|-----------------|-----------------|-------------------|-----------------|
| N topdressing | Alföld | MV Nádor | MV Karéj | MV Toborzó | MV Toldi |
| 0 | 39.73 | 44.78 | 46.70 | 45.05 | 45.81 |
| 80 | 38.73 | 43.16 | 45.58 | 42.64 | 44.20 |
| 80+40 | 38.63 | 42.79 | 45.82 | 42.42 | 45.07 |
| 120 | 38.56 | 42.10 | 45.40 | 41.44 | 43.77 |
| 120+40 | 38.83 | 42.21 | 45.36 | 42.40 | 44.40 |
| 160 | 38.02 | 41.09 | 43.78 | 42.60 | 42.88 |

9. Table Impact of N topdressing applications on thousand kernel weight of wheat varieties. 2015-2016 and 2016–2017 (Nagygyombos, Hungary)

The results obtained from experiment on site #2 during the 2018-2019 experiment shown on Table 10, the thousand kernel weight value increased up to $120 \text{ kg/ha}^{-1} \text{ N}$ undivided, and split application accordingly with a rising level of Nitrogen amount. However, after $80+40 \text{ kg/ha}^{-1}$ and $120 \text{ kg/ha}^{-1} \text{ N}$ applications, increasing level of the N supply had a slightly negative effect on the thousand kernel weight at the $120+40 \text{ kg/ha}^{-1}$ and $160 \text{ kg/ha}^{-1} \text{ N}$ applications compared to the $80+40 \text{ kg/ha}^{-1}$ and 120 kg/ha^{-1} . The highest thousand grain weight was recorded on the $80 \text{ kg kg ha}^{-1} \text{ N}$ MV Karéj plot with 45.08 g/ thousand kernel weight, and the lowest was detected on an Alföld plot treated with undivided $80 \text{ kg ha}^{-1} \text{ N}$ application, resulting in 38.00 g/ thousand kernel weight.

| Thousand kernel weight (g/1000 kernel) | | |
|---|---------------|-----------------|
| N topdressing | Alföld | MV Karéj |
| 0 | 38,75 | 40,99 |

| | | |
|---------------|-------|-------|
| 80 | 39,43 | 42,22 |
| 80+40 | 40,07 | 42,45 |
| 120 | 40,53 | 44,24 |
| 120+40 | 40,37 | 42,21 |
| 160 | 40,07 | 41,46 |

10. Table Impact of N topdressing applications on thousand kernel weight of wheat varieties. 2018-2019 (Gödöllő, Hungary) 1

Comparing the two sites, Alföld had 2.89% higher numbers at site #2 than site #1, and the MV Karéj had 7.52% higher numbers at site #1 than site #2, however, in general comparison the sites did not have significant differences between each other. Thousand kernel weight showed a significant effect on year and genotype, but no correlation was found with a different dose and type of N applications, as tracked on Table 36-54.

The significance level of the difference, based on year, site, genotypes, and treatment in terms of thousand kernel weight measured in the sample of all genotypes, was examined through ANOVA and outcomes were reflected in Table 11.


| Thousand Kernel Weight (g) | | N | M | SD | F/t | df | p | Difference |
|-----------------------------------|--------------|----------|----------|-----------|------------|-----------|----------|-------------------|
| Year | 2016 | 90 | 44.53 | 2.26 | 30.280 | 2-213 | .000 | 1 > 2, 3 |
| | 2017 | 90 | 41.14 | 4.15 | | | | |
| | 2019 | 36 | 41.07 | 2.07 | | | | |
| Site | Hatvan | 180 | 42.84 | 3.74 | 2.752 | 214 | .006 | 1 > 2 |
| | Gödöllő | 36 | 41.07 | 2.07 | | | | |
| Genotypes | Alföld | 54 | 39.12 | 2.60 | 28.322 | 4-211 | .000 | 2 > 3 > 1 |
| | Karéj | 54 | 44.51 | 2.45 | | | | |
| | Nádor | 36 | 42.69 | 2.15 | | | | |
| | Toborzó | 36 | 42.76 | 4.73 | | | | |
| | Toldi | 36 | 44.35 | 2.19 | | | | |
| Treatment | 0 kg/ha | 36 | 43.66 | 3.50 | 1.154 | 5-210 | .333 | |
| | 80 kg/ha | 36 | 42.52 | 3.43 | | | | |
| | 80+40 kg/ha | 36 | 42.67 | 3.64 | | | | |
| | 120 kg/ha | 36 | 42.28 | 3.58 | | | | |
| | 120+40 kg/ha | 36 | 42.41 | 3.59 | | | | |
| | 160 kg/ha | 36 | 41.71 | 3.67 | | | | |

11. Table Comparison of the Means of Thousand Kernel Weight Parameters Measured in the Sample of All Genotypes Based on Categorical Variables, 2016,2017 Hatvan and 2019 Gödöllő crop years.

As seen in Table 11, the outcomes reveal that there is no significant difference in statistical level in thousand kernel weight based on treatment ($F(5-210) = 1.154, p > .05$), however, there is a significant difference in statistical level in thousand kernel weight based on year ($F(2-213) = 30.280, p < .001$) and site ($t(214) = 2.752, p < .01$), genotypes ($F(4-211) = 28.322, p < .001$) in the sample of all genotypes. Accordingly, the mean values of thousand kernel weight in 2016 tend to be higher than in 2017 and 2019; Hatvan tends to be higher than Gödöllő; Karéj tends to be higher than Nádor, which tends to be lower than Alföld.


6.3. The grain moisture

Figures 19-20 and Table 12 as well as Table 31-32-32 and Figure 32 (Appendix 1) give information on N application effects on grain moisture content in the studied winter wheat varieties. Figure 19 Shows the experimental site #1's results for 2015-2016 and 2016-2017 crop seasons: the moisture content values slightly increased in some of the tested varieties by increasing the level of divided N application, but the changes found were not significant. However, the highest value was recorded on the untreated plot of MV Toldi with 17.70%, and the lowest result on the MV Nádor 80+40 kg/ha⁻¹ N applied plot with 5.30% humidity.



19. Figure Impact of N topdressing applications on moisture content of wheat varieties. 2015-2016 and 2016–2017 (Nagygyombos, Hungary)

Figure 20 shows the experimental site #2's results for the 2018-2019 crop season. Grain moisture content values changes found were not significant, however, and the highest value was recorded on 120+40 kg/ ha⁻¹ N applied plot of MV Karéj with 12.70% and the lowest result on Alföld 80 kg/ ha⁻¹ N applied plot with 9.00% humidity content.



20. Figure Impact of N topdressing applications on moisture content of wheat varieties. 2018-2019 (Gödöllő, Hungary)

Comparing the two sites, site #2 had higher grain moisture content for both varieties, Alföld had 86.33% and MV Karéj 18.30% more moisture content at site #2 than site #1; however, results showed significant differences between year, site, and genotype via the one-way ANOVA test of moisture content. But that moisture content ($F_{(5,102)} = .253, p > .05$) did not have significant effect due to different dose and application type of N applications, as is shown on Table 36-54.

The significance level of the difference based on year, site, genotypes and treatment in terms of moisture content measured in the sample of all genotypes was examined through ANOVA and outcomes were reflected in Table 12.

| Moisture Content (%) | | N | M | SD | F/t | df | p | Difference |
|----------------------|---------|-----|-------|------|--------|-------|------|------------|
| Year | 2016 | 90 | 8.19 | 1.66 | 59.097 | 2-213 | .000 | 3 > 2 > 1 |
| | 2017 | 90 | 9.77 | 0.61 | | | | |
| | 2019 | 36 | 10.57 | 1.32 | | | | |
| Site | Hatvan | 180 | 8.98 | 1.48 | -6.009 | 214 | .000 | 2 > 1 |
| | Gödöllő | 36 | 10.57 | 1.32 | | | | |
| Genotypes | Alföld | 54 | 8.56 | 1.50 | 46.976 | 4-211 | .000 | 2 > 1 |

| | | | | | | | | |
|-----------|--------------|----|-------|------|------|-------|------|--|
| | Karéj | 54 | 10.55 | 1.02 | | | | |
| | Nádor | 36 | 8.37 | 1.52 | | | | |
| | Toborzó | 36 | 8.01 | 0.79 | | | | |
| | Toldi | 36 | 10.42 | 0.23 | | | | |
| Treatment | 0 kg/ha | 36 | 9.18 | 1.52 | .063 | 5-210 | .997 | |
| | 80 kg/ha | 36 | 9.26 | 1.46 | | | | |
| | 80+40 kg/ha | 36 | 9.16 | 1.62 | | | | |
| | 120 kg/ha | 36 | 9.24 | 1.58 | | | | |
| | 120+40 kg/ha | 36 | 9.33 | 1.64 | | | | |
| | 160 kg/ha | 36 | 9.30 | 1.67 | | | | |

12. Table Comparison of the Means of Moisture Content Parameters Measured in the Sample of All Genotypes Based on Categorical Variables, 2016 and 2017 Hatvan and 2019 Gödöllő crop years.

As seen in Table 12, the outcomes reveal that there is no significant difference in statistical level in moisture content based on treatment ($F(5-210) = .063$, $p > .05$); however, there is a significant difference in statistical level in moisture content based on year ($F(2-213) = 59.097$, $p < .001$), site ($t(214) = -6.009$, $p < .001$) and genotypes ($F(4-211) = 46.976$, $p < .001$) in the sample of all genotypes. Accordingly, the mean values of moisture content in 2018-2019 tend to be higher than in 2016-2017, which tends to be higher than in 2015-2016; Gödöllő tends to be higher than Hatvan, and Karéj tends to be higher than Alföld.

6.4. The grain protein


Table 13-20 and Figure 21-22 as well as Table 34-35 and Figure 33-34-35 (Appendix 1) show the grain protein values in site #1 in 2015-2016, 2016-2017 and site #2 in 2018-2019 crop years. Protein amounts changed from 6.86% to 17.33 %.

| Protein content % 2015-2016 and 2016–2017 | | | | | |
|---|--------|----------|----------|------------|----------|
| N | Alföld | MV Nádor | MV Karéj | MV Toborzó | MV Toldi |
| 0 | 13,36 | 11,56 | 11,15 | 11,94 | 12,17 |
| 80 | 14,24 | 12,48 | 12,55 | 12,91 | 13,40 |
| 80+40 | 15,19 | 12,96 | 13,35 | 13,49 | 13,56 |
| 120 | 15,50 | 13,28 | 13,40 | 13,68 | 13,71 |
| 120+40 | 15,43 | 13,61 | 14,10 | 14,27 | 14,18 |
| 160 | 15,89 | 13,96 | 14,35 | 14,45 | 14,36 |

13. Table Impact of N topdressing applications on protein content of wheat varieties. 2015-2016 and 2016–2017 (Nagygyombos, Hungary)

| Protein content % 2018–2019 | | |
|------------------------------------|---------------|-----------------|
| N | Alföld | MV Karéj |
| 0 | 13,60 | 13,80 |
| 80 | 13,00 | 15,20 |
| 80+40 | 13,40 | 15,90 |
| 120 | 13,30 | 15,50 |
| 120+40 | 14,10 | 17,20 |
| 160 | 13,80 | 16,90 |


14. Table Impact of N topdressing applications on protein content of wheat varieties. 2018–2019 (Gödöllő, Hungary)



21. Figure Impact of N topdressing applications on protein content of wheat varieties. 2015-2016 and 2016–2017 (Nagygombos, Hungary) and 2018-2019 (Gödöllő, Hungary) combined

Based on the results of the experiment, we can conclude that increasing levels of N topdressing had a significant effect on grain protein content in all studied winter wheat varieties, either in split or undivided dose applications. The results obtained are in harmony with Dubetz et al., (1979), Pollhamer (1981), Vida et al., (1996), Varga and Svecnjak (2006), Öztürk and Gökkuş (2008), and Pepó (2010) studies. There were varietal differences, too, as untreated plots had remarkable differences between varieties such as 2016-2017 crop seasons of untreated Alföld and MV Karéj's plots showing 32.63% differences shown on Figure 22 boxplot. A special increase in dose applications had a remarkable effect in the experimental year 2017. The highest 17.33% grain protein content was observed on the Alföld plot with split 120+40 kg ha⁻¹ N application at site one 2015-2016 crop season and the lowest, 6.86 %, was obtained on Karéj

untreated plot at site #1's 2016-2017 crop season. Split dose application did not have a significant effect compared to the same total amount of undivided application; however, sites had significant effect under identical agronomical treatments. Alföld had up to 10.30% differences between the two sites regarding grain protein content, and MV Karéj plots had 19.80% greater results in the general average of grain protein content of the tested MV Karéj plots. The MV Karéj have shown greater results on site #2 compared to site #1, except for the 80 and 120 kg ha⁻¹ N single dose application according to One Way ANOVA statistics shown on Table 4.



22. Figure Impact of N topdressing applications on protein content of wheat varieties untreated plots. 2015-2016 and 2016-2017 (Nagygyombos, Hungary)

As seen in Table 36-54 the outcomes reveal that there is a significant difference in the statistical levels of grain protein content between trial years ($F_{(2,105)} = 8.829, p < .001$), site ($t_{(106)} = -2.885, p < .01$), genotype ($t_{(87.225)} = 2.094, p < .05$) and the treatment ($F_{(5,102)} = 10.238, p < .001$).

The significance level of the difference based on year, site, genotypes, and treatment by means of protein content measured in the sample of all genotypes was examined through ANOVA and outcomes were reflected in Table 15.

| Protein Content (%) | | N | M | SD | F/t | df | p | Difference |
|---------------------|--------------|-----|-------|------|--------|-------|------|---|
| Year | 2016 | 90 | 14.12 | 1.12 | 25.938 | 2-213 | .000 | 3 > 1 > 2 |
| | 2017 | 90 | 13.08 | 1.60 | | | | |
| | 2019 | 36 | 14.93 | 1.48 | | | | |
| Site | Hatvan | 180 | 13.60 | 1.47 | -4.946 | 214 | .000 | 2 > 1 |
| | Gödöllő | 36 | 14.93 | 1.48 | | | | |
| Genotypes | Alföld | 54 | 14.63 | 1.17 | 8.426 | 4-211 | .000 | 1 > 2 > 3 |
| | Karéj | 54 | 13.99 | 1.94 | | | | |
| | Nador | 36 | 12.95 | 1.01 | | | | |
| | Toborzo | 36 | 13.46 | 1.62 | | | | |
| | Toldi | 36 | 13.60 | 1.13 | | | | |
| Treatment | 0 kg/ha | 36 | 12.24 | 1.57 | 18.050 | 5-210 | .000 | 2, 3, 4, 5, 6 > 1; 4, 5, 6 > 2; 5, 6 > 3; 6 > 4 |
| | 80 kg/ha | 36 | 13.33 | 1.22 | | | | |
| | 80+40 kg/ha | 36 | 13.93 | 1.43 | | | | |
| | 120 kg/ha | 36 | 14.07 | 1.28 | | | | |
| | 120+40 kg/ha | 36 | 14.56 | 1.23 | | | | |
| | 160 kg/ha | 36 | 14.80 | 1.08 | | | | |

15. Table Comparison of the Means of Protein Content Parameters Measured in the Sample of All Genotypes Based on Categorical Variables, 2016,2017 Hatvan and 2019 Gödöllő crop years.

As seen in Table 15, outcomes revealed that there is a significant difference in statistical level in protein content based on year ($F(2-213) = 25.938$, $p < .001$), site ($t(214) = -4.946$, $p < .001$), genotypes ($F(4-211) = 8.426$, $p < .001$) and treatment ($F(5-210) = 18.050$, $p < .001$) in the sample of all genotypes. Accordingly, the mean values of protein content in 2018-2019 tend to be higher than in 2015-2016 tend to be higher than in 2016-2017; Gödöllő tends to be higher than in Hatvan; Alföld tends to be higher than in Karéj, which tends to be higher than in Nádor. 2, 3, 4, 5 and 6 tend to be higher than in 1; 4, 5 and 6 tend to be higher than in 2; 5 and 6 tend to be higher than in 3; and 6 tends to be higher than in 4.

The significance level of the difference, based on genotypes and treatment by means of protein content measured in the sample of all genotypes in 2015-2016, was examined through ANOVA and outcomes were reflected in Table 16.

| Protein Content (%) | N | M | SD | F/t | df | p | Difference |
|---------------------|---|---|----|-----|----|---|------------|
|---------------------|---|---|----|-----|----|---|------------|

| | | | | | | | | |
|-----------|--------------|----|-------|------|--------|------|------|---------------------------------|
| Genotypes | Alföld | 18 | 15.14 | 1.02 | 13.004 | 4-85 | .000 | 1 > 2, 3, 5; 3, 4 > 2; 4 > 3, 5 |
| | Karéj | 18 | 13.93 | 1.06 | | | | |
| | Nádor | 18 | 13.24 | 0.96 | | | | |
| | Toborzó | 18 | 14.64 | 0.76 | | | | |
| | Toldi | 18 | 13.64 | 0.62 | | | | |
| Treatment | 0 kg/ha | 15 | 12.92 | 0.83 | 11.433 | 5-84 | .000 | 3, 4, 5, 6 > 1, 2; 5 > 4 |
| | 80 kg/ha | 15 | 13.48 | 0.90 | | | | |
| | 80+40 kg/ha | 15 | 14.44 | 1.06 | | | | |
| | 120 kg/ha | 15 | 14.20 | 0.75 | | | | |
| | 120+40 kg/ha | 15 | 14.91 | 0.94 | | | | |
| | 160 kg/ha | 15 | 14.76 | 0.80 | | | | |

16. Table Comparison of the Means of Protein Content Parameters Measured in the Sample of All Genotypes in 2016, Based on Categorical Variables, Hatvan site.

As seen in Table 16, according to the outcomes it is found out that there is a significant difference in statistical level in protein content based on genotypes ($F(4-85) = 13.004$, $p < .001$) and treatment ($F(5-84) = 11.433$, $p < .001$) in the sample of all genotypes in 2015-2016. Accordingly, the mean values of protein content in Alföld tend to be higher than in Karéj, Nádor and Toldi, Nádor and Toborzó tend to be higher than in Karéj, Toborzó tends to be higher than in Nádor and Toldi; 3, 4, 5 and 6 tend to be higher than in 1 and 2; and 5 tends to be higher than in 4.

The significance level of the difference, based on genotypes and treatment in terms of protein content measured in the sample of Alföld and Karéj genotypes in Gödöllő in 2018-2019, was examined through Independent Samples t-Test and outcomes were reflected in Table 17.

| Protein Content (%) | | N | M | SD | F/t | df | p | Difference |
|---------------------|--------------|----|-------|------|--------|------|------|------------|
| Genotypes | Alföld | 18 | 14.21 | 0.86 | -3.285 | 34 | .002 | 2 > 1 |
| | Karéj | 18 | 15.65 | 1.64 | | | | |
| Treatment | 0 kg/ha | 6 | 13.71 | 0.92 | 2.370 | 5-30 | .063 | |
| | 80 kg/ha | 6 | 14.20 | 1.34 | | | | |
| | 80+40 kg/ha | 6 | 15.18 | 1.17 | | | | |
| | 120 kg/ha | 6 | 14.88 | 1.93 | | | | |
| | 120+40 kg/ha | 6 | 15.77 | 1.30 | | | | |
| | 160 kg/ha | 6 | 15.85 | 1.29 | | | | |

17. Table Comparison of the Means of Protein Content Parameters Measured in the Sample of Alföld and Karéj Genotypes in Gödöllő in 2019, Based on Categorical Variables, Gödöllő site.

As seen in Table 17, the outcomes revealed that there is no significant difference in statistical level in protein content based on treatment ($F(5-30) = 2.370$, $p > .05$), however, there is a

significant difference in statistical level in protein content based on genotypes ($t(34) = -3.285$, $p < .01$) in the sample of Alföld and Karéj genotypes in Gödöllő in 2018-2019. Accordingly, the mean values of protein content in Karéj tend to be higher than in Alföld.

The significance level of the difference, based on year, site and treatment in terms of protein content measured in the sample of Karéj genotype, was examined through ANOVA and outcomes were reflected in Table 18.

| Protein Content (%) | | N | M | SD | F/t | df | p | Difference |
|---------------------|--------------|----|-------|------|--------|------|------|--------------------------|
| Year | 2016 | 18 | 13.93 | 1.06 | 23.524 | 2-51 | .000 | 3 > 1 > 2 |
| | 2017 | 18 | 12.39 | 1.51 | | | | |
| | 2019 | 18 | 15.65 | 1.64 | | | | |
| Site | Hatvan | 36 | 13.16 | 1.50 | -5.561 | 52 | .000 | 2 > 1 |
| | Gödöllő | 18 | 15.65 | 1.64 | | | | |
| Treatment | 0 kg/ha | 9 | 12.09 | 1.91 | 4.060 | 5-48 | .004 | 3, 4, 5, 6 > 1; 5, 6 > 2 |
| | 80 kg/ha | 9 | 13.35 | 1.66 | | | | |
| | 80+40 kg/ha | 9 | 14.17 | 1.80 | | | | |
| | 120 kg/ha | 9 | 14.10 | 1.84 | | | | |
| | 120+40 kg/ha | 9 | 15.01 | 1.58 | | | | |
| | 160 kg/ha | 9 | 15.21 | 1.43 | | | | |

18. Table Comparison of the Means of Protein Content Parameters Measured in the Sample of Karej Genotype Based on Categorical Variables

As seen in Table 18, outcomes revealed that there is a significant difference in statistical level in protein content based on year ($F(2-51) = 23.524$, $p < .001$), site ($t(52) = -5.561$, $p < .001$) and treatment ($F(5-48) = 4.060$, $p < .01$) in the sample of Karéj genotype. Accordingly, the mean values of protein content in 2018-2019 is higher than 2015-2016, and 2015-2016 is higher than 2016-2017; Gödöllő tends to be higher than in Hatvan; 3, 4, 5 and 6 tend to be higher than in 1; and 5 and 6 tend to be higher than in 2.

6.5. The gluten content

| Gluten content % | | | | | |
|------------------|--------|----------|----------|------------|----------|
| N | Alföld | MV Nádor | MV Karéj | MV Toborzó | MV Toldi |
| 0 | 27,41 | 24,98 | 21,20 | 25,11 | 24,31 |
| 80 | 31,84 | 27,89 | 25,75 | 28,01 | 28,18 |
| 80+40 | 34,48 | 29,71 | 28,40 | 30,34 | 28,81 |

| | | | | | |
|---------------|-------|-------|-------|-------|-------|
| 120 | 35,16 | 30,33 | 28,65 | 31,27 | 28,06 |
| 120+40 | 35,28 | 30,58 | 30,35 | 31,99 | 30,44 |
| 160 | 36,37 | 32,36 | 31,65 | 33,31 | 31,45 |

19. Table Impact of N topdressing applications on gluten content of wheat varieties. 2015-2016 and 2016–2017 (Nagyombos, Hungary)

| Gluten content % | | |
|-------------------------|---------------|-----------------|
| N | Alföld | MV Karéj |
| 0 | 25,50 | 28,60 |
| 80 | 23,80 | 32,70 |
| 80+40 | 23,50 | 34,90 |
| 120 | 24,30 | 34,20 |
| 120+40 | 26,50 | 39,40 |
| 160 | 25,80 | 38,10 |

20. Table Impact of N topdressing applications on gluten content of wheat varieties. 2018–2019 (Gödöllő, Hungary) 1

Grain gluten amounts changed from 10.00 % to 41.10 %. The highest value was observed on the MV Karéj plot with split 120+40 kg ha⁻¹ N application as 41.10% and the lowest, 10.00 %, had been obtained on MV Karéj untreated plot. Grain gluten content was significantly affected by increasing doses of N applications as well with increased split dose applications. Similar examples have been reported by several authors (Györi, 2006; Kismányoky and Tóth, 2010; Rakszegi et al., 2016). Table 19-20-21-22-23-24 as well as Figure 36-40 (Appendix 1) show strong effect of N application on grain gluten content regardless of crop year, variety, or split/undivided application. Alföld had up to 34.22% differences between two sites in terms of grain gluten content, and MV Karéj plots had 25.23% higher results in the general average of protein content of the tested MV Karéj plots between two experimental sites. In addition, split dose application did not have significant effect on grain gluten content compared to the same amount of undivided application.

The significance level of the difference based on year, site, genotypes and treatment, in terms of gluten content measured in the sample of all genotypes, was examined through ANOVA and outcomes were reflected in Table 21.

| Gluten Content (%) | N | M | SD | F/t | df | p | Difference |
|---------------------------|----------|----------|-----------|------------|-----------|----------|-------------------|
| Year | 2016 | 90 | 30.73 | 24.209 | 2-213 | .000 | 3 > 1 > 2 |
| | 2017 | 90 | 28.85 | | | | |

| | | | | | | | | |
|-----------|--------------|-----|-------|------|--------|-------|------|--|
| | 2019 | 36 | 34.52 | 3.90 | | | | |
| Site | Hatvan | 180 | 29.79 | 4.28 | -6.141 | 214 | .000 | 2 > 1 |
| | Gödöllő | 36 | 34.52 | 3.90 | | | | |
| Genotypes | Alföld | 54 | 34.12 | 3.42 | 14.093 | 4-211 | .000 | 1 > 2, 3 |
| | Karěj | 54 | 29.62 | 5.23 | | | | |
| | Nádor | 36 | 29.30 | 3.01 | | | | |
| | Toborzó | 36 | 30.01 | 4.43 | | | | |
| | Toldi | 36 | 28.55 | 3.62 | | | | |
| Treatment | 0 kg/ha | 36 | 25.70 | 5.06 | 21.138 | 5-210 | .000 | 2, 3, 4, 5, 6 > 1; 3, 4, 5, 6 > 2; 5, 6 > 3; 6 > 4 |
| | 80 kg/ha | 36 | 29.00 | 3.38 | | | | |
| | 80+40 kg/ha | 36 | 31.01 | 3.62 | | | | |
| | 120 kg/ha | 36 | 31.26 | 3.40 | | | | |
| | 120+40 kg/ha | 36 | 32.78 | 3.84 | | | | |
| | 160 kg/ha | 36 | 33.72 | 2.93 | | | | |

21. Table Comparison of the Means of Gluten Content Parameters Measured in the Sample of All Genotypes Based on Categorical Variables

As seen in Table 21, outcomes revealed that there is a significant difference in statistical level in gluten content based on year ($F(2-213) = 24.209$, $p < .001$), site ($t(214) = -6.141$, $p < .001$), genotypes ($F(4-211) = 14.093$, $p < .001$) and treatment ($F(5-210) = 21.138$, $p < .001$) in the sample of all genotypes. Accordingly, the mean values of gluten content in 2018-2019 tend to be higher than in 2015-2016, which tend to be higher than in 2017. Gödöllő tends to be higher than Hatvan; Alföld tends to be higher than in Karěj and Nádor; 2, 3, 4, 5 and 6 tend to be higher than in 1; 3, 4, 5 and 6 tend to be higher than in 2; 5 and 6 tend to be higher than in 3, and 6 tends to be higher than in 4.

The significance level of the difference based on genotypes and treatment, in terms of gluten content measured in the sample of all genotypes in 2016, was examined through ANOVA and outcomes were reflected in Table 22.

| Gluten Content (%) | | N | M | SD | F/t | df | p | Difference |
|--------------------|----------|----|-------|------|--------|------|------|-------------------------------------|
| Genotypes | Alföld | 18 | 34.24 | 3.15 | 17.043 | 4-85 | .000 | 1 > 2, 3, 5; 4 > 2; 4 > 3; 3, 4 > 5 |
| | Karěj | 18 | 28.71 | 3.29 | | | | |
| | Nádor | 18 | 30.23 | 2.78 | | | | |
| | Toborzó | 18 | 32.71 | 2.50 | | | | |
| | Toldi | 18 | 27.75 | 2.04 | | | | |
| Treatment | 0 kg/ha | 15 | 26.60 | 2.78 | 11.632 | 5-84 | .000 | 3, 4, 5, 6 > 1, 2 |
| | 80 kg/ha | 15 | 28.75 | 3.01 | | | | |

| | | | | | | | | |
|--|-----------------|----|-------|------|--|--|--|--|
| | 80+40 kg/ha | 15 | 31.99 | 3.14 | | | | |
| | 120 kg/ha | 15 | 31.16 | 2.57 | | | | |
| | 120+40 kg/ha | 15 | 32.94 | 3.04 | | | | |
| | 160 kg/ha | 15 | 32.94 | 2.80 | | | | |

22. Table Comparison of the Means of Gluten Content Parameters Measured in the Sample of All Genotypes in 2016 Based on Categorical Variables

As seen in Table 22, outcomes revealed that there is a significant difference in statistical level in gluten content based on genotypes ($F(4-85) = 17.043$, $p < .001$) and treatment ($F(5-84) = 11.632$, $p < .001$) in the sample of all genotypes in 2016. Accordingly, the mean values of gluten content in Alföld tend to be higher than in Karéj, Nádor and Toldi, Toborzó tends to be higher than in Karéj, Toborzó tends to be higher than in Nádor, and Nádor and Toborzó tend to be higher than in Toldi. 3, 4, 5 and 6 tend to be higher than in 1 and 2.

The significance level of the difference based on year and treatment, in terms of gluten content measured in the sample of Toldi genotype, was examined through Independent Samples t-Test and outcomes were reflected in Table 23.

| Gluten Content (%) | | N | M | SD | F/t | df | p | Difference |
|--------------------|--------------|----|-------|------|--------|------|------|-------------|
| Year | 2016 | 18 | 27.75 | 2.04 | -1.345 | 34 | .188 | |
| | 2017 | 18 | 29.36 | 4.64 | | | | |
| Treatment | 0 kg/ha | 6 | 24.32 | 3.67 | 3.928 | 5-30 | .007 | 3, 5, 6 > 1 |
| | 80 kg/ha | 6 | 28.21 | 3.26 | | | | |
| | 80+40 kg/ha | 6 | 28.83 | 2.54 | | | | |
| | 120 kg/ha | 6 | 28.08 | 3.27 | | | | |
| | 120+40 kg/ha | 6 | 30.45 | 2.54 | | | | |
| | 160 kg/ha | 6 | 31.44 | 2.80 | | | | |

23. Table Comparison of the Means of Gluten Content Parameters Measured in the Sample of Toldi Genotype Based on Categorical Variables

As seen in Table 23, outcomes revealed that there is no significant difference in statistical level in gluten content based on year ($t(34) = -1.345$, $p > .05$), however, there is a significant difference in statistical level in gluten content based on treatment ($F(5-30) = 3.928$, $p < .01$) in the sample of Toldi genotype. Accordingly, the mean values of gluten content in 3, 5 and 6 tend to be higher than in 1.


The significance level of the difference based on genotypes and treatment, in terms of gluten content measured in the sample of Alföld and Karéj genotypes in Gödöllő in 2018-2019, was examined through Independent Samples t-Test and outcomes were reflected in Table 24.

| Gluten Content (%) | | N | M | SD | F/t | df | p | Difference |
|--------------------|--------------|----|-------|------|-------|------|------|-----------------------|
| Genotypes | Alföld | 18 | 35.53 | 2.37 | 1.589 | 34 | .121 | |
| | Karéj | 18 | 33.51 | 4.85 | | | | |
| Treatment | 0 kg/ha | 6 | 31.25 | 5.57 | 3.884 | 5-30 | .008 | 5, 6 > 1, 2, 3; 5 > 4 |
| | 80 kg/ha | 6 | 32.27 | 3.22 | | | | |
| | 80+40 kg/ha | 6 | 34.34 | 1.90 | | | | |
| | 120 kg/ha | 6 | 34.12 | 3.29 | | | | |
| | 120+40 kg/ha | 6 | 37.95 | 2.11 | | | | |
| | 160 kg/ha | 6 | 37.18 | 2.07 | | | | |

24. Table Comparison of the Means of Gluten Content Parameters Measured in the Sample of Alföld and Karéj Genotypes in Gödöllő in 2019 Based on Categorical Variables

As seen in Table 24, according to the outcomes it is found out that there is no significant difference in statistical level in gluten content based on genotypes ($t(34) = 1.589$, $p > .05$); however, there is a significant difference in statistical level in gluten content based on treatment ($F(5-30) = 3.884$, $p < .01$) in the sample of Alföld and Karéj genotypes in Gödöllő in 2018-2019. Accordingly, the mean values of gluten content in 5 and 6 tend to be higher than in 1, 2 and 3; 5 tends to be higher than in 4.

6.6 The Zeleny number (ml)



23. Figure Impact of N topdressing applications on Zeleny Nr. of wheat varieties. 2015-2016 and 2016-2017 (Nagygyombos, Hungary) 1

| Zeleny Nr/mL | | | | | |
|---------------|--------|----------|----------|------------|----------|
| N | Alföld | MV Nádor | MV Karěj | MV Toborzó | MV Toldi |
| 0 | 28,64 | 29,93 | 23,60 | 26,60 | 29,56 |
| 80 | 36,95 | 36,54 | 33,30 | 35,39 | 35,13 |
| 80+40 | 41,71 | 39,06 | 38,80 | 38,45 | 35,69 |
| 120 | 43,64 | 41,29 | 39,55 | 40,79 | 36,44 |
| 120+40 | 43,64 | 43,01 | 43,70 | 42,17 | 38,28 |
| 160 | 45,90 | 43,75 | 45,10 | 44,50 | 41,14 |

25. Table Impact of N topdressing applications on Zeleny Nr. of wheat varieties. 2015-2016 and 2016–2017 (Nagygombos, Hungary) 2

| Zeleny Nr/mL | | |
|--------------|--------|----------|
| N | Alföld | MV Karéj |
| 0 | 38,50 | 41,10 |
| 80 | 37,60 | 50,80 |
| 80+40 | 35,60 | 55,40 |
| 120 | 35,90 | 51,90 |
| 120+40 | 42,70 | 60,70 |
| 160 | 41,30 | 58,90 |

26. Table Impact of N topdressing applications on Zeleny Nr. of wheat varieties. 2015-2016 and 2018–2019 (Gödöllő, Hungary)

The significance level of the difference based on year in terms of parameters was examined through ANOVA and outcomes were reflected in Figure 23 and Table 26-30 as well as Table 36-54 (Appendix 1)

As seen in Table 36-54, outcomes revealed that there is a significant difference in statistical level in Zeleny number ($F_{(2,105)} = 18.886, p < .001$) based on year, ($t_{(103.445)} = 3.234, p < .01$) based on site and ($F_{(5,102)} = 13.148, p < .001$) based on treatment. However, no significance ($t_{(106)} = 1.635, p > .05$) based on genotypes.

The significance level of the difference based on year, site, genotypes and treatment, in terms of Zeleny number measured in the sample of all genotypes, was examined through ANOVA and outcomes were reflected in Table 27.

| Zeleny Number | | N | M | SD | F/t | df | p | Difference |
|---------------|---------|-----|-------|-------|--------|-------|------|------------|
| Year | 2016 | 90 | 35.05 | 6.04 | 31.764 | 2-213 | .000 | 2, 3 > 1 |
| | 2017 | 90 | 42.68 | 8.80 | | | | |
| | 2019 | 36 | 45.88 | 10.12 | | | | |
| Site | Hatvan | 180 | 38.87 | 8.44 | -4.391 | 214 | .000 | 2 > 1 |
| | Gödöllő | 36 | 45.88 | 10.12 | | | | |
| Genotypes | Alföld | 54 | 39.59 | 8.59 | 1.672 | 4-211 | .158 | |
| | Karéj | 54 | 42.55 | 11.45 | | | | |
| | Nádor | 36 | 38.93 | 7.62 | | | | |
| | Toborzó | 36 | 37.98 | 7.41 | | | | |

| | | | | | | | | |
|-----------|--------------|----|-------|------|--------|-------|------|---|
| | Toldi | 36 | 40.10 | 8.39 | | | | |
| Treatment | 0 kg/ha | 36 | 29.59 | 8.82 | 23.141 | 5-210 | .000 | 2, 3, 4, 5, 6 > 1; 4, 5, 6 > 2; 5, 6 > 3; 6 > 4 |
| | 80 kg/ha | 36 | 37.56 | 7.49 | | | | |
| | 80+40 kg/ha | 36 | 40.65 | 6.78 | | | | |
| | 120 kg/ha | 36 | 41.71 | 6.91 | | | | |
| | 120+40 kg/ha | 36 | 44.71 | 6.58 | | | | |
| | 160 kg/ha | 36 | 46.00 | 7.58 | | | | |

27. Table Comparison of the Means of Zeleny Nr. Parameters Measured in the Sample of All Genotypes Based on Categorical Variables

As seen in Table 27, outcomes revealed that there is no significant difference in the statistical level in Zeleny number based on genotypes ($F(4-211) = 1.672$, $p > .05$); however, there is a significant difference in statistical level in Zeleny number based on year ($F(2-213) = 31.764$, $p < .001$), site ($t(214) = -4.391$, $p < .001$) and treatment ($F(5-210) = 23.141$, $p < .001$) in the sample of all genotypes. Accordingly, the mean values of Zeleny number in 2016-2017 and 2018-2019 tend to be higher than in 2015-2016; Gödöllő tends to be higher than in Hatvan; 2, 3, 4, 5 and 6 tend to be higher than in 1; 4, 5 and 6 tend to be higher than in 2; 5 and 6 tend to be higher than in 3; and 6 tends to be higher than in 4.

The significance level of the difference based on genotypes and treatment, in terms of Zeleny number measured in the sample of Alföld and Karéj genotypes in Gödöllő in 2018-2019, was examined through Independent Samples t-Test and outcomes were reflected in Table 28.

| Zeleny Number | | N | M | SD | F/t | df | p | Difference |
|---------------|--------------|----|-------|-------|--------|------|------|------------|
| Genotypes | Alföld | 18 | 38.61 | 6.26 | -6.188 | 34 | .000 | 2 > 1 |
| | Karéj | 18 | 53.14 | 7.75 | | | | |
| Treatment | 0 kg/ha | 6 | 39.78 | 8.35 | 1.148 | 5-30 | .357 | |
| | 80 kg/ha | 6 | 44.23 | 8.03 | | | | |
| | 80+40 kg/ha | 6 | 45.52 | 11.59 | | | | |
| | 120 kg/ha | 6 | 43.88 | 9.99 | | | | |
| | 120+40 kg/ha | 6 | 51.72 | 10.56 | | | | |
| | 160 kg/ha | 6 | 50.12 | 11.06 | | | | |

28. Table Comparison of the Means of Zeleny Nr. Parameters Measured in the Sample of Alföld and Karéj Genotypes in Gödöllő in 2019 Based on Categorical Variables

As seen in Table 28, outcomes revealed that there is no significant difference in statistical level in Zeleny number based on treatment ($F(5-30) = 1.148$, $p > .05$), however, there is a significant difference in statistical level in Zeleny number based on genotypes ($t(34) = -6.188$, $p < .001$) in the sample of Alföld and Karéj genotypes in Gödöllő in 2018-2019. Accordingly, the mean values of zeleny number in Karéj tend to be higher than in Alföld.

The significance level of the difference based on genotypes and treatment, in terms of Zeleny number measured in the sample of Alföld and Karéj genotypes in 2015-2016, was examined through Independent Samples t-Test and outcomes were reflected in Table 29.

| Zeleny Number | | N | M | SD | F/t | df | p | Difference |
|---------------|--------------|----|-------|------|--------|------|------|--|
| Genotypes | Alföld | 18 | 33.24 | 6.54 | -.754 | 34 | .456 | 2 > 1 |
| | Karéj | 18 | 34.99 | 7.40 | | | | |
| Treatment | 0 kg/ha | 6 | 22.97 | 3.05 | 18.143 | 5-30 | .000 | 2, 3, 4, 5, 6 > 1; 3, 4, 5, 6 > 2; 5 > 4 |
| | 80 kg/ha | 6 | 30.02 | 4.54 | | | | |
| | 80+40 kg/ha | 6 | 37.01 | 4.94 | | | | |
| | 120 kg/ha | 6 | 35.54 | 2.89 | | | | |
| | 120+40 kg/ha | 6 | 40.06 | 3.49 | | | | |
| | 160 kg/ha | 6 | 39.08 | 3.01 | | | | |

29. Table Comparison of the Means of Zeleny Nr. Parameters Measured in the Sample of Alföld and Karéj Genotypes in 2016 Based on Categorical Variables

As seen in Table 29, outcomes revealed that there is no significant difference in statistical level in Zeleny number based on genotypes ($t(34) = -.754$, $p > .05$); however, there is a significant difference in statistical level in Zeleny number based on treatment ($F(5-30) = 18.143$, $p < .001$) in the sample of Alföld and Karéj genotypes in 2016. Accordingly, the mean values of Zeleny number in Karéj tend to be higher than in Alföld; 2, 3, 4, 5 and 6 tend to be higher than in 1; 3, 4, 5 and 6 tend to be higher than in 2; and 5 tends to be higher than in 4.

The significance level of the difference based on year, site and treatment, in terms of Zeleny number measured in the sample of Karéj genotype, was examined through ANOVA and outcomes were reflected in Table 30.

| Zeleny Number | | N | M | SD | F/t | df | p | Difference |
|---------------|--------|----|-------|-------|--------|------|------|------------|
| Year | 2016 | 18 | 34.99 | 7.40 | 21.924 | 2-51 | .000 | 3 > 1, 2 |
| | 2017 | 18 | 39.51 | 10.25 | | | | |
| | 2019 | 18 | 53.14 | 7.75 | | | | |
| Site | Hatvan | 36 | 37.25 | 9.10 | -6.338 | 52 | .000 | 2 > 1 |

| | | | | | | | | |
|-----------|-----------------|----|-------|-------|-------|------|------|-----------------------------|
| | Gödöllő | 18 | 53.14 | 7.75 | | | | |
| Treatment | 0 kg/ha | 9 | 29.05 | 11.20 | 5.969 | 5-48 | .000 | 3, 4, 5, 6 > 1; 5, 6 > 2 |
| | 80 kg/ha | 9 | 39.16 | 10.25 | | | | |
| | 80+40 kg/ha | 9 | 44.36 | 9.23 | | | | |
| | 120 kg/ha | 9 | 43.65 | 7.79 | | | | |
| | 120+40 kg/ha | 9 | 49.36 | 9.09 | | | | |
| | 160 kg/ha | 9 | 49.72 | 8.75 | | | | |

30. Table Comparison of the Means of Zeleny Nr. Parameters Measured in the Sample of Karéj Genotype Based on Categorical Variables

As seen in Table 30, outcomes revealed that there is significant difference in statistical level in Zeleny number based on year ($F(2-51) = 21.924$, $p < .001$), site ($t(52) = -6.338$, $p < .001$) and treatment ($F(5-48) = 5.969$, $p < .001$) in the sample of Karéj genotype. Accordingly, the mean values of Zeleny number in 2019 tend to be higher than in 2015-2016 and 2016-2017; Gödöllő tends to be higher than in Hatvan; 3, 4, 5 and 6 tend to be higher than in 1; and 5 and 6 tend to be higher than in 2.

7. CONCLUSION AND RECOMMENDATIONS

- Test weight measurements showed no significant differences by different dose and divided amounts of nitrogen fertilizer applications. However, year, site and genotype showed significant differences on test weight results of tested winter wheat species. Alföld and Karéj species had greater results both on sites and years.
- Thousand kernel weight measurements showed no significant differences by different dose and divided amount of nitrogen fertilizer applications. However, year, site and genotype showed significant differences on thousand kernel weight results of tested winter wheat species. Hatvan site had greater results compared to Gödöllő.
- The grain moisture measurements showed no significant differences by different dose and divided amounts of nitrogen fertilizer applications. However, year, site and genotype showed significant differences on grain moisture results of tested winter wheat species. Alföld and Karéj species had greater results both on sites and years. The Gödöllő site had greater results compared to Hatvan.
- The grain protein measurements showed significant effect on all studied years, sites, genotypes and treatment parameters. 2018-2019 was the leading year and the Gödöllő site had higher figures than Hatvan. The Alföld variety showed the best performance while Karéj was the second good performer among the five studied winter wheat varieties. Apart from this, the general results wherein all the years, sites and varieties were measured, showed the increased level of Nitrogen fertilizer supply caused an increase in the grain protein content (where no species, year sites separated). However, division of the given amounts of the nitrogen fertilizer did not have significant effect on tested winter wheat varieties. On the other hand, those at the Hatvan site's 2015-2016-year trials regarding all varieties and 2018-2019-year 80+40 kg/ha divided application, compared to 120 kg/ha un-divided applications, as well as the Karéj's

80+40 kg/ha compared to 120 kg/ha applications, showed that while increasing doses of the nitrogen fertilizer supply shows significant effect, divided doses of the nitrogen fertilizer showed greater effect compared to same amounts un-divided applications. Therefore, 80+40 kg/ha N application is recommended.

- The grain gluten measurements showed significant effect on all studied year, site, genotype and treatment parameters. 2018-2019 was the leading crop year as well as the Gödöllő site had greater figures than Hatvan. Alföld variety showed the best performance among the five studied winter wheat varieties. Apart from this, the general results wherein all the years, sites and varieties that were measured, showed that the increased level of Nitrogen fertilizer supply caused an increase in the grain protein content. However, division of the given amounts of the nitrogen fertilizer did not have a significant effect on tested winter wheat varieties. On the other hand, those at the Hatvan site's 2015-2016- crop year trials and Toldi varieties, and Hatvan's site trials results of 80+40 kg/ha divided application compared to 120 kg/ha un-divided applications, showed that while increasing doses of the nitrogen fertilizer supply shows significant effect, divided doses of the nitrogen fertilizer showed greater effect compared to the same amounts in undivided applications.

- The Zeleny number measurements showed significant effect on all studied years, sites, genotypes and treatment parameters. 2018-2019 was the leading year, and the Gödöllő site had higher figures than Hatvan. The Karéj variety showed the best performance among the five studied winter wheat varieties. Apart from that, the general results wherein all the years, sites and varieties that were measured, showed that the increased level of Nitrogen fertilizer supply caused an increase to the Zeleny number; however, division of the given amounts of the nitrogen fertilizer did not have significant effect on the tested winter wheat varieties. On the other hand, those in the Hatvan site 2015-2016-year trials, the Gödöllő site 2018-2019-year trials' results, and the Karéj varieties measurement of 80+40 kg/ha divided application compared to 120 kg/ha un-divided applications, and the 120+40 kg/ha divided applications compared to 160 kg/ha un-divided applications showed that, while increasing doses of the nitrogen fertilizer supply shows significant effect, divided doses of the nitrogen fertilizer showed greater effect compared to the same amounts in un-divided applications.

8. NEW SCIENTIFIC RESULTS

- 1- Measurements of this experiment proved that increasing amounts of Nitrogen fertilizer application raised the protein content of the tested winter varieties; however, 80+40 kg/ha Nitrogen fertilization was even more remarkable and recommended.
- 2- The Gödöllő site and Alföld variety showed the best results comparing the sites and the tested winter wheat varieties, in regard to the grain gluten content. While the grain gluten content increased with higher amounts of Nitrogen fertilizer applications, increasing the level of the Nitrogen application did not affect the result significantly.
- 3- By analyzing the Zeleny number of the chosen winter wheat varieties, results showed that the Karéj variety was significantly affected by split-dose application of Nitrogen fertilizer; however, all the varieties of Zeleny numbers rose with the increasing amount of N application.
- 4- All the tested winter wheat varieties showed significant differences, comparing all the measurement parameters (Zeleny number, Grain protein content, Grain gluten content, Test weight, Thousand kernel weight and Grain moisture) by site, genotype, and year.

9. SUMMARY

The Earth's rising population growth is contributing to increasing hunger, as well as creating insufficient and unbalanced nutrition, all of which continue to be major problems for human survival. Although different opinions are put forward for the solution, the most notable consensus among experts has been to engineer an increase in plant and animal products. Wheat, the most widely grown plant species in the world, is a plant of high strategic importance due to the fact that it has been grown since the earliest times; its agricultural process is easier than other plants; the product is more adaptable to transportation and storage conditions; and its inherent high nutrition solves one of the most pressing economic problems. Wheat is also one of the most predominant cereal grains in Hungary and Turkey, and it retains high economic value. The goal of wheat production is two-fold: to provide both quantity and quality. The aim of this study is to find the quality changes on tested winter wheat varieties with the application of Nitrogen fertilizer. As

Nitrogen fertilization has a considerably high economic and environmental effect on wheat farming and wheat quality, we therefore should find the most appropriate formula for supplying fertilizer to get the best results to be able to improve the quality of the winter wheat products, as well as consider the economic and environmental side effects of the this kind of farming approach.

Baking quality of wheat flour is determined by grain protein concentration (GPC) and its composition. It's also highly influenced by environmental factors, such as nitrogen (N) fertilization management (Xue et al., 2019). Increasing the level of protein content will raise the baking quality (Gabriel, et al., 2017; Wrigley and Batey, 2003). It's generally accepted that fertilization increases soil microbial biomass in agricultural soils on the short and long term (Nguyen et al., 2018), however, application of solely continuous N fertilizer can decrease soil microbial biomass (Wallenstein et al., 2006; Treseder, 2008). Numerous studies stipulate that overdose application, rather than the appropriate amount of nitrogen fertilizer application, had no significant positive effect on wheat quality parameters (Abedi et al., 2011; Cui et al., 2010; Chen et al., 2004; Lloveras et al., 2001) and even dropped the yield performances (Jolánkai, 1985). Nitrogen use efficiency is closely related to fertilization, the genetic and morphological characteristics of the plant, as well as the climate and soil characteristics.

Results showed that test weight, thousand kernel weight and grain moisture outcomes showed that there is no significant difference with applied dose and divided amount of nitrogen fertilizer applications. However, year, site and genotype showed significant differences.

The grain protein measurements showed significant effect on all studied years, sites, genotypes and treatment parameters. 2018-2019 was the leading year and the Gödöllő site had higher figures than Hatvan. The Alföld variety showed the best performance while Karék was the second good performer among the five studied winter wheat varieties. Apart from this, the general results wherein all the years, sites and varieties were measured, showed the increased level of Nitrogen fertilizer supply caused an increase in the grain protein content (where no species, year sites separated). However, division of the given amounts of the nitrogen fertilizer did not have significant effect on tested winter wheat varieties. On the other hand, those at the Hatvan site's 2015-2016-year trials regarding all varieties and 2018-2019-year 80+40 kg/ha divided application, compared to 120 kg/ha un-divided applications, as well as the Karék's 80+40 kg/ha compared to 120 kg/ha

applications, showed that while increasing doses of the nitrogen fertilizer supply shows significant effect, divided doses of the nitrogen fertilizer showed greater effect compared to same amounts un-divided applications. Therefore, 80+40 kg/ha N application is recommended.

- The grain gluten measurements showed significant effect on all studied year, site, genotype and treatment parameters. 2018-2019 was the leading crop year as well as the Gödöllő site had greater figures than Hatvan. Alföld variety showed the best performance among the five studied winter wheat varieties. Apart from this, the general results wherein all the years, sites and varieties that were measured, showed that the increased level of Nitrogen fertilizer supply caused an increase in the grain protein content. However, division of the given amounts of the nitrogen fertilizer did not have a significant effect on tested winter wheat varieties. On the other hand, those at the Hatvan site's 2015-2016- crop year trials and Toldi varieties, and Hatvan's site trials results of 80+40 kg/ha divided application compared to 120 kg/ha un-divided applications, showed that while increasing doses of the nitrogen fertilizer supply shows significant effect, divided doses of the nitrogen fertilizer showed greater effect compared to the same amounts in undivided applications.

- The Zeleny number measurements showed significant effect on all studied years, sites, genotypes and treatment parameters. 2018-2019 was the leading year, and the Gödöllő site had higher figures than Hatvan. The Karéj variety showed the best performance among the five studied winter wheat varieties. Apart from that, the general results wherein all the years, sites and varieties that were measured, showed that the increased level of Nitrogen fertilizer supply caused an increase to the Zeleny number; however, division of the given amounts of the nitrogen fertilizer did not have significant effect on the tested winter wheat varieties. On the other hand, those in the Hatvan site 2015-2016-year trials, the Gödöllő site 2018-2019-year trials' results, and the Karéj varieties measurement of 80+40 kg/ha divided application compared to 120 kg/ha un-divided applications, and the 120+40 kg/ha divided applications compared to 160 kg/ha un-divided applications showed that, while increasing doses of the nitrogen fertilizer supply shows significant effect, divided doses of the nitrogen fertilizer showed greater effect compared to the same amounts in un-divided applications.

10. THE PUBLICATIONS OF THE AUTHOR IN THE RESEARCH FIELD

1. Eser A. (2017): "Storage proteins in wheat (*Triticum aestivum* L.) and impact of mycotoxins affecting quality and quantity, with focus on nitrogen supply." *Columella Journal of Agricultural and Environmental Sciences*. 299. DOI: 10.18380/SZIE.COLUM.2017.4.1.suppl.
2. Eser A. – Kassai M.K. – Tarnawa Á. – Nyárai H.F. – Horváth Cs. – Jolánkai M. (2017): „Impact of nitrogen topdressing on the quality and quantity of wheat yield varieties.” *Georgikon for Agriculture* 21. 1. 109-113 pp.
3. Eser A. – Kassai M.K. – Tarnawa Á. – Nárai H.F. – Jolánkai M. (2017): “Impact of crop year and nitrogen topdressing on the quantity and quality of wheat yield.” *Columella 4.1. Suppl.* 157-162 pp.
4. Eser A. (2018): “Impact of nitrogen topdressing on the quality and quantity parameters of yield and grain protein of wheat (*Triticum aestivum* L.)” *17th Alps Adria Scientific Workshop, Hnanice, Czech Republic*. 144-145 pp. ISBN 978-963-269-734-5
5. Eser A., Kato H., Kempf L., Nyarai F., Kunos V., Szentpetery Z. (2019): “A world alimentation chance estimate based on protein production of crop species.” *Columella - Journal of Agricultural and Environmental Sciences Vol. 6, No. 1* (2019) DOI: 10.18380/SZIE.COLUM.2019.6.35
6. Eser A., Kató H., Kempf L., Jolánkai M. (2019): “Water footprint of yield protein content of twelve field crop species on a Hungarian crop site.” *Agrokémia és Talajtan*

7. Eser A. (2019): “Impact of nitrogen topdressing on the quality and quantity parameters of yield and grain protein of wheat (*Triticum aestivum L.*)” (2019) *Abstract book of the 18th Alps-Adria Scientific Workshop*. [Online ISBN 978-963-269-818-2](#). DOI: 10.34116/NTI.2019.AA.18. 52-53 pp.
8. Eser, A., Kassai, K.M., Kato, H., Kunos, V., Tarnava, A., & Jolánkai, M. (2020). “Impact of nitrogen topdressing on the quality parameters of winter wheat (*Triticum aestivum L.*) yield.” *Acta Alimentaria*, 49(3), 244-253.
<https://doi.org/10.1556/066.2020.49.3.2>
9. Eser A., Kato H., Czerödi-Kempf L., Hollo-Cseh R. (2020): “Effect of the crop year’s weather on the quality and quantity of winter wheat (*Triticum aestivum L.*)” *19th Alps-Adria Scientific Workshop Online* ISBN 978-963-269-896-0 DOI: 10.34116/NTI.2020.AA. 30.
10. Kassai M.K., Tarnawa Á., Nyárai H.F., Eser A., Kempf L., Jolánkai M. (2017): “A csapadék és a hőmérséklet évjáratati hatása őszi búza fajták fehérjetermelésére.” *Növénytermelés*. 66. 3. 1-9 pp.
11. Kassai M.K., Tarnawa Á., Nyárai H.F., Szentpétery Zs., Eser A., Kató H., Jolánkai, M. (2019): “Quality and quantity of winter wheat varieties in 22 years’ time range.” In: *Book of Proceedings. 2nd Conference on Long Term Field Experiments*. 20-21. November 2019. Nyíregyháza. Ed: Makádi M. Nyíregyháza, 28-31 pp.
12. Kassai M.K., Tarnawa Á., Nyárai H.F., Szentpétery Zs., Eser A., Kató H., Jolánkai, M. (2020): “Quality and quantity of winter wheat varieties in 22 years’ time range.” *Collumela Journal of Agricultural and Environmental Sciences*. DOI: 10.18380/SZIE.COLUM.2020.7.1.5, 5-9 pp.
13. Kató H., Eser A., Holló-Cseh R., Kassai M.K., Tarnawa Á. (2020): “Induced climate impacts on mycotoxins contamination in maize.” *19th Alps-Adria Scientific*

Workshop Online ISBN 978-963-269-896-0 DOI: 10.34116/NTI.2020.AA. 33 pp.

14. Jolánkai M., Kassai M.K., Eser A., Kempf L., Tarnawa Á. (2018): “Water footprint of protein yield of field crop species based on evapotranspiration patterns.” *Review on Agriculture and Rural Development*. 7. 1-2. 11-15 pp.
15. Jolánkai M., Tarnawa Á., Kassai M.K., Szentpétery Zs., Eser A., Kató H. (2019): “Crop year effects on the quantity and quality of winter wheat varieties.” In: *Transport of water, chemicals and energy in the soil-plant-atmosphere system*. Ed.: A. Celková. UH-SAV, Bratislava. 87-91 pp.
16. Jolánkai M., Tarnawa Á., Kassai M.K., Szentpétery Zs., Eser A., Kató H. (2020): “Crop year effects on the quality and quantity of winter wheat varieties.” *Acta Horticulturae et Regiotecturae – Special Issue*. 20–22 pp. DOI: 10.2478/ahr-2020-0017
17. Jolánkai M., Tarnawa A., Kassai M.K., Eser A., Kende Z. (2021): Water Footprint of the Protein Formation of Six of Field Crop Species. *Environ Anal Eco stud*. 8(3). EAES. 000686. 2021. DOI: 10.31031/EAES.2021.08.000686, 884-885 pp.
18. Tarnawa Á., Kassai M.K., Eser A., Kempf L., Jolánkai M. (2017): “Precíziós módszerek alkalmazása a szántóföldi növénytermesztésben.” In: *Precíziós gazdálkodás*. Ed: Milics G. Opal Média és Kommunikáció Kiadó, Budapest. 82-84 pp
19. Kunos, V., Cséplő, M., Seress, D., Eser, A., Kende, Z., Uhrin, A., Bányai, J., Bakonyi, J., Pál, M., Mészáros, K. “The Stimulation of Superoxide Dismutase Enzyme Activity and Its Relation with the *Pyrenophora teres f. teres* Infection in Different Barley Genotypes.” *Sustainability* 2022, 14, 2597.
<https://doi.org/10.3390/su14052597>
20. Khaeim H., Kende Z.*, Balla I., Csaba Gyuricza, Eser A., Tarnawa A. (2022): “The Effect of Temperature and Water Stresses on Seed Germination and Seedling Growth of Wheat (*Triticum aestivum L.*)” *MDPI Sustainability*, Sustainability 2022, 14(7),

3887; <https://doi.org/10.3390/su14073887>

11. BIBLIOGRAPHY

1. AACC (2010). Approved methods of analysis 11th ed. Determination of Falling Number Method No. 56-81.04.
2. Abad, A., J. Lloveras, A. Michelena, and J. Ferrán. (1996). Effect of nitrate levels and N fertilization on bread and durum wheat yield and quality and on residual N–NO₃ concentration under irrigation in the Ebro Valley (Spain). p. 514–515. In *Proc. Congr. Eur. Soc. of Agron.*, 4th, Veldoven, the Netherlands. 7–11 July 1996.
3. Abedi, T., Alemzadeh, A., & Kazemeini, S. A. (2011). Wheat Yield and Grain Protein Response to Nitrogen Amount and Timing. *Australian Journal of Crop Science*, 5(3), 330–336.
<https://search.informit.org/doi/10.3316/informit.279825257779534>
4. Ahmad, A., Abrol, Y. P., & Abdin, M. Z. (1999). Effect of split application of sulphur and nitrogen on growth and yield attributes of Brassica genotypes differing in time of flowering. *Canadian journal of plant science*, 79(2), 175-180.
5. Akkaya, A. (1994). Erzurum koşullarında Farklı Ekim Sıklıklarının İki Kışlık Buğday Çeşidinde Verim ve Verim Unsurlarına Etkileri. *Turkish Journal of Agriculture and Forestry* 18: 161-168.
6. Alcoz, M.M., Hons, F.M. and Haby, V.A. (1993). Nitrogen Fertilization Timing Effect on Wheat Production, Nitrogen Uptake Efficiency, and Residual Soil Nitrogen. *Agron. J.*, 85: 1198-1203. <https://doi.org/10.2134/agronj1993.00021962008500060020x>
7. Alipatra, A., Kundu, C. K., Mandal, M. K., Banerjee, H., & Bandopadhyay, P. (2013). Yield and quality improvement in fodder oats (*Avena sativa* L.) through split application of fertilizer and cutting management. *Journal of Crop and Weed*, 9(2), 193-195.
8. Al-Moshileh, A. M., Errebhi, M. A., & Motawei, M. I. (2005). Effect of various potassium and nitrogen rates and splitting methods on potato under sandy soil and arid environmental conditions. *Emirates Journal of Food and Agriculture*, 1-9.
9. Ankara, T.C. Ankara Valiliği İl Gıda Tarım ve Hayvancılık Müdürlüğü. (2015). *Buğday Yetiştiriciliği*.
<https://ankara.tarimorman.gov.tr/Belgeler/liftet/bugdayyetistiriciligi.pdf>
10. Anonymous (2012). Buğdayda Kalite ve Kaliteyi Etkileyen Faktörler. *University of Namık Kemal Faculty of Agriculture Crop Production Department*.

11. Arana, I. (ED.) (2016). Physical properties of foods: Novel measurement techniques and applications. *CRC Press*, p 420.
12. Arpacioğlu A. G. N., (2018). Determination of different nitrogen doses on yield and quality traits and nitrogen uptake/use efficiency under Eskisehir climatic conditions. Eskisehir, TR: Eskişehir Osmangazi University
13. Atar B. (2017). Gıdamız buğdayın, geçmişten geleceğe yolculuğu. *Yalvaç Akademi Dergisi*, 2(1), 1-12.
14. Atar, B., Kara, B., & Küçükşumuk, Z. (2017). Kışlık ekmeçlik buğday çeşitlerinin azot etkinliklerinin belirlenmesi. *Tarım Bilimleri Dergisi*, 23(1), 119-127.
15. Atli Arnarson, (2019). PhD -Wheat 101: Nutrition Facts and Health Effects – <https://www.healthline.com/nutrition/foods/wheat>. Date of search: 2021.05.19.
16. Awulachew, M. T. (2019). Environmental Impact on Processing Quality of Wheat Grain. *Int J Food Sci Nutr Diet. S*, 1, 1-8.
17. Ayyub, C. M., Wasim Haidar, M., Zulfiqar, F., Abideen, Z., & Wright, S. R. (2019). Potato tuber yield and quality in response to different nitrogen fertilizer application rates under two split doses in an irrigated sandy loam soil. *Journal of Plant Nutrition*, 42(15), 1850-1860.
18. Banu S., Duxbury J.M., Lauren J.G., Meisner C., Slam R.I. (2004). Wheat seed quality a study on farmer's seed. *Paper presented at 4th International Crop Science Congress, Brisbane, Queensland, Australia*.
19. Başar H., Tümsavaş Z., Katkat V.A., Özgümüş A. (1998). Effects of Various Nitrogen Sources and the Levels of Nitrogen on Yield and the Yield Components of the Wheat Cultivar Saraybosna. *Turkish Journal of Agriculture and Forestry*, 22, 59-63.
20. Bayfield, E. G. (1936). The influence of climate, soil, and fertilizers upon quality of soft winter wheat. *Ohio Agr. Exp. Sta. Bull.* 563.
21. Belete, F., Dechassa, N., Molla, A. et al. (2018). Effect of split application of different N rates on productivity and nitrogen use efficiency of bread wheat (*Triticum aestivum L.*).” *Agric & Food Secur* 7, 92. <https://doi.org/10.1186/s40066-018-0242-9>
22. Bellemare, M. F. (2015). Rising food prices, food price volatility, and social unrest. *Am. J. Agric. Econ.* 97, 1–21.
23. Biesiada, A., & Kuś, A. (2010). The effect of nitrogen fertilization and irrigation on yielding and nutritional status of sweet basil (*Ocimum basilicum L.*). *Acta Sci. Pol. Hortorum Cultus*, 9(2), 3-12.

24. Bilgin, O., Korkut, K. Z. (2005). Determination of Some Bread Quality and Grain Yield Characters in Bread Wheat (*Triticum aestivum L.*). *International Journal of Agriculture and Biology*, 7 (1): 125-128.
25. Blumenthal, J. M., Baltensperger, D. D., Cassman, K. G., Mason, S. C., & Pavlista, A. D. (2008). Importance and effect of nitrogen on crop quality and health. In *Nitrogen in the Environment* (pp. 51-70). Academic Press.
26. Brezolin, A. P.; Silva, J. A. G. da; Frantz, F. C. R.; Binelo, M. O.; Valdiero, A. C.; Zimmer, C. M.; Mantai, R. D.; Marolli, A.; Scremin, O. S.; Mazurkiewicz, G. (2016). The efficiency of wheat yields by nitrogen dose and fractionation. *African Journal of Agricultural Research*, v.11, p.3440-3449. <https://doi.org/10.5897/AJAR2016.11249>
27. Brooker, D. B., Bakker-Arkema, F. W., & Hall, C. W. (1992). Drying and storage of grains and oilseeds. *Springer Science & Business Media*.
28. Brown, B., Westcott, M., Christensen, N., Pan, B., Stark, J., (2005). *Nitrogen Management for Hard Wheat Protein Enhancement*. University of Idaho Extension, PNW 578.
29. Bulut, S. (2012). Ekmeklik buğdayda kalite. *Erciyes Üniversitesi Fen Bilimleri Enstitüsü Fen Bilimleri Dergisi*, 28 (5), 441-446
30. Burkart, M.R., Stoner, J.D., (2008). Chapter 7 – Nitrogen in Groundwater Associated with Agricultural Systems. *Nitrogen in the Environment (Second Edition)*, Academic Press, Pages 177-202, ISBN 9780123743473
31. Bushuk W., Briggs K.G. and Shebeski L.H. (1996). Protein Quantity and Quality as Factors in the Evaluation of Bread Wheats. *Canadian Journal of Plant Science*, Volume 49, Number 2, 113-122.
32. Carson, G.R., Edwards, N.M., (2009). Criteria of wheat and flour quality. In: Khan, K., Shewry, P.R. (Eds.), *Wheat: Chemistry and Technology*, fourth ed. AACC International, Inc., St. Paul, MN, USA, pp. 97e118.
33. Çetin, Ö., Uygan, D., Boyacı, H., Öretir K., (1999). Kılık budayda sulama-azot ve bazı önemli iklim özellikleri arasındaki ilikiler. Türkiye 3. Tarla Bitkileri Kongresi, 15-18 Kasım 1999, Cilt I, Genel ve Tahıllar, 151-156, Adana.
34. Chen, X., Zhou, J., Wang, X., Blackmer, A. M., & Zhang, F. (2004). Optimal rates of nitrogen fertilization for a winter wheat-corn cropping system in Northern China. *Communications in Soil Science and Plant Analysis*, 35(3-4), 583-597.
35. Cui, Z., Zhang, F., Chen, X., Dou, Z., & Li, J. (2010). In-season nitrogen management strategy for winter wheat: Maximizing yields, minimizing environmental impact in an

- over-fertilization context. *Field Crops Research*, 116, 140-146., ISSN 0378-4290, <https://doi.org/10.1016/j.fcr.2009.12.004>.
36. Csaba Horváth, - PhD thesis, SZIU, Gödöllő, (2016). Impact of nitrogen supply on the performance of quality parameters and protein compounds of five winter wheat (*Triticum aestivum L.*) varieties.
37. Decker, W. L.: (1994). Developments in Agricultural Meteorology as a Guide to its Potential for the Twenty-First Century, *Agric. For. Meteorol.* 69, 9–25.
38. Denčić, S., Mladenov, N., Kobiljski, B. (2012). Effects of genotype and environment on breadmaking quality in wheat. *International Journal of Plant Production*, 5(1), 71-82. doi: 10.22069/ijpp.2012.721
39. Djaman, K., Mel, V. C., Ametonou, F. Y., El-Namaky, R., Diallo, M. D., & Koudahe, K. (2018). Effect of nitrogen fertilizer dose and application timing on yield and nitrogen use efficiency of irrigated hybrid rice under semi-arid conditions.
40. Drezner, G., Dvojkovic, K., Horvat, D. et al. (2007). Environmental Impacts on Wheat Agronomic and Quality Traits. *Cereal Research Communications* 35, 357–360. <https://doi.org/10.1556/CRC.35.2007.2.48>
41. Dubetz, S., Gardiner, E.E., Flynn, D., and Dela Roche, A. Ian (1979). Effect of nitrogen fertilizer on nitrogen fractions and amino acid composition on spring wheat. *Can. J. Plant Sci.* 59: 299-305.
42. Edwards, N.M., Gianibelli, M.C., McCaig, T.N., Clarke, J.M., Ames, N.P., Larroque, O.R., and Dexter, J.E., (2007). Relationships between dough strength, polymeric protein quantity and composition for diverse durum wheat genotypes. *Journal of Cereal Science*, 45, 140-149.
43. Ercan, R, Bildik, E. (1993). Azotlu Gübre Uygulamasının Ekmeklik Buğday Kalitesine Etkisi. *Gıda*, 18(3). Retrieved from <https://dergipark.org.tr/en/pub/gida/issue/6971/92918>
44. Erekul, O., Yiğit, A., Koca, Y. O., Ellmer, F., & Wei, K. (2016). Bazı ekmeklik buğday (*Triticum aestivum L.*) çeşitlerinin kalite potansiyelleri ve beslenme fizyolojisi açısından önemi.” *Tarla Bitkileri Merkez Araştırma Enstitüsü Dergisi*, 25(ÖZEL SAYI-1), 31-36.
45. Ertugay Z. (1982). Buğday un ve ekmek arasındaki kalite ilişkileri. *Atatürk Üniv. Ziraat Fak. Ziraat Dergisi*, 13(1-2): 165 176.

46. Eser A. – Kassai M.K. – Tarnawa Á. – Nárai H.F. – Jolánkai M. (2017): Impact of crop year and nitrogen topdressing on the quantity and quality of wheat yield. *Columella 4.1. Suppl.* 157-162 pp.
47. Eser, D., Adak, M. S., and Biesantz, A. (1999). Orta Anadolu'nun kuru tarım bölgelerinde nadas, kışlık mercimek, azotlu gübreleme ve farklı toprak iş lemenin buğday verimine etkileri. *Turkish J. of Agriculture and Forestry*, 23(ek sayı 3) 567-576.
48. Fao, Ifad, Unicef, Wfp & Who. (2017). Building Resilience for Peace and Food Security (Fao, Rome, 2017).
49. Fao, (2018). *Food and Agriculture Organization of the United Nations, Land Resources*. FAOSTATS-Crops. <http://faostat.fao.org/site/567/default.aspx#ancor> (last accessed: 15. October 2020)
50. Fao. (2021). World cereal production. Crop Prospects and Food Situation. *Quarterly Global Report No. 1*, March 2021. Rome. <https://doi.org/10.4060/cb3672en>
51. Fao (2022). Food and Agriculture Organization of the United Nations, World Food Situation. <https://www.fao.org/worldfoodsituation/csdb/en/>
52. Ferrari, M.; Nardino, M.; Carvalho, I. R.; Szarecki, V. J.; Pelegrin, A. J. de; Souza, V. Q. de. (2016). Manejos e fontes de nitrogênio nos componentes de afilamento de trigo. *Revista Brasileira de Ciências Agrárias*, v.11, p.178-185. <https://doi.org/10.5039/agraria.v11i3a5379>
53. Fowler, D. B., Brydon, J., Darroch, B. A., Entz, M. H., & Johnston, A. M. (1990). Environment and genotype influence on grain protein concentration of wheat and rye. *Agronomy Journal*, 82(4), 655-664.
54. Gabriel, D., Pfitzner, C., Haase, N. U., Hüskén, A., Prüfer, H., Greef, J.-M., & Rühl, G. (2017). New strategies for a reliable assessment of baking quality of wheat – Rethinking the current indicator protein content. *Journal of Cereal Science*, 77, 126–134. doi:10.1016/j.jcs.2017.08.002
55. Gadisa, A. (2019). Review on the effect of seed source and size on grain yield of bread wheat (*Triticum aestivum* L.). *J Ecol & Nat Resour*, 3(1), 000155.
56. Gaju, O., Allard, V., Martre, P., Snape, J., Heumez, E., LeGouis, J., Moreau, D., Bogard, M., Griffiths, S. and Orford, S. (2011). Identification of traits to improve the nitrogen-use efficiency of wheat genotypes. *Field Crops Research*, 123 (2), 139-152.

57. Gallagher, L., V.V., Soliman, K. M., Rains, D. W., Qualset, C. O., and Huffaker, R. C. (1983). Nitrogen assimilation in common wheats differing in potential nitrate reductase activity and tissue nitrate concentrations. *Crop Science*, 23: 913-919.
58. Gauer, L. E., Grant, C. A., Gehl, T. D., and Bailey, L. D. (1992). Effect of nitrogen fertilization on grain protein content, nitrogen uptake, and nitrogen use efficiency of six spring wheat (*Triticum aestivum L.*) cultivars, in relation to estimated moisture supply. *Canadian. J. Plant Sci.*, 72: 235-241.
59. Gaur, A.; Kumar, A.; Kiran, R.; Kumari, P. (2020). Importance of seed-borne diseases of agricultural crops: Economic losses and impact on society. In *Seed-Borne Diseases of Agricultural Crops: Detection, Diagnosis & Management*; Springer: Singapore. pp. 3–23.
60. Gelinas P, Morin C, Reid JF, Lachance P (2009). Wheat cultivars grown under organic agriculture and the bread making performance of stone-ground whole wheat flour. *Int J Food Sci Tech* 44: 525–530.
61. Godfrey D, Hawkesford MJ, Powers SJ, Millar S, Shewry PR. (2010). Effects of crop nutrition on wheat grain composition and end use quality. *J Agric Food Chem*. 2010 Mar 10;58(5):3012-21. doi: 10.1021/jf9040645. PMID: 20131902.
62. Gooding, M. J., & Davies, W. P. (1997). Wheat production and utilization: systems, quality and the environment. *CAB international*. Wallingford, UK ISBN 0 85199 155 6. *The Journal of Agricultural Science*, 131(2), 245-249. doi: 10.1017/S0021859698225792
63. Gooding, M.J., and W.P. Davies. (1992). Foliar urea fertilization of cereals: A review.” *Fert. Res.* 32:209–222
64. Goos, R.J., Westfall, D.G., Ludwick, A.E. Goris, J.E., (1982). Grain Protein Content as an Indicator of N Sufficiency for Winter Wheat. *Agronomy Journal*, 74: 130-133.
65. Grabowska L. & Koziara W. (2006) The Effect of Nitrogen Dose, Sowing Density and Time of Harvest on Development and Yields of Hemp Cultivar Bialobrzeskie, *Journal of Natural Fibers*, 2:4, 1-17, DOI: 10.1300/J395v02n04_01
66. Grimwade, B., Tatham, A.S., Freedman, R.B., Sheyry, P.R. & Napier, J.A. (1996). Comparison of the expression patterns of wheat gluten proteins and proteins involved in the secretory pathway in developing caryopses of wheat. *Plant Mol. Biol.*, 30, 1067–1073.

67. Gündoğmuş, E. (1998). Ankara ili Akyurt ilçesi tarım işletmelerinde ekmeklik buğday (*Triticum aestivum L.*) üretiminin fonksiyonel analizi ve üretim maliyetinin hesaplanması. *Tr. J. of Agriculture and Forestry*, 22, 251-260.
68. Györi, Z. (2006). A trágyázás hatása az őszi búza minőségére. (Impacts of fertilizer application on winter wheat quality). *Agrofórum*, 17(9), 14–16.
69. Hassoon, W.H.; Dziki, D.; Miś, A., Biernacka, B. (2021). Wheat Grinding Process with Low Moisture Content: A New Approach for Wholemeal Flour Production. *Processes*. 9, 32. <https://doi.org/10.3390/pr9010032>
70. Hagin J. and Tucker B. (1982). Fertilization of dryland and irrigated soils. *Adv Ser Agric Sci 12*, Springer Verlag, pp. 188
71. Hallikeri, S. S., Halemani, H. L., Patil, V. C., Palled, Y. B., Patil, B. C., & Katageri, I. S. (2010). Effect of nitrogen levels, split application of nitrogen and de-topping on seed cotton yield and fiber quality in Bt-cotton. *Karnataka J. Agric. Sci*, 23(3), 418-422.
72. Hamlin, H.J., (2006). Nitrate toxicity in Siberian sturgeon (*Acipenser baeri*). *Aquaculture*. 253, 688-693
73. Hammad, H. M., Ahmad, A., Wajid, A., & Akhter, J. A. V. A. I. D. (2011). Maize response to time and rate of nitrogen application. *Pak. J. Bot*, 43(4), 1935-1942.
74. Hooser K. and Schafer B., (1969). Yield and quality in multifactorial winter wheat national variety trials, 1963-1966. *Bayer Landw. Ib* 46, No. 2, 191-201.
75. Horváth, Cs., Kis, J., Tarnawa, Á., Kassai, K., Nyárai, H. & Jolánkai, M. (2014): The effect of nitrogen fertilization and crop year precipitation on the protein and wet gluten content of wheat (*Triticum aestivum L.*) grain. *Agrokem. Talajtan*, 63(1), 159-164. <https://akjournals.com/view/journals/0088/63/1/article-p159.xml>
76. Hossein AF, Payam M, Kasra M (2011). Effect of seed size on seedling production in wheat. *Adv Environ Biol* 5(7): 1711-1715. 28. S
77. Hussain, I., Khan, M.A. & Khan, E.A. (2006). Bread wheat varieties as influenced by different nitrogen levels. *J. Zhejiang Univ. - Sci. B* 7, 70–78. <https://doi.org/10.1631/jzus.2006.B0070>
78. Iso (2019). Cereals — determination of bulk density, called mass per hectoliter.” ISO 7971-3:2019
79. Jia, Y. Q., Fabre, J. L., & Aussenac, T. (1996). Effects of growing location on response of protein polymerization to increased nitrogen fertilization for the common

- wheat cultivar Soissons: relationship with some aspects of the breadmaking quality. *Cereal Chemistry*, 73(5), 526-532.
80. Johansson, E.; Branlard, G.; Cuniberti, M.; Flagella, Z.; Hüsken, A.; Nurit, E.; Peña, R.J.; Sissons, M.; Vazquez, D. (2020). Genotypic and Environmental Effects on Wheat Technological and Nutritional Quality. In: Igrejas, G., Ikeda, T., Guzmán, C. (eds) *Wheat Quality For Improving Processing And Human Health*. Springer, Cham. https://doi.org/10.1007/978-3-030-34163-3_8
 81. Johansson, E., Prieto-Linde, M. L., & Jönsson, J. Ö. (2001). Effects of wheat cultivar and nitrogen application on storage protein composition and breadmaking quality. *Cereal Chemistry*, 78(1), 19-25.
 82. Johnson, V. A., Dreier, A. F., and Granbouski, P. H. (1973). Yield and protein responses to nitrogen fertilizer of two winter wheat varieties differing in inherent protein content of their grain. *Agron. J.* 65: 259-263.
 83. Jolánkai M (1985). Differences in fertilizer response due to winter wheat varieties. *Agrokémia és Talajtan* 34:(Supplement) 57-60.
 84. Jolánkai M, Tarnawa Á, Horváth C, Nyárai FH, Kassai K. (2016). Impact of climatic factors on yield quantity and quality of grain crops." *Quarterly J Hungarian Meteorological Service*. 2016; 120(1):73-84.
 85. Joshi, A., Gupta, J. K., Choudhary, S. K., & Paliwal, D. K. (2014). Efficiency of different nitrogen source, doses and split application on growth and yield of maize (*Zea mays* L.) in the Malwa region of Madhya Pradesh. *IOSR J. Agric. Veter. Sci*, 7(2), 2319-2372.
 86. Jung, H.; Lee, Y.J.; Yoon, W.B. (2018). Effect of Moisture Content on the Grinding Process and Powder Properties in Food: A Review. 6, 69. <https://doi.org/10.3390/pr6060069>
 87. Kalkuhl, M., von Braun, J. & Torero, M. (eds). (2016). Food Price Volatility and Its Implications for Food Security and Policy," 3–31. *Springer*, Cham, Switzerland.
 88. Kent, N. L. (1984). *Technology of Cereals*, 3rd ed., Pergamon Press, Oxford.
 89. Khalilzadeh, G., Arshad, Y., Rezaei, M., & Eivazi, A. (2012). Evaluation of Yield, Yield Components, Nitrogen Uptake and Use Efficiency in Bread Wheat (*Triticum aestivum* L.) Cultivars. *Journal of Research in Crop Sciences*, 4(14), 121-138. <https://www.sid.ir/en/journal/ViewPaper.aspx?id=318686>
 90. Khanzada, A. N. A. B., Ansari, M. A., Chang, B. H., Rajput, A. M. M. A. R. A., Magsi, F. H., & Ali, U. M. E. D. (2016). Evaluating right timing and splitting nitrogen

- application rates for enhanced growth and yield of sunflower. *Euro Academ. Res*, 4(7), 5986-6007.
91. Kismányoky T., Tóth Z. (2010). Effect of mineral and organic fertilization on soil fertility as well as on the biomass production and N utilization of winter wheat (*Triticum aestivum L.*) in a long-term cereal crop rotation experiment (IOSDV). *Arch Agron Soil Sci* 56:473–479. <https://doi.org/10.1080/03650340903296819>
 92. Koochekzadeh, A., Fathi, G., Gharineh, M. H., Siadat, S. A., Jafari, S., & Alami-Saeid, K. H. (2009). Impacts of rate and split application of N fertilizer on sugarcane quality. *International Journal of Agricultural Research*, 4(3), 116-123.
 93. Kumar D., Seth R. (2004). Seed yield response of fodder cowpea (*Vigna unguiculata (L.) Walp*) varieties to varying seed rate and seed size. *Seed Res* 32: 149- 153.
 94. Langenkämper G., Zörb C., (2019). Modern aspects of wheat grain proteins.” *Journal of Applied Botany and Food Quality* 92, 240-245 (2019), DOI:10.5073/JABFQ.2019.092.033
 95. Large, E.G., (1954). Growth Stages in Cereals: Illustration of the Feeke's scale. *Pl. Path.*, 3, 128–129.
 96. Liu, H., Wang, Z., Yu, R., Li, F., Li, K., Cao, H., Yang, N., Li, M., Dai, J., Zan, Y., Li, Q., Xue, C., He, G., Huang, D., Huang, M., Liu, J., Qiu, W., Zhao, H., & Mao, H. (2016). Optimal nitrogen input for higher efficiency and lower environmental impacts of winter wheat production in China. *Agriculture, Ecosystems & Environment*, 224, 1-11.
 97. Lloveras, J., Lopez, A., Ferrán, J., Espachs, S., Solsova, J., (2001). Breadmaking quality of wheat and soil nitrate as affected by nitrogen fertilization in irrigated Mediterranean conditions. *Agron. J.* 93: 1183-1190
 98. Lopez-Bellido L., Lopez-Bellido R J., Redondo R. (2005). Nitrogen efficiency in wheat under rainfed Mediterranean conditions as affected by split nitrogen application *Field Crops Research*, 94, pp. 86-97
 99. Lošák, T., & Richter, R. (2004). Split nitrogen doses and their efficiency in poppy (*Papaver somniferum L.*) nutrition. *Growth*, 600, 49.
 100. Lv, Z., Liu, X., Cao, W., & Zhu, Y. (2013). Climate change impacts on regional winter wheat production in main wheat production regions of China. *Agricultural and forest meteorology*, 171, 234-248.
 101. M.N. Khan, M. Mobin, Z.K. Abbas, S.A. Alamri, (2018). Fertilizers and Their Contaminants in Soils, Surface and Groundwater, *Encyclopedia of the Anthropocene*,

Elsevier, Pages 225-240, ISBN 9780128135761, <https://doi.org/10.1016/B978-0-12-809665-9.09888-8>.

102. Ma, W., Yu, Z., She, M., Zhao, Y., & Islam, S. (2019). Wheat gluten protein and its impacts on wheat processing quality. *Frontiers of Agricultural Science and Engineering*.
103. Madan, H. S., & Munjal, R. (2009). Effect of split doses of nitrogen and seed rate on protein content, protein fractions and yield of wheat. *Journal of Agriculture and Biological Science*, 4(1), 26-31.
104. Mandic, V., Krnjaja, V., Tomic, Z., Bijelic, Z., Simic, A., Ruzic Muslic, D., & Gogic, M. (2015). Nitrogen fertilizer influence on wheat yield and use efficiency under different environmental conditions. *Chilean journal of agricultural research*, 75(1), 92-97.
105. Mantai, R. D., Gonzalez da Silva, J. A., Arenhardt, E. G., Heck, T. G., Sausen, A. T. Z. R., Krüger, C. A. M. B., Cardoso, A. M., Neto, C. J. G., & Kryszun, D. K. (2015). The effect of nitrogen dose on the yield indicators of oats. *African Journal of Agricultural Research*, 10(39), 3773-3781.
106. McDonald, G. K., & Hussein, S. (2017). The effects of wheat seed quality are greatest in high yielding environments." In "Doing More with Less, *Proceedings of the 18th Australian Agronomy Conference 2017*, Ballarat, Victoria, Australia, 24-28 September 2017 (pp. 1-4). Australian Society of Agronomy Inc.
107. Miezian, K., Heyne, E. G., & Finney, K. F. (1977). Genetic and Environmental Effects on the Grain Protein Content in Wheat 1. *Crop Science*, 17(4), 591-593.
108. Mughal MH. (2019). Wheat compounds – A comprehensive review. *Integr Food Nutr Metab* 6: DOI: 10.15761/IFNM.1000270
109. Mut, Z., Sezer, I., & Gulumser, A. (2005). Effect of different sowing rates and nitrogen levels on grain yield, yield components and some quality traits of triticale. *Asian Journal of Plant Sciences*. ISSN: 1812-5697. p.533-539.
110. Nasa. (N.D.) Expert Group- Growing Wheat. https://gpm.nasa.gov/education/sites/default/files/lesson_plan_files/water-for-wheaties/PR_AG_HS_GrowingWheat.pdf
111. Nass, H., MacLeod, J., and Suzuki, M., (1976). Effects of Nitrogen application on yield, plant characters, and N levels in grain of six spring wheat cultivars 1. *Crop science*, 16 (6), 877- 879.

112. Nguyen, L.T.T., Osanai, Y., Lai, K., Anderson, I.C., Bange, M. P., Tissue, D.T., and Singh, B.K. (2018). Responses of the soil microbial community to nitrogen fertilizer regimes and historical exposure to extreme weather events: flooding or prolonged drought. *Soil Biol Biochem* 118: 227–236.
<https://doi.org/10.1016/j.soilbio.2017.12.016>.
113. Olesen, J. E., Jensen, T., & Petersen, J. (2000). Sensitivity of field-scale winter wheat production in Denmark to climate variability and climate change. *Climate Research*, 15(3), 221-238.
114. Orloff, S., Wright, S., & Ottman, M. (2012). Nitrogen Management Impacts on Wheat Yield and Protein. In Proceedings, California Alfalfa & Grains Symposium, Sacramento, CA, December, 11-12.
115. Ortiz, R., Sayre, K. D., Govaerts, B., Gupta, R., Subbarao, G. V., Ban, T., ... & Reynolds, M. (2008). Climate change: can wheat beat the heat? *Agriculture, Ecosystems & Environment*, 126(1-2), 46-58.
116. Osborne, T.B., (1907). *The proteins of the wheat kernel*. Carnegie Institution, Washington. DOI: 10.5962/bhl.title.26152
117. Özden A. B. S., (2012). Role of nitrogen nutrition in biofortification of durum wheat with iron.” Istanbul, TR: Sabancı University
118. Öztürk, İ., Gökkuş, A. (2008). Azotla gübrelemenin bazı ekmeklik buğday çeşitlerinin verimi ve kalitesine etkileri. *Ankara Üniversitesi Ziraat Fakültesi Tarım Bilimleri Dergisi*, 14(4): 334-340.
119. Panozzo J.F. and Eagles H.A., (2000). Cultivar and environmental effects on quality characters in wheat.” *Australian Journal of Agricultural Research*, 51: 629-636
120. Paredes-López, O., Covarrubias-Alvarez, M. M. and Barquín-Carmona J. (1985). Influence of nitrogen fertilization on the physico-chemical and functional properties of bread wheats. *Cereal Chem.*, 62, 427–430.
121. Pepó, P. (2010). Adaptive capacity of wheat (*Triticum aestivum* L.) and maize (*Zea mays* L.) crop models to ecological conditions.” *Növénytermelés*, 59. Suppl., 325–328.
122. Peterson, C. J., Graybosch, R. A., Baenziger, P. S., & Grombacher, A. W. (1992). Genotype and environment effects on quality characteristics of hard red winter wheat. *Crop Science*, 32(1), 98-103.
123. Pollhamer, E. (1981): “A búza és a liszt minősége. (Quality of wheat and flour).” *Mezőgazdasági Kiadó*. Budapest. 203 pages.

124. Posner, E. S., & Hibbs, A. N. (2005). Wheat flour milling (No. Ed. 2). *American Association of Cereal Chemists, Inc.*
125. Posner, E.S. (2003). Principles of milling. *Encyclopedia of food science, food technology and nutrition*. Academic Press, Harcourt Brace Jovanovich Publishers. London, 3980–3986.
126. Quail K. (2018). The importance of grain moisture to the quality of end products. Australian Export Gains Innovation Centre. GroundCover Issue 132. GRDC Project Code: AEG00006
127. Quievreux, D., (1997). Réseau d’observation de parcelles de céréales.” *Résultats de la Récolte, 1997*, Chambre d’Agriculture de la Somme, Amiens, France
128. Rahman, M., Sarker, M., Amin, M., Jahan, A., & Akhter, M. (1) (2011). Yield Response and Nitrogen Use Efficiency of Wheat Under Different Doses and Split Application of Nitrogen Fertilizer. *Bangladesh Journal of Agricultural Research*, 36(2), 231-240. <https://doi.org/10.3329/bjar.v36i2.9249>
129. Rakszegi, M., Mikó, P., Löschenberger, F., Hiltbrunner, J., Aebi, R. ... & Bedő, Z. (2016). Comparison of quality parameters of wheat varieties with different breeding origin under organic and low-input conventional conditions. *J. Cereal Sci.*, 69.297–305.
130. Reddecliffe, T.M., Moot, D.J., Wilson, D.R. and Scott, W.R. (2000). Grain yield and quality of two durum wheat cultivars grown in Canterbury. *Proceedings of the Thirtieth Annual Conference, Agronomy Society of New Zealand, Palmerston North, New Zealand*. Agronomy-New-Zealand. 30: 77-82.
131. Rehman R., (2019). Uptake, transport and seed deposition of zinc in wheat and maize under varied zinc and nitrogen supply. Istanbul, TR: Sabancı University
132. Robinson, H.C., Lawler, M.R., Chenoweth, W.L. and Garwick, A.E. (1986). *Proteins and Amino Acids: 44-63, in Normal and Therapeutic Nutrition*. McMillan Publishing Company, New York.
133. Sathiya, K., & Ramesh, T. (2009). Effect of split application of nitrogen on growth and yield of aerobic rice. *Asian J. Exp. Sci*, 23(1), 303-306.
134. Schofield, J. D. and Greenwell, P. (1987). Wheat starch granule proteins and their technological significance. *Cereals in a European Context*, (ed. I. D. Morton) VCH Verlagsgesellschaft mbH, Weinheim and Ellis Horwood, Chichester, pp. 407-420.

135. Shaiful I. M., Hasanuzzaman, M., Rokonuzzaman, M., Nahar, K. (2012). Effect of split application of nitrogen fertilizer on morpho-physiological parameters of rice genotypes. *International Journal of Plant Production*, 3(1), 51-62. doi: 10.22069/ijpp.2012.631
- Shewry, P. & Halford, N. (2002): "Cereal seed storage proteins: structures, properties and role in grain utilization." *J. Exp. Bot.*, 53(370), 947–958.
136. Shewry, P. R., (2009). *Wheat, Journal of Experimental Botany*, Volume 60, Issue 6, April 2009, pp 1537–1553, <https://doi.org/10.1093/jxb/erp058>
137. Simmone R, Steege HT, Werger M (2000). Survival and growth in gaps: A case study for tree seedlings of 8 species in the Guyanese tropical rainforest in seed seedlings and gap size matters. *Tropenbos-Guyana Programmes*, Guyana.
138. Singandhupe, R. B., Rao, G. G. S. N., Patil, N. G., & Brahmanand, P. S. (2003). Fertilization studies and irrigation scheduling in drip irrigation system in tomato crop (*Lycopersicon esculentum* L.). *European Journal of Agronomy*, 19(2), 327-340.
139. Singh, H., & Singh, G. (2013a). Effect of potassium and split application of nitrogen on yield attributes and yield of soybean [*Glycine max* (L.) Merrill]. *Agricultural Science Digest*, 33(4).
140. Singh, J., Mahal, S. S., & Singh, A. (2013b). Productivity and quality of malt Barley (*Hordeum vulgare*) as affected by sowing date, rate and stage of nitrogen application. *Indian Journal of Agronomy*, 58(1), 72-80.
141. Singh, R., La, M., Singh, G., & Kumar, T. (2019). Effect of nitrogen and phosphorus on growth parameter and yield of canola (*Brassica napus* L.). *J. Pharmacogn Phytochem*, 8, 380-384.
142. Slafer, G.A., Satorre, E.H., Andrade, F.H., (1994). Increases in grain yield in bread wheat and associated physiological changes. *Improvement of Field Crops* (G.A. Slafer, ed.). Marcel Dekker: 1-68.
143. Sohail, M., Hussain, I., Tanveer, S. K., Abbas, S. H., Qamar, M., Ahmed, M. S., Waqar, S. (2018). Effect of Nitrogen Fertilizer Application Methods on Wheat Yield and Quality. *Science, Technology and Development*, 37 (2): 89-92.
144. Šramková, Z., Gregová, E., Šturdík, E. (2009). Chemical Composition and Nutritional Quality of Wheat Grain. *Acta Chimica Slovaca*, 2 (1): 115- 138.
145. Styk, B. and Dziamba, S. (1984). Effect of some agronomic practices and irrigation on yield variability, 1000-grain weight, hectoliter weight and protein content in two spring wheat cultivars. *Field Crops Abstracts*, Vol. 37: 82.

146. Szentpétery, Zs., Kleinheincz, Cs., Szöllősi, G., & Jolánkai, M. (2005). Effect of nitrogen top-dressing on winter wheat yield, quantity and quality. *Acta Alimentaria*, 34(2), 177-185.
147. Szulc, P., Bocianowski, J., & Nowosad, K. (2022). Evaluation of the effect of nitrogen fertilizers on nitrogen use efficiency in grain maize. *Maydica*, 66(3), 8.
148. Terman, G. L. (1979). Yields and Protein Content of Wheat Grain as Affected by Cultivar, N, and Environmental Growth Factors 1. *Agronomy Journal*, 71(3), 437-440.
149. Thaler, S., Eitzinger, J., Trnka, M., & Dubrovsky, M. (2012). Impacts of climate change and alternative adaptation options on winter wheat yield and water productivity in a dry climate in Central Europe. *The Journal of Agricultural Science*, 150(5), 537.
150. Todab, (n.d.) (2021). Buğday ve Arpa Tarımı. Section 5.3. <https://todab.org.tr/files/BUGDAY%20VE%20ARPA%20TARIMI/B%c3%b61%c3%bcm-5.pdf> Search engine: Google. Key words: ‘buğday azot kullanım etkinliği.’ Date of search: 2021.05.10.
151. Treseder, K.K. (2008). Nitrogen additions and microbial biomass: a meta-analysis of ecosystem studies. *Ecol Lett* 11: 1111–1120. <https://doi.org/10.1111/j.1461-0248.2008.01230.x>
152. Triboi, E., Abad, A., Michelena, A., Lloveras, J., Ollier, J. L., & Daniel, C. (2000). Environmental effects on the quality of two wheat genotypes: 1. Quantitative and qualitative variation of storage proteins. *European Journal of Agronomy*, 13(1), 47-64.
153. T.C. Tarım ve Orman Bakanlığı. (n.d.). Bugday. <https://istanbul.tarimorman.gov.tr/belgeler/kutumenu/brosurler/tarlabitkileri/bugday.pdf>
154. Ünal, S., (2002). Importance of wheat quality and methods in wheat quality determination. *Hububat Ürünleri Teknolojisi Kongre ve Sergisi 3-4 Ekim 2002*, pp 25-37, Gaziantep, Turkey.
155. Van Lill, D., Purchase, J. L., (1995). Directions in breeding for winter wheat yield and quality in South Africa from 1930 to 1990. *Euphytica* 82, 79-87.
156. Várallyay, G. (2008). Extreme soil moisture regime as limiting factor of the plants’ water uptake. *Cereal Res. Commun.*, 36(2), 3–6.

157. Varga B., Svecnjak Z., (2006). The effect of late-season urea spraying on grain yield and quality of winter wheat cultivars under low and high basal nitrogen fertilization. *Field Crops Research* 96:125-132.
158. Vázquez, D., Berger, A. G., Cuniberti, M., Bainotti, C., de Miranda, M. Z., Scheeren, P. L., ... Peña, R. J. (2012). Influence of cultivar and environment on quality of Latin American wheats. *Journal of Cereal Science*, 56(2), 196–203.
doi:10.1016/j.jcs.2012.03.004
159. Vida, Gy., Bedő, Z. & Jolánkai, M. (1996). Agronómiai kezeléskombinációk őszi búzafajták sütőipari minőségére gyakorolt hatásának elemzése főkomponens-analízissel. (Investigation of the effect of agronomical treatment combinations on the winter wheat's baking qualities with main component analysis). *Növénytermelés*, 45(6), 453–462.
160. Wallenstein, M.D., McNulty, S., Fernandez, I.J., Boggs, J., and Schlesinger, W.H. (2006). Nitrogen fertilization decreases forest soil fungal and bacterial biomass in three long-term experiments. *For Ecol Manage* 222: 459–468.
<https://doi.org/10.1016/j.foreco.2005.11.002>.
161. Wang, C., Liu, D., and Bai, E. (2018b). Decreasing soil microbial diversity is associated with decreasing microbial biomass under nitrogen addition. *Soil Biol Biochem* 120: 126–133. <https://doi.org/10.1016/j.soilbio.2018.02.003>.
162. Wang, F., Chen, S., Wang, Y., Zhang, Y., Hu, C., and Liu, B. (2018a). Long-term nitrogen fertilization elevates the activity and abundance of nitrifying and denitrifying microbial communities in an upland soil: implications for nitrogen loss from intensive agricultural systems. *Front Microbiol* 9: 2424.
<https://doi.org/10.3389/fmicb.2018.02424>.
163. Wang, H., Zhang, Y., Chen, A., Liu, H., Zhai, L., Lei, B., & Ren, T. (2017). An optimal regional nitrogen application threshold for wheat in the North China Plain considering yield and environmental effects. *Field Crops Research*, 207, 52-61.
164. Washington Wheat Facts 2016/2017. Washington Grain Commission
<http://wagrains.org/wp-content/uploads/2015/04/WGC-2016-17WF4WebFinal.pdf>
(last accessed: 10 October 2019).
165. Wrigley, C., & Batey, I. (2003). Assessing grain quality. *Bread Making*, pp 71–96. doi:10.1533/9781855737129.1.71
166. Wrigley, C.W. (2007). Mitigating the damaging effects of growth and storage conditions on grain quality. In: Buck, H.T., Nisi, J.E., Salomón, N. (Eds.), *Wheat*

- Production in Stress Environments. Developments in Plant Breeding, vol. 12.*
Springer, Dordrecht, The Netherlands, pp. 425e439.
167. Xu, J., Cai, H., Wang, X., Ma, C., Lu, Y., Ding, Y., ... & Saddique, Q. (2020). Exploring optimal irrigation and nitrogen fertilization in a winter wheat-summer maize rotation system for improving crop yield and reducing water and nitrogen leaching. *Agricultural Water Management*, 228, 105904.
168. Xue, C., Auf'm Erley, G. S., Rossmann, A., Schuster, R., Koehler, P., & Mühling, K. H. (2016). Split Nitrogen Application Improves Wheat Baking Quality by Influencing Protein Composition Rather Than Concentration. *Frontiers in plant science*, 7, 738.
<https://doi.org/10.3389/fpls.2016.00738>
169. Xue, C., Matros, A., Mock, H. P., & Mühling, K. H. (2019). Protein Composition and Baking Quality of Wheat Flour as Affected by Split Nitrogen Application. *Frontiers in plant science*, 10, 642.
<https://doi.org/10.3389/fpls.2019.00642>
170. Yara, (2011). Effect of applied nitrogen on winter wheat grain protein.
<https://www.yara.co.uk/crop-nutrition/wheat/increasing-hagberg-falling-number/>
171. Yara, (n.d.) "How to increase wheat protein content."
<https://www.yara.co.uk/crop-nutrition/wheat/increasing-wheat-protein/> Search engine: Google. Key words: 'wheat grain protein.' Date of search: 2021.05.05.
172. Yevdokimov, I., Gatteringer, A., Buegger, F. et al. (2008). Changes in microbial community structure in soil as a result of different amounts of nitrogen fertilization. *Biol Fertile Soils* 44, 1103. <https://doi.org/10.1007/s00374-008-0315-1>
173. Yılmaz, N., Şimşek, S. (2013). Sivas ekolojik koşullarında ekmeklik buğdayda (*Triticum aestivum L.*) üst gübrelemede kullanılacak azotlu gübre form ve miktarının belirlenmesi. *Akademik Ziraat Dergisi*, 1(2), 91-96.
174. Yousaf, M., Fahad, S., Shah, A. N., Shaaban, M., Khan, M. J., Sabiel, S. A. I., ... & Osman, K. A. (2014). The effect of nitrogen application rates and timings of first irrigation on wheat growth and yield. *Int. J. Agric. Innovat. Res*, 2(4), 645-65.
175. Zadoks, J.C., Chang, T.T., Konzak, C.F., (1974). A Decimal Code for Growth Stage of Cereals. *Weed. Res.*, 14: 415-421.
176. Zecevic, V., Knezevic, D., Boskovic, J. Micanovic D. & Dozet G. (2010). Effect of nitrogen fertilization on winter wheat quality. *CEREAL RESEARCH COMMUNICATIONS* 38, 243–249. <https://doi.org/10.1556/CRC.38.2010.2.10>

177. Zende N. B., Sethi H. N., Karunakar. A. P. and Jiotode, D. J. (2005). Effect of sowing time and fertility levels on yield and quality of durum wheat genotypes. *Res. on Crops*. 6(2): 194-196.

12. ACKNOWLEDGEMENT

- I would like to express my profound gratitude to my supervisor Prof. Dr. Marton Jolánkai for his inspiration and continuous guidance from the first day until the end.


- I express my gratitude to the professors and staff members of Department of Crop Production and Horticultural Sciences who have willingly helped me out with their kindness.

- I am thankful to the Tempus Public Foundation for providing a Stipendium Hungaricum scholarship and study opportunity at the Hungarian University of Agriculture and Life Sciences (formerly Szent Istvan University), Hungary.


- I express deep and sincere gratitude to members of my family; without their blessing and inspiration, this study would not have been possible.

13. APPENDIX 1 – TABLES AND FIGURES


12.1. Test Weight




24. Figure Impact of N topdressing applications on wheat grain test weight. 2015-2016 (Nagyombos, Hungary)



25. Figure Impact of N topdressing applications on wheat grain test weight. 2016-2017 (Nagygyombos, Hungary)




26. Figure Impact of N topdressing applications on wheat grain test weight. 2018-2019 (Gödöllő, Hungary) 2




27. Figure Impact of N topdressing applications on wheat grain test weight. 2018–2019 (Gödöllő, Hungary) Box Plot Alföld and MV Karéj


12.2. Thousand Kernel Weight




28. Figure Impact of N topdressing applications on thousand kernel weight of wheat varieties. 2015-2016 (Nagygyombos, Hungary)



29. Figure Impact of N topdressing applications on thousand kernel weight of wheat varieties. 2016-2017 (Nagygyombos, Hungary)



30. Figure Impact of N topdressing applications on thousand kernel weight of wheat varieties. 2018-2019 (Gödöllő, Hungary) 2



31. Figure Impact of N topdressing applications on thousand kernel weight of wheat varieties. 2018-2019 (Gödöllő, Hungary) 3

12.3. The Grain Moisture

| Moisture content % 2015-2016 | | | | | |
|-------------------------------------|---------------|-----------------|-----------------|-------------------|-----------------|
| N | Alföld | MV Nádor | MV Karéj | MV Toborzó | MV Toldi |
| 0 | 6,50 | 6,90 | 9,80 | 7,30 | 10,40 |
| 80 | 6,50 | 7,10 | 9,70 | 7,70 | 10,60 |
| 80+40 | 6,30 | 6,60 | 9,70 | 7,20 | 10,60 |
| 120 | 6,50 | 6,90 | 9,90 | 7,20 | 10,50 |
| 120+40 | 6,70 | 7,00 | 9,70 | 7,10 | 10,60 |
| 160 | 6,50 | 6,90 | 9,80 | 7,30 | 10,40 |

31. Table Impact of N topdressing applications on moisture content of wheat varieties. 2015-2016 (Nagygyombos, Hungary)


| Moisture content % 2016-2017 | | | | | |
|-------------------------------------|---------------|-----------------|-----------------|-------------------|-----------------|
| N | Alföld | MV Nádor | MV Karéj | MV Toborzó | MV Toldi |
| 0 | 9,87 | 9,83 | 10,30 | 8,65 | 10,22 |
| 80 | 9,73 | 9,84 | 10,30 | 8,80 | 10,32 |
| 80+40 | 9,50 | 9,88 | 10,30 | 8,70 | 10,23 |
| 120 | 9,75 | 9,83 | 10,20 | 8,75 | 10,36 |
| 120+40 | 9,80 | 9,85 | 10,20 | 8,69 | 10,40 |
| 160 | 9,73 | 9,79 | 10,10 | 8,78 | 10,44 |

32. Table Impact of N topdressing applications on moisture content of wheat varieties. 2016-2017 (Nagygyombos, Hungary)

| Moisture content % 2018-2019 | | |
|-------------------------------------|---------------|-----------------|
| N | Alföld | MV Karéj |
| 0 | 15,03 | 11,77 |
| 80 | 15,30 | 11,80 |
| 80+40 | 15,33 | 11,87 |
| 120 | 14,97 | 11,83 |
| 120+40 | 15,13 | 11,80 |

| | | |
|------------|-------|-------|
| 160 | 15,03 | 11,90 |
|------------|-------|-------|

33. Table Impact of N topdressing applications on moisture content of wheat varieties. 2018-2019 (Gödöllő, Hungary) 2



32. Figure Impact of N topdressing applications on moisture content of wheat varieties. 2018-2019 (Gödöllő, Hungary) 3

12.4. Protein Content


| Protein content % 2015-2016 | | | | | |
|-----------------------------|--------|----------|----------|------------|----------|
| N | Alföld | MV Nádor | MV Karéj | MV Toborzó | MV Toldi |
| 0 | 13,60 | 12,00 | 12,40 | 13,80 | 12,70 |
| 80 | 14,50 | 12,60 | 13,30 | 13,80 | 13,30 |
| 80+40 | 15,70 | 13,60 | 14,40 | 14,90 | 13,90 |
| 120 | 15,30 | 13,50 | 13,90 | 14,60 | 13,80 |

| | | | | | |
|---------------|-------|-------|-------|-------|-------|
| 120+40 | 16,00 | 13,90 | 14,90 | 15,50 | 14,30 |
| 160 | 15,80 | 14,20 | 14,60 | 15,30 | 14,00 |


34. Table Impact of N topdressing applications on protein content of wheat varieties. 2015-2016 (Nagygombos, Hungary)

| Protein content % 2016-2017 | | | | | |
|-----------------------------|--------|----------|----------|------------|----------|
| N | Alföld | MV Nádor | MV Karéj | MV Toborzó | MV Toldi |
| 0 | 13,13 | 11,12 | 9,90 | 10,09 | 11,65 |
| 80 | 13,98 | 12,36 | 11,80 | 12,03 | 13,50 |
| 80+40 | 14,68 | 12,33 | 12,30 | 12,09 | 13,22 |
| 12 | 15,70 | 13,07 | 12,90 | 12,77 | 13,62 |
| 120+40 | 14,87 | 13,32 | 13,30 | 13,04 | 14,06 |
| 160 | 15,99 | 13,73 | 14,10 | 13,60 | 14,73 |


35. Table Impact of N topdressing applications on protein content of wheat varieties. 2016-2017 (Nagygombos, Hungary)



33. Figure Impact of N topdressing applications on protein content of wheat varieties. 2018-2019 (Gödöllő, Hungary) 2




34. Figure Impact of N topdressing applications on protein content in Alföld. 2015-2016 and 2016-2017 (Nagygyombos, Hungary) and 2018-2019 (Gödöllő, Hungary) combined.




35. Figure Impact of N topdressing applications on protein content of MV Karéj. 2015-2016 and 2016–2017 (Nagygombos, Hungary) and 2018-2019 (Gödöllő, Hungary) combined.


12.5. Gluten Content




36. Figure Impact of N topdressing applications on gluten content of wheat varieties. 2015-2016 (Nagyombos, Hungary)




37. Figure Impact of N topdressing applications on gluten content of wheat varieties. 2016-2017 (Nagyombos, Hungary)



38. *Figure Impact of N topdressing applications on gluten content of wheat varieties. 2018-2019 (Gödöllő, Hungary) 2*



39. Figure Impact of N topdressing applications on gluten content of wheat varieties. 2018–2019 (Gödöllő, Hungary) 3



40. Figure Impact of N topdressing applications on gluten content of wheat varieties. 2015-2016 and 2016–2017 (Nagygombos, Hungary) and 2018-2019 (Gödöllő, Hungary) combined.

12.6. Anova-Spss outputs

12.6.1. Examination of Parameters According to Categorical Variables in Different Years

| | | N | M | SD | F | df | p |
|----------------------------|------|----------|----------|-----------|----------|-----------|----------|
| Test Weight (hl) | 2016 | 36 | 78.07 | 0.67 | 86.509 | 2.105 | .000 |
| | 2017 | 36 | 76.42 | 2.94 | | | |
| | 2019 | 36 | 71.33 | 2.52 | | | |
| Thousand Kernel Weight (g) | 2016 | 36 | 43.18 | 2.33 | 3.910 | 2.105 | .023 |
| | 2017 | 36 | 41.20 | 5.39 | | | |
| | 2019 | 36 | 41.07 | 2.07 | | | |
| Falling Number | 2016 | 36 | 365.47 | 56.47 | 42.200 | 2.105 | .000 |
| | 2017 | 36 | 528.32 | 116.18 | | | |
| | 2019 | 36 | 396.30 | 49.66 | | | |
| Protein Content (%) | 2016 | 36 | 14.53 | 1.19 | 8.829 | 2.105 | .000 |
| | 2017 | 36 | 13.47 | 1.82 | | | |
| | 2019 | 36 | 14.93 | 1.48 | | | |
| Moisture Content (%) | 2016 | 36 | 8.12 | 1.67 | 83.241 | 2.105 | .000 |
| | 2017 | 36 | 9.97 | 0.35 | | | |
| | 2019 | 36 | 13.13 | 2.33 | | | |
| Gluten Content (%) | 2016 | 36 | 31.47 | 4.24 | 297.147 | 2.105 | .000 |
| | 2017 | 36 | 29.61 | 5.38 | | | |
| | 2019 | 36 | 10.57 | 1.32 | | | |
| Zeleny Number | 2016 | 36 | 34.11 | 6.94 | 18.886 | 2.105 | .000 |
| | 2017 | 36 | 43.22 | 9.38 | | | |
| | 2019 | 36 | 34.52 | 3.90 | | | |

36. Table Impact of undivided/split N topdressing applications on wheat grain based on years, 2015-2016, 2016-2017 and 2018-2019 crop seasons, Spss-One Way Anova

12.6.2. Examination of Parameters According to Categorical Variables in Different Sites-Hatvan

12.6.2.1. Comparison of the Means of Parameters Measured in Hatvan-Nagygyombos Based on Year

| | | N | M | SD | t | df | p |
|----------------------------|------|----------|----------|-----------|----------|-----------|----------|
| Test Weight (hl) | 2016 | 36 | 78.07 | 0.67 | 3.280 | 38.601 | .002 |
| | 2017 | 36 | 76.42 | 2.94 | | | |
| Thousand Kernel Weight (g) | 2016 | 36 | 43.18 | 2.33 | 2.026 | 47.655 | .048 |
| | 2017 | 36 | 41.20 | 5.39 | | | |

| | | | | | | | |
|----------------------|------|----|--------|--------|--------|--------|------|
| Falling Number | 2016 | 36 | 365.47 | 56.47 | -7.564 | 50.660 | .000 |
| | 2017 | 36 | 528.32 | 116.18 | | | |
| Protein Content (%) | 2016 | 36 | 14.53 | 1.19 | 2.919 | 60.304 | .005 |
| | 2017 | 36 | 13.47 | 1.82 | | | |
| Moisture Content (%) | 2016 | 36 | 8.12 | 1.67 | -6.508 | 38.017 | .000 |
| | 2017 | 36 | 9.97 | 0.35 | | | |
| Gluten Content (%) | 2016 | 36 | 31.47 | 4.24 | 1.635 | 70 | .107 |
| | 2017 | 36 | 29.61 | 5.38 | | | |
| Zeleny Number | 2016 | 36 | 34.11 | 6.94 | -4.682 | 70 | .000 |
| | 2017 | 36 | 43.22 | 9.38 | | | |

37. Table Impact of undivided/split N topdressing applications on wheat grain based on Hatvan-Nagyombos site, 2015-2016 and 2016-2017 crop seasons, Spss-One Way Anova

12.6.2.2. Comparison of the Means of Parameters Measured in Hatvan-Nagyombos Based on Genotype

| | | N | M | SD | t | df | p |
|----------------------------|--------|----|--------|--------|---------|--------|------|
| Test Weight (hl) | Alföld | 36 | 78.83 | 0.83 | 8.278 | 45.180 | .000 |
| | Karéj | 36 | 75.66 | 2.15 | | | |
| Thousand Kernel Weight (g) | Alföld | 36 | 38.75 | 3.03 | -11.836 | 55.378 | .000 |
| | Karéj | 36 | 45.63 | 1.72 | | | |
| Falling Number | Alföld | 36 | 521.01 | 119.56 | 6.447 | 55.885 | .000 |
| | Karéj | 36 | 372.79 | 68.79 | | | |
| Protein Content (%) | Alföld | 36 | 14.85 | 1.26 | 5.159 | 70 | .000 |
| | Karéj | 36 | 13.16 | 1.50 | | | |
| Moisture Content (%) | Alföld | 36 | 8.11 | 1.66 | -6.618 | 38.268 | .000 |
| | Karéj | 36 | 9.98 | 0.36 | | | |
| Gluten Content (%) | Alföld | 36 | 33.41 | 3.66 | 6.117 | 70 | .000 |
| | Karéj | 36 | 27.67 | 4.28 | | | |
| Zeleny Number | Alföld | 36 | 40.08 | 9.58 | 1.286 | 70 | .203 |
| | Karéj | 36 | 37.25 | 9.10 | | | |

38. Table Impact of undivided/split N topdressing applications on wheat grain based on Hatvan-Nagyombos site by Genotype, 2015-2016 and 2016-2017 crop seasons, Spss-One Way Anova

12.6.2.3. Comparison of the Means of Parameters Measured in Hatvan-Nagyombos Based on Treatment

| | | N | M | SD | F | df | p |
|------------------|-------------|----|-------|------|------|------|------|
| Test Weight (hl) | 0 kg/ha | 12 | 77.36 | 2.35 | .124 | 5.66 | .986 |
| | 80 kg/ha | 12 | 77.43 | 2.45 | | | |
| | 80+40 kg/ha | 12 | 77.38 | 2.31 | | | |

| | | | | | | | |
|----------------------------|--------------|----|--------|--------|--------|------|-------|
| | 120 kg/ha | 12 | 76.97 | 2.33 | | | |
| | 120+40 kg/ha | 12 | 77.43 | 2.35 | | | |
| | 160 kg/ha | 12 | 76.91 | 2.29 | | | |
| Thousand Kernel Weight (g) | 0 kg/ha | 12 | 43.22 | 4.23 | .208 | 5.66 | .958 |
| | 80 kg/ha | 12 | 42.16 | 4.12 | | | |
| | 80+40 kg/ha | 12 | 42.23 | 4.43 | | | |
| | 120 kg/ha | 12 | 41.98 | 4.34 | | | |
| | 120+40 kg/ha | 12 | 42.09 | 4.50 | | | |
| | 160 kg/ha | 12 | 41.45 | 4.55 | | | |
| Falling Number | 0 kg/ha | 12 | 389.38 | 86.66 | .813 | 5.66 | .545 |
| | 80 kg/ha | 12 | 437.33 | 110.76 | | | |
| | 80+40 kg/ha | 12 | 446.99 | 123.54 | | | |
| | 120 kg/ha | 12 | 463.99 | 133.08 | | | |
| | 120+40 kg/ha | 12 | 462.00 | 127.34 | | | |
| | 160 kg/ha | 12 | 481.69 | 148.08 | | | |
| Protein Content (%) | 0 kg/ha | 12 | 12.03 | 1.61 | 9.227 | 5.66 | .000 |
| | 80 kg/ha | 12 | 13.39 | 1.23 | | | |
| | 80+40 kg/ha | 12 | 14.26 | 1.49 | | | |
| | 120 kg/ha | 12 | 14.45 | 1.22 | | | |
| | 120+40 kg/ha | 12 | 14.78 | 1.11 | | | |
| | 160 kg/ha | 12 | 15.11 | 0.95 | | | |
| Moisture Content (%) | 0 kg/ha | 12 | 9.12 | 1.63 | .023 | 5.66 | 1.000 |
| | 80 kg/ha | 12 | 9.06 | 1.56 | | | |
| | 80+40 kg/ha | 12 | 8.93 | 1.64 | | | |
| | 120 kg/ha | 12 | 9.08 | 1.58 | | | |
| | 120+40 kg/ha | 12 | 9.08 | 1.49 | | | |
| | 160 kg/ha | 12 | 9.01 | 1.55 | | | |
| Gluten Content (%) | 0 kg/ha | 12 | 24.28 | 4.71 | 10.307 | 5.66 | .000 |
| | 80 kg/ha | 12 | 28.79 | 3.89 | | | |
| | 80+40 kg/ha | 12 | 31.43 | 4.19 | | | |
| | 120 kg/ha | 12 | 31.89 | 3.64 | | | |
| | 120+40 kg/ha | 12 | 32.84 | 3.28 | | | |
| | 160 kg/ha | 12 | 34.02 | 2.83 | | | |
| Zeleny Number | 0 kg/ha | 12 | 25.83 | 7.65 | 13.155 | 5.66 | .000 |
| | 80 kg/ha | 12 | 35.14 | 7.03 | | | |
| | 80+40 kg/ha | 12 | 40.27 | 6.09 | | | |
| | 120 kg/ha | 12 | 41.58 | 7.79 | | | |
| | 120+40 kg/ha | 12 | 43.66 | 4.83 | | | |
| | 160 kg/ha | 12 | 45.53 | 7.49 | | | |

39. Table Impact of undivided/split N topdressing applications on wheat grain, based on Hatvan-Nagyombos site by Treatments, 2015-2016 and 2016-2017 crop seasons, Spss-One Way Anova

12.6.3. Examination of Parameters According to Categorical Variables in Different Sites-Gödöllő

12.6.3.1. Comparison of the Means of Parameters Measured in Gödöllő Based on Genotypes

| | | N | M | SD | t | df | p |
|----------------------------|--------|----|--------|-------|--------|--------|------|
| Test Weight (hl) | Alföld | 18 | 72.22 | 1.90 | 2.259 | 30.046 | .031 |
| | Karéj | 18 | 70.43 | 2.78 | | | |
| Thousand Kernel Weight (g) | Alföld | 18 | 39.87 | 1.09 | -4.213 | 25.211 | .000 |
| | Karéj | 18 | 42.26 | 2.15 | | | |
| Falling Number | Alföld | 18 | 424.37 | 42.50 | 4.080 | 34 | .000 |
| | Karéj | 18 | 368.22 | 40.03 | | | |
| Protein Content (%) | Alföld | 18 | 14.21 | 0.86 | -3.285 | 25.677 | .003 |
| | Karéj | 18 | 15.65 | 1.64 | | | |
| Moisture Content (%) | Alföld | 18 | 12.21 | 2.78 | -2.549 | 34 | .015 |
| | Karéj | 18 | 14.05 | 1.27 | | | |
| Gluten Content (%) | Alföld | 18 | 9.47 | 0.21 | -9.210 | 18.451 | .000 |
| | Karéj | 18 | 11.67 | 0.99 | | | |
| Zeleny Number | Alföld | 18 | 35.53 | 2.37 | 1.589 | 24.662 | .125 |
| | Karéj | 18 | 33.51 | 4.85 | | | |

40. Table Impact of undivided/split N topdressing applications on wheat grain based on Gödöllő site by Genotype, 2018-2019 crop seasons, Spss-One Way Anova

12.6.3.2. Comparison of the Means of Parameters Measured in Gödöllő Based on Treatment

| | | N | M | SD | F | df | p |
|----------------------------|--------------|---|--------|-------|-------|------|------|
| Test Weight (hl) | 0 kg/ha | 6 | 70.03 | 1.98 | 1.220 | 5.30 | .324 |
| | 80 kg/ha | 6 | 71.99 | 2.62 | | | |
| | 80+40 kg/ha | 6 | 72.33 | 2.53 | | | |
| | 120 kg/ha | 6 | 72.48 | 2.59 | | | |
| | 120+40 kg/ha | 6 | 71.10 | 2.81 | | | |
| | 160 kg/ha | 6 | 70.02 | 2.26 | | | |
| Thousand Kernel Weight (g) | 0 kg/ha | 6 | 39.87 | 1.84 | .946 | 5.30 | .466 |
| | 80 kg/ha | 6 | 40.83 | 2.35 | | | |
| | 80+40 kg/ha | 6 | 41.26 | 2.49 | | | |
| | 120 kg/ha | 6 | 42.39 | 2.12 | | | |
| | 120+40 kg/ha | 6 | 41.29 | 1.82 | | | |
| | 160 kg/ha | 6 | 40.77 | 1.75 | | | |
| Falling Number | 0 kg/ha | 6 | 369.67 | 44.91 | 1.252 | 5.30 | .310 |
| | 80 kg/ha | 6 | 373.67 | 21.93 | | | |
| | 80+40 kg/ha | 6 | 414.44 | 57.19 | | | |

| | | | | | | | |
|----------------------|--------------|---|--------|-------|-------|------|------|
| | 120 kg/ha | 6 | 416.39 | 51.77 | | | |
| | 120+40 kg/ha | 6 | 417.11 | 70.28 | | | |
| | 160 kg/ha | 6 | 386.50 | 29.91 | | | |
| Protein Content (%) | 0 kg/ha | 6 | 13.71 | 0.92 | 2.370 | 5.30 | .063 |
| | 80 kg/ha | 6 | 14.20 | 1.34 | | | |
| | 80+40 kg/ha | 6 | 15.18 | 1.17 | | | |
| | 120 kg/ha | 6 | 14.88 | 1.93 | | | |
| | 120+40 kg/ha | 6 | 15.77 | 1.30 | | | |
| | 160 kg/ha | 6 | 15.85 | 1.29 | | | |
| Moisture Content (%) | 0 kg/ha | 6 | 12.19 | 0.84 | 1.165 | 5.30 | .349 |
| | 80 kg/ha | 6 | 12.72 | 1.02 | | | |
| | 80+40 kg/ha | 6 | 13.58 | 0.90 | | | |
| | 120 kg/ha | 6 | 11.89 | 5.17 | | | |
| | 120+40 kg/ha | 6 | 14.32 | 1.26 | | | |
| | 160 kg/ha | 6 | 14.09 | 0.92 | | | |
| Gluten Content (%) | 0 kg/ha | 6 | 10.20 | 1.24 | .370 | 5.30 | .865 |
| | 80 kg/ha | 6 | 10.25 | 1.30 | | | |
| | 80+40 kg/ha | 6 | 10.48 | 1.12 | | | |
| | 120 kg/ha | 6 | 10.53 | 1.42 | | | |
| | 120+40 kg/ha | 6 | 10.95 | 1.57 | | | |
| | 160 kg/ha | 6 | 11.02 | 1.59 | | | |
| Zeleny Number | 0 kg/ha | 6 | 31.25 | 5.57 | 3.884 | 5.30 | .008 |
| | 80 kg/ha | 6 | 32.27 | 3.22 | | | |
| | 80+40 kg/ha | 6 | 34.34 | 1.90 | | | |
| | 120 kg/ha | 6 | 34.12 | 3.29 | | | |
| | 120+40 kg/ha | 6 | 37.95 | 2.11 | | | |
| | 160 kg/ha | 6 | 37.18 | 2.07 | | | |

41. Table Impact of undivided/split N topdressing applications on wheat grain based on Gödöllő site by Treatment, 2018-2019 crop seasons, Spss-One Way Anova

12.6.4. Examination of Parameters According to Categorical Variables in Different Genotypes

12.6.4.1. Comparison of the Means of Parameters Measured for Alföld Genotype Based on Year

| | | N | M | SD | F | df | p |
|----------------------------|------|----|-------|------|---------|------|------|
| Test Weight (hl) | 2016 | 18 | 78.47 | 0.44 | 166.885 | 2.51 | .000 |
| | 2017 | 18 | 79.19 | 0.97 | | | |
| | 2019 | 18 | 72.22 | 1.90 | | | |
| Thousand Kernel Weight (g) | 2016 | 18 | 41.20 | 0.81 | 46.646 | 2.51 | .000 |
| | 2017 | 18 | 36.30 | 2.36 | | | |
| | 2019 | 18 | 39.87 | 1.09 | | | |

| | | | | | | | |
|----------------------|------|----|--------|-------|---------|------|------|
| Falling Number | 2016 | 18 | 418.46 | 18.03 | 82.101 | 2.51 | .000 |
| | 2017 | 18 | 623.56 | 82.67 | | | |
| | 2019 | 18 | 424.37 | 42.50 | | | |
| Protein Content (%) | 2016 | 18 | 15.14 | 1.02 | 3.065 | 2.51 | .055 |
| | 2017 | 18 | 14.56 | 1.43 | | | |
| | 2019 | 18 | 14.21 | 0.86 | | | |
| Moisture Content (%) | 2016 | 18 | 6.49 | 0.20 | 56.663 | 2.51 | .000 |
| | 2017 | 18 | 9.73 | 0.27 | | | |
| | 2019 | 18 | 12.21 | 2.78 | | | |
| Gluten Content (%) | 2016 | 18 | 34.24 | 3.15 | 395.263 | 2.51 | .000 |
| | 2017 | 18 | 32.58 | 4.03 | | | |
| | 2019 | 18 | 9.47 | 0.21 | | | |
| Zeleny Number | 2016 | 18 | 33.24 | 6.54 | 30.463 | 2.51 | .000 |
| | 2017 | 18 | 46.93 | 6.85 | | | |
| | 2019 | 18 | 35.53 | 2.37 | | | |

42. Table Impact of undivided/split N topdressing applications on wheat grain based on Alföld genotype by years, 2015-2016, 2016-2017 and 2018-2019 crop seasons, Spss-One Way Anova

12.6.4.2. Comparison of the Means of Parameters Measured for Alföld Genotype Based on Site

| | | N | M | SD | t | df | p |
|----------------------------|---------|----|--------|--------|--------|--------|------|
| Test Weight (hl) | Hatvan | 36 | 78.83 | 0.83 | 14.089 | 20.283 | .000 |
| | Gödöllő | 18 | 72.22 | 1.90 | | | |
| Thousand Kernel Weight (g) | Hatvan | 36 | 38.75 | 3.03 | -1.976 | 48.718 | .054 |
| | Gödöllő | 18 | 39.87 | 1.09 | | | |
| Falling Number | Hatvan | 36 | 521.01 | 119.56 | 4.333 | 48.541 | .000 |
| | Gödöllő | 18 | 424.37 | 42.50 | | | |
| Protein Content (%) | Hatvan | 36 | 14.85 | 1.26 | 1.917 | 52 | .061 |
| | Gödöllő | 18 | 14.21 | 0.86 | | | |
| Moisture Content (%) | Hatvan | 36 | 8.11 | 1.66 | -6.786 | 52 | .000 |
| | Gödöllő | 18 | 12.21 | 2.78 | | | |
| Gluten Content (%) | Hatvan | 36 | 33.41 | 3.66 | 39.102 | 35.438 | .000 |
| | Gödöllő | 18 | 9.47 | 0.21 | | | |
| Zeleny Number | Hatvan | 36 | 40.08 | 9.58 | 2.693 | 42.747 | .010 |
| | Gödöllő | 18 | 35.53 | 2.37 | | | |

43. Table Impact of undivided/split N topdressing applications on wheat grain based on Alföld genotype by Site, 2015-2016, 2016-2017 and 2018-2019 crop seasons, Spss-One Way Anova

12.6.4.3. Comparison of the Means of Parameters Measured for Alföld Genotype Based on Treatment

| | | N | M | SD | F | df | p |
|----------------------------|--------------|----------|----------|-----------|----------|-----------|----------|
| Test Weight (hl) | 0 kg/ha | 9 | 76.34 | 4.35 | .171 | 5.48 | .972 |
| | 80 kg/ha | 9 | 77.08 | 3.36 | | | |
| | 80+40 kg/ha | 9 | 76.94 | 3.05 | | | |
| | 120 kg/ha | 9 | 76.81 | 2.81 | | | |
| | 120+40 kg/ha | 9 | 76.83 | 3.42 | | | |
| | 160 kg/ha | 9 | 75.77 | 3.98 | | | |
| Thousand Kernel Weight (g) | 0 kg/ha | 9 | 39.40 | 1.89 | .082 | 5.48 | .995 |
| | 80 kg/ha | 9 | 38.96 | 2.28 | | | |
| | 80+40 kg/ha | 9 | 39.11 | 2.82 | | | |
| | 120 kg/ha | 9 | 39.21 | 2.86 | | | |
| | 120+40 kg/ha | 9 | 39.34 | 3.27 | | | |
| | 160 kg/ha | 9 | 38.70 | 2.95 | | | |
| Falling Number | 0 kg/ha | 9 | 431.35 | 72.43 | .765 | 5.48 | .580 |
| | 80 kg/ha | 9 | 468.65 | 103.24 | | | |
| | 80+40 kg/ha | 9 | 503.40 | 103.38 | | | |
| | 120 kg/ha | 9 | 514.01 | 115.96 | | | |
| | 120+40 kg/ha | 9 | 506.40 | 111.29 | | | |
| | 160 kg/ha | 9 | 508.98 | 148.43 | | | |
| Protein Content (%) | 0 kg/ha | 9 | 13.08 | 1.15 | 10.543 | 5.48 | .000 |
| | 80 kg/ha | 9 | 13.96 | 0.75 | | | |
| | 80+40 kg/ha | 9 | 14.96 | 0.87 | | | |
| | 120 kg/ha | 9 | 15.09 | 0.71 | | | |
| | 120+40 kg/ha | 9 | 15.20 | 0.86 | | | |
| | 160 kg/ha | 9 | 15.50 | 0.69 | | | |
| Moisture Content (%) | 0 kg/ha | 9 | 9.44 | 2.48 | .268 | 5.48 | .928 |
| | 80 kg/ha | 9 | 9.48 | 2.48 | | | |
| | 80+40 kg/ha | 9 | 9.62 | 2.98 | | | |
| | 120 kg/ha | 9 | 8.52 | 3.69 | | | |
| | 120+40 kg/ha | 9 | 9.93 | 2.92 | | | |
| | 160 kg/ha | 9 | 9.86 | 3.00 | | | |
| Gluten Content (%) | 0 kg/ha | 9 | 21.36 | 9.39 | .308 | 5.48 | .906 |
| | 80 kg/ha | 9 | 24.32 | 11.37 | | | |
| | 80+40 kg/ha | 9 | 26.19 | 12.58 | | | |
| | 120 kg/ha | 9 | 26.60 | 12.81 | | | |
| | 120+40 kg/ha | 9 | 26.70 | 13.04 | | | |
| | 160 kg/ha | 9 | 27.42 | 13.41 | | | |
| Zeleny Number | 0 kg/ha | 9 | 30.88 | 7.47 | 3.195 | 5.48 | .014 |
| | 80 kg/ha | 9 | 35.90 | 6.72 | | | |
| | 80+40 kg/ha | 9 | 39.58 | 6.69 | | | |
| | 120 kg/ha | 9 | 40.57 | 9.09 | | | |
| | 120+40 kg/ha | 9 | 41.70 | 5.73 | | | |
| | 160 kg/ha | 9 | 42.77 | 8.53 | | | |

44. Table Impact of undivided/split N topdressing applications on wheat grain based on Alföld genotype by Treatment, 2015-2016, 2016-2017 and 2018-2019 crop seasons, Spss-One Way Anova

12.6.4.4. Comparison of the Means of Parameters Measured for Karéj Genotype Based on Year

| | | N | M | SD | F | df | p |
|----------------------------|------|----------|----------|-----------|----------|-----------|----------|
| Test Weight (hl) | 2016 | 18 | 77.67 | 0.61 | 81.669 | 2.51 | .000 |
| | 2017 | 18 | 73.65 | 0.76 | | | |
| | 2019 | 18 | 70.43 | 2.78 | | | |
| Thousand Kernel Weight (g) | 2016 | 18 | 45.16 | 1.49 | 21.077 | 2.51 | .000 |
| | 2017 | 18 | 46.09 | 1.85 | | | |
| | 2019 | 18 | 42.26 | 2.15 | | | |
| Falling Number | 2016 | 18 | 312.49 | 17.15 | 53.958 | 2.51 | .000 |
| | 2017 | 18 | 433.09 | 41.82 | | | |
| | 2019 | 18 | 368.22 | 40.03 | | | |
| Protein Content (%) | 2016 | 18 | 13.93 | 1.06 | 23.524 | 2.51 | .000 |
| | 2017 | 18 | 12.39 | 1.51 | | | |
| | 2019 | 18 | 15.65 | 1.64 | | | |
| Moisture Content (%) | 2016 | 18 | 9.75 | 0.32 | 171.195 | 2.51 | .000 |
| | 2017 | 18 | 10.22 | 0.22 | | | |
| | 2019 | 18 | 14.05 | 1.27 | | | |
| Gluten Content (%) | 2016 | 18 | 28.71 | 3.29 | 128.146 | 2.51 | .000 |
| | 2017 | 18 | 26.63 | 4.96 | | | |
| | 2019 | 18 | 11.67 | 0.99 | | | |
| Zeleny Number | 2016 | 18 | 34.99 | 7.40 | 2.879 | 2.51 | .065 |
| | 2017 | 18 | 39.51 | 10.25 | | | |
| | 2019 | 18 | 33.51 | 4.85 | | | |

45. Table Impact of undivided/split N topdressing applications on wheat grain based on Karéj genotype by Year, 2015-2016, 2016-2017 and 2018-2019 crop seasons, Spss-One Way Anova

12.6.4.5. Comparison of the Means of Parameters Measured for Karéj Genotype Based on Site

| | | N | M | SD | t | df | p |
|----------------------------|---------|----------|----------|-----------|----------|-----------|----------|
| Test Weight (hl) | Hatvan | 36 | 75.66 | 2.15 | 7.000 | 27.452 | .000 |
| | Gödöllő | 18 | 70.43 | 2.78 | | | |
| Thousand Kernel Weight (g) | Hatvan | 36 | 45.63 | 1.72 | 5.788 | 28.231 | .000 |
| | Gödöllő | 18 | 42.26 | 2.15 | | | |
| Falling Number | Hatvan | 36 | 372.79 | 68.79 | .307 | 50.641 | .760 |
| | Gödöllő | 18 | 368.22 | 40.03 | | | |

| | | | | | | | |
|----------------------|---------|----|-------|------|---------|--------|------|
| Protein Content (%) | Hatvan | 36 | 13.16 | 1.50 | -5.561 | 52 | .000 |
| | Gödöllő | 18 | 15.65 | 1.64 | | | |
| Moisture Content (%) | Hatvan | 36 | 9.98 | 0.36 | -13.349 | 18.380 | .000 |
| | Gödöllő | 18 | 14.05 | 1.27 | | | |
| Gluten Content (%) | Hatvan | 36 | 27.67 | 4.28 | 21.316 | 41.942 | .000 |
| | Gödöllő | 18 | 11.67 | 0.99 | | | |
| Zeleny Number | Hatvan | 36 | 37.25 | 9.10 | 1.971 | 51.711 | .054 |
| | Gödöllő | 18 | 33.51 | 4.85 | | | |

46. Table Impact of undivided/split N topdressing applications on wheat grain based on Karéj genotype by Site, 2015-2016, 2016-2017 and 2018-2019 crop seasons, Spss-One Way Anova

12.6.4.6. Comparison of the Means of Parameters Measured for Karéj Genotype Based on Treatment

| | | N | M | SD | F | df | p |
|----------------------------|--------------|---|--------|-------|-------|------|------|
| Test Weight (hl) | 0 kg/ha | 9 | 73.48 | 3.66 | .112 | 5.48 | .989 |
| | 80 kg/ha | 9 | 74.15 | 3.35 | | | |
| | 80+40 kg/ha | 9 | 74.45 | 3.35 | | | |
| | 120 kg/ha | 9 | 74.13 | 3.13 | | | |
| | 120+40 kg/ha | 9 | 73.81 | 3.97 | | | |
| | 160 kg/ha | 9 | 73.45 | 3.92 | | | |
| Thousand Kernel Weight (g) | 0 kg/ha | 9 | 44.80 | 3.52 | .280 | 5.48 | .922 |
| | 80 kg/ha | 9 | 44.46 | 2.33 | | | |
| | 80+40 kg/ha | 9 | 44.70 | 2.42 | | | |
| | 120 kg/ha | 9 | 45.02 | 1.32 | | | |
| | 120+40 kg/ha | 9 | 44.31 | 2.39 | | | |
| | 160 kg/ha | 9 | 43.75 | 2.74 | | | |
| Falling Number | 0 kg/ha | 9 | 334.27 | 35.14 | 1.103 | 5.48 | .371 |
| | 80 kg/ha | 9 | 363.57 | 48.24 | | | |
| | 80+40 kg/ha | 9 | 368.89 | 51.92 | | | |
| | 120 kg/ha | 9 | 382.23 | 62.81 | | | |
| | 120+40 kg/ha | 9 | 387.68 | 77.96 | | | |
| | 160 kg/ha | 9 | 390.95 | 73.29 | | | |
| Protein Content (%) | 0 kg/ha | 9 | 12.09 | 1.91 | 4.060 | 5.48 | .004 |
| | 80 kg/ha | 9 | 13.35 | 1.66 | | | |
| | 80+40 kg/ha | 9 | 14.17 | 1.80 | | | |
| | 120 kg/ha | 9 | 14.10 | 1.84 | | | |
| | 120+40 kg/ha | 9 | 15.01 | 1.58 | | | |
| | 160 kg/ha | 9 | 15.21 | 1.43 | | | |
| Moisture Content (%) | 0 kg/ha | 9 | 10.84 | 1.25 | .206 | 5.48 | .958 |
| | 80 kg/ha | 9 | 11.08 | 1.76 | | | |
| | 80+40 kg/ha | 9 | 11.33 | 2.13 | | | |
| | 120 kg/ha | 9 | 11.51 | 2.33 | | | |
| | 120+40 kg/ha | 9 | 11.72 | 2.72 | | | |
| | 160 kg/ha | 9 | 11.55 | 2.47 | | | |

| | | | | | | | |
|--------------------|--------------|---|-------|------|--------|------|------|
| Gluten Content (%) | 0 kg/ha | 9 | 17.82 | 5.96 | .909 | 5.48 | .483 |
| | 80 kg/ha | 9 | 20.90 | 7.60 | | | |
| | 80+40 kg/ha | 9 | 22.71 | 8.91 | | | |
| | 120 kg/ha | 9 | 22.93 | 8.70 | | | |
| | 120+40 kg/ha | 9 | 24.39 | 9.10 | | | |
| | 160 kg/ha | 9 | 25.28 | 9.73 | | | |
| Zeleny Number | 0 kg/ha | 9 | 24.39 | 5.92 | 14.195 | 5.48 | .000 |
| | 80 kg/ha | 9 | 32.47 | 5.19 | | | |
| | 80+40 kg/ha | 9 | 37.00 | 4.73 | | | |
| | 120 kg/ha | 9 | 37.62 | 5.52 | | | |
| | 120+40 kg/ha | 9 | 41.82 | 4.29 | | | |
| | 160 kg/ha | 9 | 42.72 | 6.47 | | | |

47. Table Impact of undivided/split N topdressing applications on wheat grain based on Karéj genotype by Treatment, 2015-2016, 2016-2017 and 2018-2019 crop seasons, Spss-One Way Anova

| | | ANOVA by treatments | | | | | |
|--------|---------------------------|----------------------------|----------------|-----|-------------|--------|------|
| | | | Sum of Squares | df | Mean Square | F | Sig. |
| Alföld | Test weight(hl) | Between Groups | 11,987 | 5 | 2,397 | 2,193 | ,061 |
| | | Within Groups | 111,484 | 102 | 1,093 | | |
| | | Total | 123,471 | 107 | | | |
| | Thousand kernel weight(g) | Between Groups | 27,954 | 5 | 5,591 | ,513 | ,766 |
| | | Within Groups | 1110,874 | 102 | 10,891 | | |
| | | Total | 1138,829 | 107 | | | |
| | Falling number | Between Groups | 160764,750 | 5 | 32152,950 | 2,220 | ,058 |
| | | Within Groups | 1477427,352 | 102 | 14484,582 | | |
| | | Total | 1638192,102 | 107 | | | |
| | Protein | Between Groups | 111,323 | 5 | 22,265 | 18,682 | ,000 |
| | | Within Groups | 121,559 | 102 | 1,192 | | |
| | | Total | 232,882 | 107 | | | |
| | Gluten | Between Groups | 987,878 | 5 | 197,576 | 21,084 | ,000 |
| | | Within Groups | 955,809 | 102 | 9,371 | | |
| | | Total | 1943,687 | 107 | | | |
| | Zeleny | Between Groups | 3659,712 | 5 | 731,942 | 9,528 | ,000 |
| | | Within Groups | 7835,598 | 102 | 76,820 | | |
| | | Total | 11495,310 | 107 | | | |

| | | | | | | | | |
|----------|---------------------------|-----------------|----------------|-------|-----------|--------|------|------|
| MV-Nádor | Test weight(hl) | Between Groups | 9,561 | 5 | 1,912 | 1,503 | ,195 | |
| | | Within Groups | 129,750 | 102 | 1,272 | | | |
| | | Total | 139,312 | 107 | | | | |
| | Thousand kernel weight(g) | Between Groups | 138,837 | 5 | 27,767 | 6,244 | ,000 | |
| | | Within Groups | 453,604 | 102 | 4,447 | | | |
| | | Total | 592,440 | 107 | | | | |
| | Falling number | Between Groups | 5606,264 | 5 | 1121,253 | 0,095 | ,993 | |
| | | Within Groups | 1206753,401 | 102 | 11830,916 | | | |
| | | Total | 1212359,666 | 107 | | | | |
| | Protein | Between Groups | 65,112 | 5 | 13,022 | 16,210 | ,000 | |
| | | Within Groups | 81,941 | 102 | 0,803 | | | |
| | | Total | 147,052 | 107 | | | | |
| | Gluten | Between Groups | 592,986 | 5 | 118,597 | 15,686 | ,000 | |
| | | Within Groups | 771,184 | 102 | 7,561 | | | |
| | | Total | 1364,170 | 107 | | | | |
| | Zeleny | Between Groups | 2374,650 | 5 | 474,930 | 9,763 | ,000 | |
| | | Within Groups | 4962,066 | 102 | 48,648 | | | |
| | | Total | 7336,717 | 107 | | | | |
| | MV-Karéj | Test weight(hl) | Between Groups | 3,233 | 5 | ,647 | ,126 | ,986 |

| | | | | | | | |
|------------|---------------------------|----------------|------------|-----|----------|--------|------|
| | | Within Groups | 521,596 | 102 | 5,114 | | |
| | | Total | 524,829 | 107 | | | |
| | Thousand kernel weight(g) | Between Groups | 81,695 | 5 | 16,339 | 1,337 | ,255 |
| | | Within Groups | 1246,449 | 102 | 12,220 | | |
| | | Total | 1328,144 | 107 | | | |
| | Falling number | Between Groups | 44637,317 | 5 | 8927,463 | 1,811 | ,117 |
| | | Within Groups | 502698,593 | 102 | 4928,418 | | |
| | | Total | 547335,909 | 107 | | | |
| | Protein | Between Groups | 121,323 | 5 | 24,265 | 13,861 | ,000 |
| | | Within Groups | 178,559 | 102 | 1,751 | | |
| | | Total | 299,883 | 107 | | | |
| | Gluten | Between Groups | 1279,148 | 5 | 255,830 | 19,902 | ,000 |
| | | Within Groups | 1311,162 | 102 | 12,855 | | |
| | | Total | 2590,309 | 107 | | | |
| | Zeleny | Between Groups | 6202,355 | 5 | 1240,471 | 22,893 | ,000 |
| | | Within Groups | 5526,898 | 102 | 54,185 | | |
| | | Total | 11729,253 | 107 | | | |
| MV-Toborzó | Test weight(hl) | Between Groups | 2,106 | 5 | ,421 | ,081 | ,995 |
| | | Within Groups | 529,535 | 102 | 5,192 | | |
| | | Total | 531,641 | 107 | | | |

| | | | | | | | | |
|--|---------------------------|------------------|----------------|---------|-----------|--------|-------|------|
| | Thousand kernel weight(g) | Between Groups | 131,000 | 5 | 26,200 | 1,048 | ,394 | |
| | | Within Groups | 2551,082 | 102 | 25,011 | | | |
| | | Total | 2682,082 | 107 | | | | |
| | Falling number | Between Groups | 33203,598 | 5 | 6640,720 | ,493 | ,781 | |
| | | Within Groups | 1374205,673 | 102 | 13472,605 | | | |
| | | Total | 1407409,271 | 107 | | | | |
| | Protein | Between Groups | 75,826 | 5 | 15,165 | 6,574 | ,000 | |
| | | Within Groups | 235,308 | 102 | 2,307 | | | |
| | | Total | 311,134 | 107 | | | | |
| | Gluten | Between Groups | 797,881 | 5 | 159,576 | 9,944 | ,000 | |
| | | Within Groups | 1636,827 | 102 | 16,047 | | | |
| | | Total | 2434,708 | 107 | | | | |
| | Zeleny | Between Groups | 3676,516 | 5 | 735,303 | 22,665 | ,000 | |
| | | Within Groups | 3309,076 | 102 | 32,442 | | | |
| | | Total | 6985,592 | 107 | | | | |
| | MV-Toldi | Test weight (hl) | Between Groups | 20,703 | 5 | 4,141 | ,702 | ,623 |
| | | | Within Groups | 601,572 | 102 | 5,898 | | |
| | | | Total | 622,275 | 107 | | | |
| | | | Between Groups | 93,379 | 5 | 18,676 | 3,692 | ,004 |

| | | | | | | | |
|--|---------------------------|----------------|-------------|-----|-----------|--------|------|
| | Thousand kernel weight(g) | Within Groups | 515,915 | 102 | 5,058 | | |
| | | Total | 609,294 | 107 | | | |
| | Falling number | Between Groups | 26999,588 | 5 | 5399,918 | ,152 | ,979 |
| | | Within Groups | 3624628,161 | 102 | 35535,570 | | |
| | | Total | 3651627,749 | 107 | | | |
| | Protein | Between Groups | 53,587 | 5 | 10,717 | 8,847 | ,000 |
| | | Within Groups | 123,571 | 102 | 1,211 | | |
| | | Total | 177,159 | 107 | | | |
| | Gluten | Between Groups | 545,280 | 5 | 109,056 | 7,548 | ,000 |
| | | Within Groups | 1473,787 | 102 | 14,449 | | |
| | | Total | 2019,068 | 107 | | | |
| | Zeleny | Between Groups | 3088,550 | 5 | 617,710 | 10,289 | ,000 |
| | | Within Groups | 6123,849 | 102 | 60,038 | | |
| | | Total | 9212,399 | 107 | | | |

48. Table Impact of undivided/split N topdressing applications on wheat grain, 2015-2016 and 2016-2017 crop seasons, Spss-One Way Anova

| ANOVA ^a | | | | | | | |
|--------------------|----------------|----------------|----|-------------|------|------|--|
| | | Sum of Squares | df | Mean Square | F | Sig. | |
| Test weight (hl) | Between Groups | 3,010 | 1 | 3,010 | ,729 | ,441 | |
| | Within Groups | 16,523 | 4 | 4,131 | | | |

| | | | | | | |
|----------------------------|----------------|-----------|---|----------|--------|------|
| | Total | 19,534 | 5 | | | |
| Thousand kernel weight (g) | Between Groups | 7,504 | 1 | 7,504 | 3,195 | ,148 |
| | Within Groups | 9,394 | 4 | 2,349 | | |
| | Total | 16,898 | 5 | | | |
| Falling number | Between Groups | 9178,074 | 1 | 9178,074 | 40,445 | ,003 |
| | Within Groups | 907,704 | 4 | 226,926 | | |
| | Total | 10085,778 | 5 | | | |
| Protein content (%) | Between Groups | ,256 | 1 | ,256 | ,260 | ,637 |
| | Within Groups | 3,936 | 4 | ,984 | | |
| | Total | 4,192 | 5 | | | |
| Moisture content (%) | Between Groups | ,205 | 1 | ,205 | ,250 | ,643 |
| | Within Groups | 3,286 | 4 | ,821 | | |
| | Total | 3,491 | 5 | | | |
| Gluten content (%) | Between Groups | 5,227 | 1 | 5,227 | 8,522 | ,043 |
| | Within Groups | 2,453 | 4 | ,613 | | |
| | Total | 7,680 | 5 | | | |
| Zeleny number | Between Groups | 103,335 | 1 | 103,335 | 8,017 | ,047 |
| | Within Groups | 51,560 | 4 | 12,890 | | |
| | Total | 154,895 | 5 | | | |

49. Table Impact of undivided/split N topdressing applications on wheat grain, 2018-2019 crop seasons 0kg/ha, Spss-One Way Anova

| | | ANOVA ^a | | | | |
|-------|--|--------------------|----|-------------|---|------|
| Treat | | Sum of Squares | df | Mean Square | F | Sig. |

| | | | | | | | |
|--|----------------------------|----------------|----------|---|---------|-------|------|
| | Test weight (hl) | Between Groups | 4,770 | 1 | 4,770 | ,644 | ,467 |
| | | Within Groups | 29,627 | 4 | 7,407 | | |
| | | Total | 34,397 | 5 | | | |
| | Thousand kernel weight (g) | Between Groups | 11,676 | 1 | 11,676 | 2,926 | ,162 |
| | | Within Groups | 15,962 | 4 | 3,990 | | |
| | | Total | 27,638 | 5 | | | |
| | Falling number | Between Groups | 711,407 | 1 | 711,407 | 1,680 | ,265 |
| | | Within Groups | 1693,481 | 4 | 423,370 | | |
| | | Total | 2404,889 | 5 | | | |
| | Protein content (%) | Between Groups | 3,496 | 1 | 3,496 | 2,580 | ,183 |
| | | Within Groups | 5,419 | 4 | 1,355 | | |
| | | Total | 8,915 | 5 | | | |
| | Moisture content (%) | Between Groups | 1,771 | 1 | 1,771 | 2,085 | ,222 |
| | | Within Groups | 3,399 | 4 | ,850 | | |
| | | Total | 5,170 | 5 | | | |
| | Gluten content (%) | Between Groups | 5,434 | 1 | 5,434 | 7,328 | ,054 |
| | | Within Groups | 2,966 | 4 | ,742 | | |
| | | Total | 8,400 | 5 | | | |
| | Zeleny number | Between Groups | 13,500 | 1 | 13,500 | 1,406 | ,301 |
| | | Within Groups | 38,393 | 4 | 9,598 | | |
| | | Total | 51,893 | 5 | | | |

50. Table Impact of undivided/split N topdressing applications on wheat grain, 2018-2019 crop seasons 80kg/ha, Spss-One Way Anova

| | |
|--|--------------------------|
| | ANOVA^a |
|--|--------------------------|

| | | Sum of Squares | df | Mean Square | F | Sig. | |
|-------------------------|----------------------------|----------------|-----------|-------------|-----------|-------|------|
| Treatment = 80+40 kg/ha | Test weight(hl) | Between Groups | 3,682 | 1 | 3,682 | ,521 | ,510 |
| | | Within Groups | 28,257 | 4 | 7,064 | | |
| | | Total | 31,938 | 5 | | | |
| | Thousand kernel weight (g) | Between Groups | 8,520 | 1 | 8,520 | 1,523 | ,285 |
| | | Within Groups | 22,372 | 4 | 5,593 | | |
| | | Total | 30,892 | 5 | | | |
| | Falling number | Between Groups | 11208,963 | 1 | 11208,963 | 8,716 | ,042 |
| | | Within Groups | 5144,296 | 4 | 1286,074 | | |
| | | Total | 16353,259 | 5 | | | |
| | Protein content (%) | Between Groups | 2,707 | 1 | 2,707 | 2,647 | ,179 |
| | | Within Groups | 4,091 | 4 | 1,023 | | |
| | | Total | 6,798 | 5 | | | |
| | Moisture content (%) | Between Groups | 1,335 | 1 | 1,335 | 1,965 | ,234 |
| | | Within Groups | 2,717 | 4 | ,679 | | |
| | | Total | 4,052 | 5 | | | |
| | Gluten content (%) | Between Groups | 4,335 | 1 | 4,335 | 8,969 | ,040 |
| | | Within Groups | 1,933 | 4 | ,483 | | |
| | | Total | 6,268 | 5 | | | |
| | Zeleny number | Between Groups | 5,549 | 1 | 5,549 | 1,771 | ,254 |
| | | Within Groups | 12,535 | 4 | 3,134 | | |
| | | Total | 18,084 | 5 | | | |

51. Table Impact of undivided/split N topdressing applications on wheat grain, 2018-2019 crop seasons 80+40 kg/ha, Spss-One Way Anova

| | | ANOVA ^a | | | | | |
|-----------------------|----------------------------|--------------------|----------------|----|-------------|--------|------|
| | | | Sum of Squares | df | Mean Square | F | Sig. |
| Treatment = 120 kg/ha | Test weight (hl) | Between Groups | 4,507 | 1 | 4,507 | ,621 | ,475 |
| | | Within Groups | 29,027 | 4 | 7,257 | | |
| | | Total | 33,533 | 5 | | | |
| | Thousand kernel weight (g) | Between Groups | 20,572 | 1 | 20,572 | 44,062 | ,003 |
| | | Within Groups | 1,868 | 4 | ,467 | | |
| | | Total | 22,440 | 5 | | | |
| | Falling number | Between Groups | 7049,796 | 1 | 7049,796 | 4,440 | ,103 |
| | | Within Groups | 6351,407 | 4 | 1587,852 | | |
| | | Total | 13401,204 | 5 | | | |
| | Protein content (%) | Between Groups | 1,938 | 1 | 1,938 | ,466 | ,532 |
| | | Within Groups | 16,627 | 4 | 4,157 | | |
| | | Total | 18,565 | 5 | | | |
| | Moisture content (%) | Between Groups | 40,925 | 1 | 40,925 | 1,768 | ,254 |
| | | Within Groups | 92,579 | 4 | 23,145 | | |
| | | Total | 133,504 | 5 | | | |
| | Gluten content (%) | Between Groups | 6,000 | 1 | 6,000 | 5,892 | ,072 |
| | | Within Groups | 4,073 | 4 | 1,018 | | |
| | | Total | 10,073 | 5 | | | |
| | Zeleny number | Between Groups | ,735 | 1 | ,735 | ,055 | ,826 |
| | | Within Groups | 53,433 | 4 | 13,358 | | |
| | | Total | 54,168 | 5 | | | |

52. Table Impact of undivided/split N topdressing applications on wheat grain, 2018-2019 crop seasons 120 kg/ha, Spss-One Way Anova

| | | ANOVA ^a | | | | | |
|--------------------------|----------------------------|--------------------|----------------|----|-------------|---------|-------|
| | | | Sum of Squares | df | Mean Square | F | Sig. |
| Treatment = 120+40 kg/ha | Test weight(hl) | Between Groups | 10,935 | 1 | 10,935 | 1,528 | ,284 |
| | | Within Groups | 28,620 | 4 | 7,155 | | |
| | | Total | 39,555 | 5 | | | |
| | Thousand kernel weight (g) | Between Groups | 5,078 | 1 | 5,078 | 1,772 | ,254 |
| | | Within Groups | 11,461 | 4 | 2,865 | | |
| | | Total | 16,539 | 5 | | | |
| | Falling number | Between Groups | 4231,185 | 1 | 4231,185 | ,827 | ,415 |
| | | Within Groups | 20464,074 | 4 | 5116,019 | | |
| | | Total | 24695,259 | 5 | | | |
| | Protein content (%) | Between Groups | 6,242 | 1 | 6,242 | 11,355 | ,028 |
| | | Within Groups | 2,199 | 4 | ,550 | | |
| | | Total | 8,441 | 5 | | | |
| | Moisture content (%) | Between Groups | 5,684 | 1 | 5,684 | 9,873 | ,035 |
| | | Within Groups | 2,303 | 4 | ,576 | | |
| | | Total | 7,987 | 5 | | | |
| | Gluten content (%) | Between Groups | 12,042 | 1 | 12,042 | 136,321 | <,001 |
| | | Within Groups | ,353 | 4 | ,088 | | |
| | | Total | 12,395 | 5 | | | |
| | Zeleny number | Between Groups | ,202 | 1 | ,202 | ,037 | ,858 |
| | | Within Groups | 22,093 | 4 | 5,523 | | |
| | | Total | 22,295 | 5 | | | |

53. Table Impact of undivided/split N topdressing applications on wheat grain, 2018-2019 crop seasons 120+40 kg/ha, Spss-One Way Anova

| | | ANOVA ^a | | | | | |
|-----------------------|----------------------------|--------------------|----------------|----|-------------|---------|-------|
| | | | Sum of Squares | df | Mean Square | F | Sig. |
| Treatment = 160 kg/ha | Test weight (hl) | Between Groups | 3,682 | 1 | 3,682 | ,671 | ,459 |
| | | Within Groups | 21,947 | 4 | 5,487 | | |
| | | Total | 25,628 | 5 | | | |
| | Thousand kernel weight (g) | Between Groups | 2,926 | 1 | 2,926 | ,950 | ,385 |
| | | Within Groups | 12,323 | 4 | 3,081 | | |
| | | Total | 15,249 | 5 | | | |
| | Falling number | Between Groups | 1242,241 | 1 | 1242,241 | 1,538 | ,283 |
| | | Within Groups | 3230,370 | 4 | 807,593 | | |
| | | Total | 4472,611 | 5 | | | |
| | Protein content (%) | Between Groups | 6,998 | 1 | 6,998 | 20,445 | ,011 |
| | | Within Groups | 1,369 | 4 | ,342 | | |
| | | Total | 8,368 | 5 | | | |
| | Moisture content (%) | Between Groups | 3,154 | 1 | 3,154 | 11,485 | ,028 |
| | | Within Groups | 1,098 | 4 | ,275 | | |
| | | Total | 4,252 | 5 | | | |
| | Gluten content (%) | Between Groups | 12,615 | 1 | 12,615 | 688,091 | <,001 |
| | | Within Groups | ,073 | 4 | ,018 | | |
| | | Total | 12,688 | 5 | | | |

| | | | | | | | |
|--|---------------|-------------------|--------|---|-------|------|------|
| | Zeleny number | Between Groups | 3,082 | 1 | 3,082 | ,672 | ,458 |
| | | Within Groups | 18,347 | 4 | 4,587 | | |
| | | Total | 21,428 | 5 | | | |

54. Table Impact of undivided/split N topdressing applications on wheat grain, 2018-2019 crop seasons 160 kg/ha, Spss-One Way Anova