

Doctoral (PhD) dissertation

Ribal Masri

Gödöllő

2023



Hungarian University of Agriculture and Life Sciences

A transcriptomic study exploring the role of the *Vitis vinifera* NAC genes in response to stresses: case of *Botrytis cinerea*

DOI: 10.54598/004270

Doctoral (PhD) dissertation

Ribal Masri

Gödöllő

2023

The PhD School

Name: Doctoral School of Biological Sciences

Discipline: Plant Biotechnology

Head: **Prof. Dr. Zoltán Nagy**
Director of the institute, professor
MATE, Gödöllő

Supervisor: **Prof. Dr. Erzsébet Kiss**
Professor emeritus
MATE, Gödöllő
Institute of Genetics and Biotechnology

.....
Prof. Dr. Zoltán Nagy
Approval of the School Leader

.....
Prof. Dr. Erzsébet Kiss
Approval of the Supervisor

LIST OF ABBREVIATIONS

ABA	Abscisic acid
ABF	ABA binding factor
ABRE	abscisic acid responsive element
AREB	ABA-responsive element
BAP	6-Benzylaminopurine
Bgh	Blumeria graminis f. sp. Hordei
CARE's	Cis-acting regulatory elements
CDS	Coding DNA sequence
DPE	Downstream promoter elements
DRE	Dehydration-responsive elements
DREB	Dehydration responsive binding element
DREB	DRE Binding
ERE	Ethylene responsive element
ET	Ethylene
HR	Hypersensitive response
ID	Intrinsic disorder
Inr	Initiator region
JA	Jasmonates
LTREs	Low-temperature responsive elements
MiRNA	Micro- RNA
MW	Molecular weight
MYB	Myeloblastosis
MYC	Myelocytomatosis
NAA	Naphthalene acetic acid

NACRS	NAC-recognition sequence for drought response
NACs	NAM – no apical meristem, ATAF – <i>Arabidopsis</i> transcription activation factor, and CUC – cup-shaped cotyledon
NGS	Next-generation sequencing
NPR1	Natriuretic peptide receptor 1
ORF	Open reading frame
Pi	Isoelectric point
PM	Powdery Mildew
PR	Pathogen related
SA	Salicylic acid
SAGE	Serial Analysis of Gene Expression
SAR	Systemic acquired resistance
SGFP	Synthetic green fluorescent protein
SiRNA	Small interfering RNA
STRE	stress responsive elements
TF	Transcription factor
TFDB	transcription factor database
TILLING	Targeted Induced Local Lesions in Genome
TMH	Transmembrane helices
TR	Transcription regulatory
TRR	Transcription regulatory regions
TSS	Transcription start site
Vv	<i>Vitis vinifera</i> L.
WRE3	Wound-responsive elements 3

TABLE OF CONTENT

LIST OF ABBREVIATIONS	4
1. INTRODUCTION	8
1.1 Objectives	9
2. LITERATURE REVIEW	10
2.1. Plant defence mechanisms	10
2.1.1. Biotic Stress defence mechanisms	10
2.1.2. Abiotic stress defence mechanisms	10
2.1.3. Crosstalk between biotic and abiotic stress responses.....	11
2.2. Cis-acting regulatory elements of plants	11
2.2.1. Abiotic stress and CAREs	12
2.2.2. Biotic stress and CAREs	12
2.3. Bioinformatics exploration	12
2.3.1. Areas of bioinformatics	12
2.3.2. Bioinformatic bases	13
2.3.3. Bioinformatic applications.....	13
2.3.4. Bioinformatics and stresses.....	14
2.3.5. Bioinformatics examples	15
2.4. NAC transcription factors	16
2.4.1. Structure of NAC proteins	16
2.4.2. NAC regulation	16
2.4.3. NAC genes expression	17
2.4.4. Abiotic and biotic stresses response	17
2.4.5. NAC TFs in multiple processes	18
2.5. <i>Vitis vinifera</i> L.	19
2.5.1. Description	19
2.5.2. Abiotic and biotic stresses.....	19
2.5.3. Botrytis cinerea	20
2.5.4. Applied breeding methods and objectives in <i>Vitis vinifera</i> L.	20
2.6. <i>In silico</i> methods: Definition, applications, and objectives.....	21
3. MATERIALS AND METHODS	23
3.1. <i>In silico</i> part	23
3.1.1. Detection and data analyses of NAC family genes	23
3.1.2. Conserved motifs and cis-acting regulatory elements (CAREs) of VvNAC genes.....	23
3.1.3. Protein–protein interaction analysis.....	24

3.1.4. In silico expression profiles of VvNAC genes and statistical analysis	24
3.2. In vitro part.....	24
3.2.1. Construction of plant expression vectors.....	24
3.2.2. Transient Expression	26
4. RESULTS.....	28
4.1. Identification and genomic distribution of NAC proteins	28
4.2. TMHs and KaKs value of VvNAC genes	28
4.3. Conserved motifs of VvNAC genes	30
4.4. Protein–protein interaction of VvNAC	35
4.5. Promoter regions detection and analysis of cis-acting regulatory elements	36
4.5.1. Stress responsive cis-acting regulatory elements	36
4.5.2. Cis-acting regulatory elements in hormonal regulation.....	37
4.5.3. Role of cis-acting regulatory elements in cellular development:.....	37
4.5.4. In silico analysis of VvNAC gene expression.....	37
4.5.5. Subcellular localization prediction	38
4.5.6. Statistical Analysis	39
4.5.7. In vitro verifications: VvNAC36 model	41
5. DISCUSSION AND LIMITATIONS	43
5.1. VvNAC36 protein	46
5.2. Study limitations	46
6. CONCLUSIONS AND RECOMMENDATIONS	47
6.1. New scientific results	48
7. APPENDIXES.....	49
A1: Bibliography.....	49
A2: Supplementary data	64
8. ACKNOWLEDGMENTS.....	75

1. INTRODUCTION

Biotic and abiotic stresses are the main factors that significantly limit growth and fruit quality of essential crops worldwide. Plants have developed the ability to cope with the harmful effects of stresses by activating a number of defense mechanisms which include phytohormone signaling networks (Masri and Kiss 2023).

Transcription factor (TFs) proteins are essential regulators of gene expression in all living organisms, with roles in plant growth, cell cycling, cell signaling, and stress response (Li et al., 2021) by binding to specific short sequence motifs mostly detected within the promoters of the target genes, known as cis-regulatory sequences (Priest et al., 2009). Thus, the interaction between them, formed a crucial functional link of gene regulatory networks in plant defence response by controlling various stresses (Kaur et al., 2016).

The NACs (NAM – no apical meristem, ATAF – *Arabidopsis* transcription activation factor, and CUC – cup-shaped cotyledon) are considered as one of the largest TF families that consists of N-terminus and C-terminal. The NAC domain is highly conserved, comprise of 160 amino acid residues that classified into five subdomains from A to E (Puranik et al., 2012). The researchers discovered several NAC genes in various plant species using genomic sequencing, including 151 in rice, 117 in *Arabidopsis*, 74 in grape (Shao et al., 2015; Ju et al., 2020).

Different roles for NAC proteins in response to abiotic and biotic stresses, as well as in developmental processes have been reported such as, leaf senescence (Breeze et al. 2011), cell division (Kim et al. 2006), seed development (Sperotto et al. 2009), fiber development (Ko et al. 2007), shoot apical meristem formation (Kim et al. 2006) and embryo development (Duval et al. 2002).

Grapevine (*Vitis vinifera* L.), a member of the *Vitaceae* family and one of the world's most valuable fruit crops with high economic value is prone to a wide range of pathogens that cause yield and quality losses (Ju et al., 2020).

Plant pathogens are categorized to different classes according to their modes of nutrition, such as: biotrophic, necrotrophic, hemibiotrophic and saprophytic pathogens.

The necrotrophic fungus *Botrytis cinerea*, is the main cause of the grey mold disease in more than 200 plants species (Elad et al., 2004) and one of the most crucial pathogens in grapevine (Haile et al., 2017) with a short biotrophic phase (Veloso and van Kan, 2018).

However, in the vineyards, the infection of *B. cinerea* targets primarily the ripe berries; the pathogen inoculation occurs during the onset of the grape berry development (Keller et al., 2003; Pezet et al., 2003).

The *B. cinerea* weakens the natural pathogen resistance system of grape berries and imposes different modifications such as lowering the mechanical resistance of the cell-wall, compacting the bunches and increasing the sugar levels and the concentration of organic acids (Prusky et al., 2013; Blanco-Ulate et al., 2015). Additionally, speeding up the ripening process or promoting the programmed cell death machinery of the grape berries can be manipulated by this pathogen during the infection (Veloso and van Kan, 2018).

Many studies have reported the role of *V. vinifera* NAC genes in response to biotic stress. For example, the *VvNAC1* gene (known as *VvNAC60* in Genoscope database) showed resistance to *B. cinerea* and *Hyaloperonospora arabidopsidis* (Le Henanff et al., 2013), *VvNAC36* gene in *A. thaliana* was up-regulated in response to powdery mildew colonization (Tóth et al., 2016).

Therefore, studying NAC transcription factor promoters can reveal useful information about the genes and signaling networks involved in abiotic and biotic stress responses. Web-based databases such as, Genomatix, PLACE and Plant CARE have provided a convenient and easy *in silico* way to search motifs and detect cis-acting regulatory elements (CAREs) of the promoters (Ibraheem et al., 2010). Recently, many studies for elucidation and annotation of gene functions have been carried out by use of *in silico* methods. Fortunately, the past decade has seen a revolution in omics technologies that have generated abundant amounts of useful data for *in silico* function predictions (Rhee et al., 2014).

1.1 Objectives

The NAC proteins play an important role in providing resistance to plants. Till now, few works have been reported on cis-acting regulatory elements present in *Vitis* NAC genes. Therefore, our aim in the present work was to perform a transcriptomic study of the *VvNAC* genes to report their responses against *B.cinerea* through an *in silico* approach by characterizing the cis-acting regulatory elements of the *Vitis vinifera* L. NAC genes, followed by an *in vitro* work wherever is available in order to find a link between both approaches giving results that might be complemented. This work throws light on the promoter regions of grape NAC genes which further can provide new ways for the plant genetic engineering technology for protection of crops against stresses and especially *B.cinerea*.

1. Characterizing of the *Vitis vinifera* L. NAC TF genes through an *in silico* approach.
 - Identifying the cis-acting regulatory elements (CAREs) of the *Vitis vinifera* L. NAC genes.
 - Analyzing the microarray expression profiles of the *Vitis vinifera* L. NAC genes in response to *B. cinerea*.
 - Interpreting the up-regulations' results by linking them to the existence of specific CAREs in the promoters of the up-regulated *VvNAC* genes.
2. Detailed interpretation of the function of the *VvNAC* genes.
 - Detecting the remarkable responses whereas up-regulation or down- regulation of the *VvNAC* genes is induced by *B. cinerea*.
 - Predicting new genes with a function that might respond to stresses and especially to *B. cinerea*.
3. Generating *in vitro* approach.
 - Determination of the subcellular localization through a transient expression for the *VvNAC36* gene in a model tobacco plant, *N. benthamiana* L.
 - Comparing the transient expression results with the computational results to validate or find a link between both approaches.
 -

2. LITERATURE REVIEW

2.1. Plant defence mechanisms

Plants, being sessile organisms, have evolved complex defence mechanisms to protect themselves against a wide array of biotic and abiotic stresses in their environment. These defence mechanisms are the result of millions of years of co-evolution with various stress factors, and they play a critical role in the survival and adaptation of plants (Hönig et al., 2023).

2.1.1. Biotic Stress defence mechanisms

Biotic stresses in plants arise from living organisms, including pathogens, herbivores, and pests. To combat these threats, plants have developed an arsenal of defence mechanisms. One of the most well-known and immediate responses to pathogen attack is the hypersensitive response (HR), characterized by rapid cell death at the infection site. This localized cell death helps to restrict pathogen spread (Pruitt et al., 2021).

HR is often associated with the production of reactive oxygen species (ROS), which can have direct antimicrobial effects. Plant hormones like salicylic acid (SA) play a pivotal role in activating defence responses against pathogens. For instance, SA induces the expression of defence-related genes, such as pathogenesis-related (PR) proteins.

Systemic acquired resistance (SAR) is another defence strategy where the entire plant becomes more resistant to subsequent pathogen attacks following a localized infection, a phenomenon partly mediated by SA. In addition to SA, jasmonic acid (JA) and ethylene (ET) are key players in plant defence against herbivores and necrotrophic pathogens (Yildiz et al., 2021).

These hormones trigger the expression of genes associated with defence against herbivores, such as proteinase inhibitors and volatile organic compounds that attract predators of herbivores. The intricate crosstalk between SA, JA, and ET signalling pathways allows plants to fine-tune their responses based on the nature of the biotic stress.

Furthermore, plants engage in a form of "priming," wherein they pre-activate defence responses in anticipation of future attacks. This priming involves epigenetic modifications and the accumulation of signalling molecules. Once primed, plants can mount faster and stronger defences when challenged by pathogens or herbivores (Hönig et al., 2023).

2.1.2. Abiotic stress defence mechanisms

Abiotic stresses, including drought, salinity, extreme temperatures, and heavy metal toxicity, pose significant challenges to plant growth and productivity. To withstand these stresses, plants have evolved various strategies at the cellular and metabolic levels.

Drought tolerance is a crucial aspect of plant adaptation to arid environments. Plants employ a range of mechanisms to conserve water and cope with dehydration. These mechanisms include stomatal closure, which reduces water loss through transpiration, and the synthesis of compatible solutes like proline and trehalose, which help maintain cellular turgor (Sako et al., 2021).

Dehydration also triggers the production of abscisic acid (ABA), a hormone that induces stomatal closure and activates stress-responsive genes. Salinity stress disrupts cellular ion homeostasis.

To counteract this, plants employ ion transporters and channels to regulate the uptake and compartmentalization of ions, such as sodium and potassium. Additionally, they accumulate osmoprotectants like glycine betaine and mannitol to maintain osmotic balance. In some halophytic plants, specialized salt glands secrete excess salt through trichomes, preventing salt build-up in the plant tissues (Rhaman et al., 2021).

Extreme temperatures, both cold and heat, can damage plant cells and disrupt physiological processes. Plants respond to temperature stress by producing heat shock proteins (HSPs), which assist in protein folding and prevent damage to cellular structures. In cold stress, the production of antifreeze proteins and changes in membrane lipid composition help plants avoid freezing-induced cell damage.

Heavy metal toxicity is a concern in contaminated soils. To mitigate heavy metal stress, plants employ mechanisms such as chelation, where metal-binding molecules like phytochelatins sequester toxic ions, and efflux pumps, which transport metals out of the cell (Kerchev et al., 2020).

2.1.3. Crosstalk between biotic and abiotic stress responses

It's essential to note that plant responses to biotic and abiotic stresses are interconnected. For example, the activation of SA-mediated defence responses against pathogens can lead to increased susceptibility to abiotic stresses like drought. Conversely, JA-mediated defences against herbivores can enhance plant tolerance to some abiotic stresses. Understanding this crosstalk is crucial for developing crop varieties with enhanced stress resilience (Koley et al., 2022).

2.2. Cis-acting regulatory elements of plants

In plants the gene expression is controlled by the existing link between the cis-acting regulatory elements and transcription factors (Ijaz et al., 2020). Thus, the TFs by interacting with the regulatory elements can mediate the target gene activation or repression in a specific tissue or any organ type.

Therefore, recognizing and characterizing these regulatory elements in the genome is important in order to understand more the levels of gene expression (Levine, 2010; Biłas et al., 2016). The CAREs are short linear fragments of non-coding DNA with a variable length, that contain a specific binding site for TFs in the regions of gene promoter (Biłas et al., 2016).

The promoter sequences of a gene are located upstream of the coding region. Through a specific sites in these promoters, the RNA Pol II (RNA polymerase II) enzyme binds to initiate the transcription. Normally, the eukaryotic gene promoter is divided into two regions: core promoter and distal promoter region.

The core promoter is important for initiation the transcription and it consists of: TATA box (present upstream of transcription start site or TSS), TFIIB recognition element (BRE), initiator (Inr) and downstream promoter element (DPE) (Kutach and Kadonaga, 2000; Pandey et al., 2019). The distal promoter or the regulatory region that contains enhancers and silencers, is responsible for gene expression regulation under different conditions (Biłas et al., 2016).

2.2.1. Abiotic stress and CAREs

Plant growth and development are affected by different abiotic stresses such as, drought, temperature and salinity (Zhu, 2002). Thus in order to survive during abiotic stress or any other stresses, plants have made various biochemical and physiological changes to develop a mechanism that detects any growth modifications which may occur through promoting a cascade of signaling pathways controlled by TFs and CAREs, activating several genes for stress tolerance (Gao et al., 2007; Gujjar et al., 2014).

Various cis-acting regulatory elements that showed responded to abiotic stress have been indentified. For example in *Arabidopsis*, dehydration responsive binding element (DREB1, DREB2) regulons and MYB (myeloblastosis)/MYC (myelocytomatosis) regulons plus ABA-responsive element (AREB) and ABA binding factor (ABF) transcription factors (Lenka and Bansal, 2019). The transcription factor NAC was also reported to play a crucial role in abiotic stress. Various transcriptomic studies have identified *NAC* genes responded to drought and salt stresses in the case of soybean, *Oryza sativa* and *Arabidopsis* (Yuan et al., 2019).

2.2.2. Biotic stress and CAREs

In the case of biotic stress, the defense system is complicated whereas different genes are prepared to be expressed to resist any foreign organism. Started by the hypersensitive response (HR) as primary infection fights and followed up by systemic acquired resistance (SAR) as a general protection mechanism for the uninfected parts of plants (Cantu et al., 2013).

The detection of pathogen invasion activates the plant cell defense, followed by an accumulation of salicylic acid (SA) in the cell to change the cytosolic cellular redox potential (Garretón et al., 2002). With these changes the natriuretic peptide receptor 1 (NPR1) protein that is involved in defense mechanisms becomes active and launch the onset of SAR after many steps (Fan and Dong, 2002; Mou et al., 2003).

The gene expression of NPR1 is regulated by TFs through an interaction with W box present in its promoter (Pandey and Somssich, 2009). Similarly, in the case of abiotic stress, TFs also play an important role resisting to biotic stress. For example, in *Arabidopsis* 74 TFs were detected responding to bacterial invasion, in rice with *OsEREBP1* gene that showed an expression in *Magnaporthe grisea* and in tobacco with *SINAC35* gene that improved the resistance to leaf curl virus in transgenic tobacco (Yuan et al., 2019).

2.3. Bioinformatics exploration

2.3.1. Areas of bioinformatics

Bioinformatics can be defined by analyzing and interpreting biological data through the application of a wide range of tools of computation and it is usually an interdisciplinary domain with a mix of computer science, mathematics, and biology that provides different methods in mapping, analyzing and comparing DNA and protein sequences (Luscombe et al., 2001).

Bioinformatics is divided into two complementary subfields: starting with the formation of computational tools and databases, then the application of them in order to produce biological knowledge that helps in better understanding living systems (Xiong, 2009). The first subfield

consists of writing software for sequencing and functional analyzing as well as the constructing biological databases.

These developed tools play key roles in three areas of genomic and molecular biology including sequence, structural and functional analysis (Xiong, 2009). Plenty of examples can be listed to explain the diverse roles in each of these three areas: firstly, for the sequence analysis that includes gene and promoter finding, motif and pattern discovery, sequence database searching and alignment.

Secondly, in the case of structural analysis including the prediction, comparison and analysis of protein and nucleic acid structures. Finally with the functional analysis that contains prediction of the subcellular localization of the proteins, determination of the protein-protein interaction and the gene expression profiling (Rao et al., 2008; Xiong, 2009).

The importance of bioinformatics in generating huge databases and biological knowledge leads to a deep understanding of the genetic and molecular basis of all biological processes in plants, helping the biotechnologists engineering new cultivars firstly with high quality and less economic and environmental costs and resistant to biotic and abiotic stresses (Koltai and Volpin, 2003).

2.3.2. Bioinformatic bases

Bioinformatics encompasses three main elements: databases, computational tools, and software. Databases store vast and complex data, categorized as genome, microarray, and sequence databases (Bourne, 2005). Examples include Xenbase, SEED, ArrayExpress, and Gene Expression Omnibus.

Computational tools play a crucial role in converting these databases into knowledge by utilizing pattern recognition, data mining, and machine learning. Patterns are identified and analyzed through supervised and unsupervised methods, followed by extracting important patterns using techniques like cluster analysis and frequent pattern analysis.

Machine learning, employing statistics and probability theory, helps discover complex patterns. Software provides various programs and tools for database analysis, such as BLAST, SSEARCH, and ENTREZ, which aid in protein and sequence searches, sequence comparison, and retrieving biological information from different databases (Raut et al., 2010).

2.3.3. Bioinformatic applications

As a powerful tool, bioinformatics provide the possibility to study how the cellular activities are altered in different developmental phases, thus help the biologists to increase the understanding of biological processes with computationally intensive techniques including sequence alignment, gene finding, genome and gene expression analysis.

In sequence alignment, the word sequence is defined by the nucleotide sequence present in DNA, RNA and amino acids in protein. In addition, alignment indicates the possibility of changes between the two homologous sequences and a common ancestor sequence that could have occurred during evolution.

The aim of the alignment is to generate a comparison between nucleotides sequences in order to find similarity (Mount et al., 2001). Therefore, a similarity that can be explained by the existence of the same functions between the sequences or by verifying homology due to the origin from a

common ancestor. The multiple sequence alignments option is frequently used recently in different sequence analysis areas because it is more accurate because of the combination process between three-dimensional structure information with primary sequences (Wallace et al., 2005). Once the sequence has been obtained, gene finding through databases is a crucial step in understanding the genome of a species (Korf, 2004).

Therefore, starting by identification genes and predicting their functions is important in any study in order to analyze a genome of an organism. Followed by comparison of the identified genes with genome of other organisms to find if they share similarity in different functions such as resistance against various diseases or stresses (Raut et al., 2010).

The gene functions and expressions have always been a target for biotechnologists to analyze, due to their importance in which the gene information is used for synthesis of proteins. For this purpose they created various techniques and tools such as, DNA Microarray, SAGE and Tilling array (Raut et al., 2010).

The synthesis of proteins or the process from DNA to RNA and then to protein, is called the central dogma of molecular biology. The process is started in the nucleus with transcription when the genes are active they produce RNA after copying coding and non-coding sequences. Followed in the cytoplasm with the translation to produce protein from the RNA after many steps. Thousands of messenger RNAs are produced when a single gene is regulated, encoding different proteins and each one of them has a specific role to play (Raut et al., 2010).

In the case of the microarray, thousands of DNA molecules that have been taken from the cells in different conditions are spotted on a glass slide as array followed by pouring a solutions of fluorescently labeled DNA or RNA over the array and each molecule in the solution searches for matching partner on the surface (Raut et al., 2010).

2.3.4. Bioinformatics and stresses

Bioinformatics plays an indispensable role in elucidating the molecular intricacies of plant responses to both biotic and abiotic stresses, providing valuable insights into stress tolerance mechanisms and enhancing crop resilience (Jiang & Tyler, 2012).

In the realm of biotic stresses, bioinformatics aids in the identification and characterization of pathogen effectors and resistance genes within plant genomes, facilitating the development of disease-resistant crop varieties. High-throughput omics technologies, such as RNA-seq and mass spectrometry, enable comprehensive analyses of transcriptomes and proteomes during plant-pathogen interactions, with bioinformatics pipelines essential for the interpretation of large-scale data (Bigeard et al., 2015).

Additionally, metabolomics studies, guided by bioinformatics tools, shed light on the metabolic adjustments that bolster plant defence. In the context of abiotic stresses, bioinformatics leverages genome-wide association studies (GWAS) to identify genetic loci associated with stress tolerance, facilitating the selection of resilient crop genotypes. Functional genomics studies, driven by bioinformatics, unravel gene functions under abiotic stress conditions, including the role of transcription factors (Shinozaki et al., 2022).

Machine learning and predictive models, integrating multi-omics data, enable the forecasting of stress responses and aid in the selection of stress-tolerant cultivars. Notably, bioinformatics promotes the integration of biotic and abiotic stress response data, unveiling shared regulatory networks and crosstalk between these pathways (Kole et al., 2015). As technology advances, bioinformatics remains pivotal for unravelling the intricate web of plant stress responses, contributing to sustainable agriculture and global food security.

2.3.5. Bioinformatics examples

Bioinformatics web tools and databases are crucial components of modern molecular biology and genomics research. PLANT CARE, for instance, significantly contributes to the understanding of gene expression and genome characterization by enabling the identification of cis-acting regulatory elements (CAREs) in plant genes (Lescot et al., 2002).

These CAREs are essential components of gene regulation, and their discovery via PLANT CARE informs researchers about the mechanisms controlling gene expression (Lescot et al., 2002). For example, a researcher studying stress responses in a particular plant species may use PLANT CARE to identify specific CAREs associated with stress-related genes.

MEME, on the other hand, plays a pivotal role in elucidating conserved sequence motifs within DNA, RNA, and protein sequences (Bailey et al., 2009). These motifs often serve as binding sites for transcription factors or other regulatory elements, providing insights into gene expression regulation (Bailey et al., 2009). For instance, MEME can be employed to analyze a set of co-regulated genes and discover common motifs, such as promoter elements shared among genes activated during a stress response.

Moreover, Expression Atlas databases, like the one hosted at EBI, are invaluable resources for studying gene expression and genome characterization (Papatheodorou et al., 2020). These databases compile high-throughput gene expression data from various experiments and organisms, allowing researchers to explore how genes are expressed under different conditions, tissues, or stressors (Papatheodorou et al., 2020).

For instance, a scientist investigating the response of a particular set of genes to a specific pathogen could leverage Expression Atlas to examine the expression patterns of these genes across various infection time points or in different tissues, thereby gaining insights into their regulatory networks and functions.

Additionally, NCBI Gene Expression Omnibus (GEO) and ArrayExpress serve as expansive repositories of high-throughput gene expression data, permitting comparisons of experimental results with publicly available datasets.

Ensembl contributes not only genomic sequences but also RNA-seq data, enabling researchers to explore gene expression across diverse tissues and developmental stages. Focused on developmental biology, the Gene Expression Database (GXD) meticulously catalogues spatial and temporal gene expression patterns during mouse embryo development (Bult et al., 2019).

The STRING Database predicts protein-protein interactions, offering insights into gene expression networks and functional relationships (Szklarczyk et al., 2019). DAVID (Database for Annotation, Visualization, and Integrated Discovery) aids in interpreting experimental outcomes by performing gene ontology enrichment and pathway analysis (Huang et al., 2009).

Furthermore, Genevestigator provides a comprehensive platform for exploring gene expression across various plant and animal species, facilitating the study of tissue-specific expression and responses to different treatments or stressors (Hruz et al., 2008).

In conclusion, these bioinformatics tools and databases play pivotal roles in advancing our understanding of gene expression and genome characterization, and their applications are diverse, ranging from uncovering regulatory elements to discovering conserved motifs and elucidating gene expression patterns under various conditions.

2.4. NAC transcription factors

The NAC is one of the largest TF families in plants that associated with different biological and developmental processes like senescence (Kjaersgaard et al., 2011; Yang et al., 2011), formation of secondary walls (Zhong et al., 2010) and biotic (Christianson et al., 2010) and abiotic stress response (Tran et al., 2010). Plant biotechnologists have succeeded to identify plenty of NAC genome sequences in plant species (Tran et al., 2004; Fujita et al., 2004). For example, they found 117 NAC genes in *Arabidopsis*, 151 in rice 163 in poplar, 152 in tobacco (*Nicotiana tabacum*) and 74 in grape (*Vitis vinifera* L.) (Nuruzzaman et al., 2010; Rushton et al., 2008).

2.4.1. Structure of NAC proteins

Normally, the structure of NAC protein consists of two terminals: N terminal NAC domain with almost 150 amino acids and C terminal. The NAC domain in his turn is divided into 5 subdomains (A to E). The subdomains C and D are highly conserved positively charged that bind to DNA, while for the subdomain A that might be involved in the formation of functional dimer and for subdomains B and E with their different roles in NAC genes functional diversity (Chen et al., 2011; Ooka et al., 2003).

The C terminal contains the transcription regulatory regions (TRRs) that can activate or repress the transcription. The TRR has different specific motifs that are rich in repeats of proline-glutamine, serine-threonine or acidic residues. For example, ten C terminal motifs were found in the rice NAC proteins (Fang et al., 2008), but it should be taken in consideration that these motifs may vary from subfamily to another one (Shen et al., 2009).

Due to the high degree of intrinsic disorder (ID) of TRRs, they are capable to interact with different target proteins (Jensen et al., 2010). Thus, some NAC proteins have ability for protein binding in their TRRs (Kim et al., 2007; Kleinow et al., 2009), like the case of the NTLs; NAC proteins with a helical transmembrane (TM) motif the responsible for plasma membrane anchoring that may have also regulatory roles under different environmental conditions (Seo et al., 2008).

2.4.2. NAC regulation

The gene is firstly regulated in transcription after binding of TFs to its promoter regulator region (Nakashima et al., 2012). Moreover, the cis-acting regulatory elements (CAREs) present in the promoter region including LTREs (Low-temperature responsive elements), MYB (Myeloblastosis) and MYC (Myelocytomatosis) binding sites, ABREs (ABA-responsive elements), DREs (Dehydration responsive elements) and W-box etc., are responsible for the regulation of the NACs stress responsive genes (Tran et al., 2010).

For example, different studies have reported that *OsNAC3* gene is expressed under abiotic stress due to the presence of DREB (DRE binding) regulon (Takasaki et al., 2010), while the expression of *ONAC045* gene under same conditions was linked to the presence of MYB and MYC core binding sites in the promoter (Zheng et al., 2009).

Secondly, the genes undergo to a post-transcriptional regulation through micro-RNA (miRNA)-mediated cleavage. Therefore, the most recognized miRNA is miRNA 164 that showed high affinity in developmental and stress regulation, it regulates the post-transcriptional RNA processing of *NAC* TF genes (Khraiweh et al., 2011).

Finally, the *NAC* TF genes are subjected to a post-translational regulation which includes protein degradation mediated by ubiquitins, dimerization (Hegedus et al., 2003) and interaction with other non-*NAC* proteins (Greve et al., 2003).

2.4.3. *NAC* genes expression

The role of *NAC* TF genes have been reported in many studies, that verified the expression of these genes at least under one type of stress like salinity, drought, fungi etc. (Fang et al., 2008). The analysis of the *NAC* gene expression by using microarray has shown for example, in the case of rice more than 45 *NAC* genes were induced in abiotic and 26 in biotic stresses (Nuruzzaman et al., 2010). Moreover, in *Arabidopsis* the majority of the *NAC* genes expressed under salt and high temperature stresses (Zeller et al., 2009).

In addition, the expression level of *NAC* genes also has been reported in response to attacks of different bacteria, viruses and fungi (Ren et al., 2000; Oh et al., 2005) and by the application also of the exogenous phytohormones like abscisic acid (ABA), ethylene (Et) and salicylic acid (SA) and jasmonic acid (JA) in different species (Yoshii et al., 2010). The phytohormones are associated with a cascade of signaling response in the case of any stresses by acting in conjunction or opposition to each other to maintain cellular homeostasis (Fujita et al., 2006).

2.4.4. Abiotic and biotic stresses response

Plenty of positive results have demonstrated the role of *NAC* TFs in response to abiotic stress after overexpression in *Arabidopsis* and rice without showing any growth retardation (Liu et al., 2011). Whereas 38 *NAC* genes were found to be involved in soybean drought response (Le et al., 2011), and 40 *NAC* genes were involved in rice drought response (Le et al., 2011); (Fang et al., 2008). Furthermore, 33 *NAC* genes in *Arabidopsis* responded to salt stress (Jiang and Deyholos 2006).

For example in rice, drought, salt tolerance and higher seed production in dry field showed by the *SNAC1* transgenic lines (Hu et al., 2006) and same in the case of *OsNAC10* (Jeong et al., 2010; Song et al., 2011). In transgenic *Arabidopsis* transformed with the wheat *NAC* transcription factor *TaSNAC4-3A* gene demonstrated drought tolerance (Mei et al., 2021). Overexpression of *ANAC019*, *ANAC055/AtNAC3*, or *AtRD26* enhanced drought tolerance in *Arabidopsis* (Tran et al., 2004). In soybeans, the role of the *GmNAC8* gene as a drought-stress positive regulator was also observed (Yang et al., 2020).

Conversly, dwarfing, late flowering and lower seed yield in the transgenics have been identified after *NACs* overexpression in some cases but it might be solved by utilizing stress-inducible or tissue-specific promoters like *OsNAC6* or *RCc3* (Nakashima et al., 2007).

Biological stress arises when a number of viruses and living organisms, such as bacteria, fungi, and harmful insects, invade and damage plants. As a result, the plant will activate a complex defense mechanism in response (Masri and Kiss 2023). Moreover, RNA interference/knockouts and overexpression studies have revealed the function of NAC TFs in various plant–pathogen interactions (Collinge and Boller 2001).

Numerous NAC proteins are known to interact directly with virus-encoded proteins to either enhance or inhibit virus replication (Yoshii et al., 2009). Despite the significance of *NAC* TFs in stress responses, there are few studies indicating *NAC* gene responses to pathogens. The most well-known researches were undertaken in rice, where the value of the *ONAC122* and *ONAC131* genes in *M. grisea* resistance responses was discovered using virus-induced gene silencing (VIGS) (Sun et al., 2013).

Furthermore, the same (VIGS) method was used to show that *TaNAC35* acts as a negative regulator for leaf rust resistance in a compatible relationship between common wheat and *Puccinia triticina* in wheat (Zhang et al., 2021). Additionally, overexpression of *ONAC066* gene in transgenic rice positively regulated blast and bacterial blight resistance by inhibiting ABA signaling pathways (Liu et al. 2018), and with *OsWRKY67* gene it positively regulates the same disease by direct activation of PR genes (Liu et al., 2018).

Finally, in *Arabidopsis*, overexpression of *ATAF1* reduced resistance to the necrotrophic fungi *Botrytis cinerea* and *Alternaria brassicicola*, suggesting that *ATAF1* is a negative regulator of necrotrophic pathogen defense (Wang et al., 2009; Wu et al., 2009).

2.4.5. *NAC* TFs in multiple processes

A crosstalk signaling between abiotic and biotic stresses have been identified in *NAC* TFs. Thus, the responses against these stresses might be discrete from the transgenic plants after overexpression of the *NAC* genes. For example, in *Arabidopsis*, the overexpression of *ATAF1* have reported to increase the drought tolerance and at same time the sensitivity to ABA, salinity, necrotrophic fungus (*B. cinerea*).

Thus, it negatively regulates ABA levels for efficient basal defense against *Blumeria graminis f.sp. graminis* (*Bgh*) (Wu et al., 2009). Moreover, *ANAC019* and *ANAC055* were associated with drought tolerance, but their overproduction also decreased resistance to *B. cinerea*. Finally, with the overexpressing of *ATAF1* in rice, *OsNAC6* that showed a slight increase of the resistance to the blast disease in addition to higher tolerance to drought and high salinity (Fujita et al., 2004).

As plants often encounter multiple stresses concurrently, such multi-functionality becomes significant for survival under extreme stress conditions. Several *NAC* TFs also integrate responses to environmental stresses into modulation of plant development processes, like lateral root development, seed germination (Balazadeh et al., 2010), flowering (Kim et al., 2007) and senescence. Such versatility of *NAC* functions may have evolved to ensure plants' longevity, survival and reproductive success under environmental stresses.

2.5. *Vitis vinifera* L.

2.5.1. Description

Vitis vinifera L. (grapevine), a member of the *Vitaceae* family, is a deciduous, woody vine with a southwestern Asian origin was transported by the European to many places over the world. Normally, they can grow to form long spiral tendrils up to 150 cm long and spread horizontally over low-growing bushes and they will be trimmed much smaller down to 20 cm in the case of plants grown for wine production.

Additionally, they start blooming during the period between May and June with small hermaphroditic flower that form very soft pulpy grapes in different sizes and colors which ripen in summer. *Vitis vinifera* L. is one of the world's most valuable fruit crops, with high economic, nutritional value, and a reputation for producing high-quality wines (Ju et al., 2020).

For instance, Pinot Blanc, Pinot Gris, Sauvignon Blanc and Chardonnay in the case of white wine varieties and Cabernet Sauvignon, Merlot and Pinot Noir for the red wine varieties, but beside these very famous grape cultivars the number of varieties has increased to about 10 000 during the centuries of viticulture.

2.5.2. Abiotic and biotic stresses

As mentioned before considering the grape as one of the largest and most important agricultural fruit crop in the world, and, like other crop, grapevines are permanently challenged by various pathogens and changing environmental conditions that cause significant yield and quality losses (Masri and Kiss 2023). Starting firstly by abiotic stresses, such as light, temperature, and drought. The grapevines undergo chemical, physical, and physiological changes in order to resist these stresses and require intensive growing conditions for producing fruit and wine of high quality.

Grapes are susceptible to many insects, disease pests and viruses particularly in humid summer climates including black rot, powdery mildew, gray mold, crown gall, botrytis bunch rot, phylloxera, grape berry moth and Japanese beetle. Several grapevines recognize the pathogen components, transduce the stress signal, and induce a defense response through hormones regulation.

Therefore, many studies have reported the role of *V. vinifera* genes regarding the response to abiotic and biotic stresses (Fang et al., 2016). For example, *VvNAC1* (*VvNAC60*) overexpression increased the osmotic, salt, and cold stress resistance in *A. thaliana* (Le Henanff et al., 2013). Overexpression of *VvNAC17* gene isolated from grapevine improved resistance to salinity, freezing, drought and also regulates an ABA-mediated pathway in *A. thaliana* (Ju et al., 2020) and the overexpression of grapevine *VvNAC08* in transgenic *Arabidopsis* increased drought tolerance (Ju et al., 2020).

Additionally, the public microarray data showed that the expression of *VvNAC26* was highly induced under water deficit, cold temperature, and high salinity stresses (Wang et al., 2013). A recent study proved a remarkable regulation of the *VvNAC36* gene in *A. thaliana* in response to powdery mildew colonization, the up-regulation of *VvNAC36* 1.5-fold in the case of powdery mildew (PM) while no response was detected for salicylic acid (SA) (Tóth et al., 2016). Finally, the overexpression of *VvNAC1* (*VvNAC60*) increased the resistance to *B. cinerea* and *Hyaloperonospora arabidopsidis* (Le Henanff et al., 2013).

2.5.3. *Botrytis cinerea*

Botrytis cinerea (*B.cinerea*) is one of the most crucial necrotrophic fungal pathogens in grapevine and in more than 200 plant species in different temperate or subtropical regions causing gray mold disease, pre and postharvest decay and fruit quality damaging in fields, greenhouses and even during storage (Williamson et al., 2007).

B.cinerea infection may occur slowly to wide range of hosts like the horticultural crops and fruit crops in their flowers, leaves, shoots and fruits. These symptoms are visible where the fungus begins to rot the plant at wound sites (Sutton 1998).

Two different types of mold disease are identified on the infected grapes. The first is noble rot, that occurs in drier condition following by wetter one resulting of accumulation of aroma and sugar concentration (Breia et al., 2021). The second one is grey rot or bunch rot, caused by heavy rainfalls and high humidity conditions leading to severe losses of the infected bunches (Gubler et al., 2013). Conversely to noble rot, the grey rot affects the production of wine badly by negatively modifying the fermentation process thus the sensory properties of the final product (Hornsey, 2007; Morales-Valle et al., 2011).

Regarding the two types, the rot development mechanism in the infected grape leading to the formation of noble or grey rot has not been well clarified yet but it might be influenced by the intrinsic characteristics of the grape cultivar or different environmental effects (Sipiczki, 2006).

2.5.4. *Applied breeding methods and objectives in Vitis vinifera* L.

Vitis vinifera L., has been a focal point of applied breeding efforts for centuries due to its economic importance in wine and table grape production. These breeding programs aim to develop grapevine cultivars with improved traits such as disease resistance, yield, fruit quality, and adaptability to changing environmental conditions. Several breeding methods and objectives have been employed to achieve these goals.

Breeding methods:

1. **Conventional Breeding:** Conventional breeding, also known as classical or traditional breeding, involves the controlled cross-pollination of two grapevine varieties with desirable traits. The resulting progeny are evaluated for specific characteristics, and the best-performing individuals are selected as potential new cultivars (Emanuelli et al., 2013).
2. **Marker-Assisted Breeding:** With advancements in genomics, marker-assisted breeding has become a powerful tool. DNA markers linked to important traits, such as resistance to pests or diseases, can be identified and used to select promising individuals at an early stage, reducing the time and resources required for breeding (Bettoni et al., 2021).
3. **Genome Editing:** Emerging technologies like CRISPR-Cas9 have the potential to revolutionize grapevine breeding. Genome editing allows for precise modification of genes responsible for specific traits, offering the opportunity to create customized grapevine varieties with improved characteristics (Ren et al., 2016).

Breeding objectives:

1. **Disease Resistance:** One of the primary objectives in *Vitis vinifera* L. breeding is enhancing resistance to diseases like downy mildew (*Plasmopara viticola*) and powdery mildew (*Erysiphe*

necator). Developing cultivars with innate resistance reduces the reliance on chemical treatments (Feechan et al., 2013).

2. **Abiotic Stress Tolerance:** Climate change and shifting environmental conditions pose challenges for grape cultivation. Breeding for abiotic stress tolerance, such as drought resistance, ensures stable grape production in the face of changing climates (Cramer et al., 2007).

3. **Improved Fruit Quality:** Enhanced fruit quality attributes, including sugar content, flavor compounds, and aroma profiles, are essential for wine and table grape varieties. Breeding programs target these traits to meet consumer preferences and market demands (Ryan et al., 2015).

4. **Yield and Productivity:** Increasing grapevine yield without compromising fruit quality is a consistent breeding objective. High-yielding varieties contribute to the economic sustainability of vineyards (de Oliveira et al., 2020).

5. **Adaptation to New Regions:** As grapevine cultivation expands to new regions, breeding objectives include developing cultivars that can thrive in diverse soil types and microclimates, allowing for more extensive grape production (This et al., 2006).

In conclusion, applied breeding methods and objectives in *Vitis vinifera* L. are driven by the need for sustainable and resilient grape production. Through conventional breeding, marker-assisted techniques, and emerging genome editing technologies, breeders work toward creating grapevine cultivars with improved disease resistance, abiotic stress tolerance, fruit quality, yield, and adaptability. These efforts ensure the continued success of grape cultivation in a dynamic and evolving agricultural landscape.

2.6. *In silico* methods: Definition, applications, and objectives

Definition:

In silico methods in plant biotechnology and molecular biology refer to computational techniques and simulations used to analyze, model, and predict biological processes and phenomena in plants. The term "*in silico*" is derived from Latin, meaning "in silicon," signifying that these methods are conducted in a computer-based, virtual environment. In the context of plant science, *in silico* methods involve the application of computer algorithms and data-driven approaches to understand, study, and manipulate various aspects of plant biology

Applications:

1. **Genome Analysis:** *In silico* methods are widely applied in plant genome analysis. This includes genome sequencing, annotation, and comparative genomics. Researchers can predict genes, identify regulatory elements, and explore genetic diversity within plant genomes using computational tools (Edwards and Batley, 2010).

2. **Functional Genomics:** These methods are crucial for interpreting high-throughput data generated by techniques like transcriptomics and proteomics. Researchers can identify and characterize genes, pathways, and regulatory networks associated with plant development, responses to stresses, and other biological processes (You, 2018).

3. **Protein Structure and Function:** *In silico* methods play a significant role in predicting the structure and function of plant proteins. Predictive modeling, molecular docking, and structural

bioinformatics aid in understanding protein interactions, enzymatic activities, and potential drug targets within plants (Kumari et al., 2013).

4. Crop Improvement: In plant breeding and crop improvement programs, *in silico* methods are used to identify candidate genes associated with desirable traits such as disease resistance, yield, and nutritional quality. These methods accelerate the breeding process by narrowing down potential targets (Schreiber et al, 2018).

5. Metabolic Pathway Engineering: Researchers utilize *in silico* approaches to analyze and engineer plant metabolic pathways. Constraint-based modeling and metabolic flux analysis aid in optimizing metabolic networks for the production of biofuels, pharmaceuticals, and other valuable compounds (Araus et al., 2022).

6. Systems Biology: *In silico* modeling is integral to systems biology, where researchers seek to understand the behavior of entire plant systems. These models integrate data from various omics levels (genomics, transcriptomics, metabolomics) to simulate and predict plant responses to changing environmental conditions (Atkin and Macherel, 2009).

Objectives:

1. Hypothesis Testing: *In silico* methods allow researchers to test hypotheses and generate predictions. For example, they can predict the effects of specific genetic modifications on plant phenotypes or assess the impact of environmental factors on gene expression (Tutar, 2014).

2. Data Integration: Given the wealth of biological data generated in plant research, one objective of *in silico* methods is to integrate and make sense of diverse data types. This integration helps researchers gain a holistic view of plant biology (Walls et al., 2012).

3. Model Construction: Researchers aim to construct accurate mathematical and computational models of plant processes. These models help simulate and understand the dynamic behavior of plants, from growth and development to responses to biotic and abiotic stresses (Gonzalez-Meler et al., 2014).

4. Crop Enhancement: In agriculture and plant biotechnology, the objective is to use *in silico* methods to identify genes and traits that can enhance crop productivity, nutritional content, and stress tolerance. This contributes to global food security and sustainable agriculture (Cai et al., 2018).

5. Drug Discovery: In the context of medicinal plants, *in silico* methods are employed to discover bioactive compounds and understand their interactions with human receptors. This objective supports the development of plant-based medicines (wan et al., 2019).

6. Environmental Impact Assessment: Researchers use *in silico* methods to assess the ecological and environmental impact of genetically modified plants or novel agricultural practices. This contributes to responsible biotechnology adoption (Denby et al., 2004).

In conclusion, *in silico* methods are indispensable tools in plant biotechnology and molecular biology. Their applications span from fundamental research to practical applications in crop improvement, drug discovery, and environmental sustainability. The objectives of *in silico* methods encompass hypothesis testing, data integration, model construction, and the enhancement of plant traits, ultimately contributing to our understanding and manipulation of plant biology for diverse purposes

3. MATERIALS AND METHODS

3.1. *In silico* part

3.1.1. *Detection and data analyses of NAC family genes*

A local blast analysis was conducted to locate *NAC* genes within the grapevine genome. The NAM domain and 26,346 predicted genes from the 12X assembled *Vitis vinifera* Pinot Noir PN40024 genome were utilized as search queries and databases. As a result, 74 genes were successfully identified and designated from *VvNAC01* to *VvNAC74*, each associated with a unique gene locus number (Wang et al., 2013).

In this study, we utilized as input the gene loci of the 74 *Vitis vinifera* *NAC* genes to retrieve the protein sequences from genoscope, the grapevine genomic database available at (<http://www.genoscope.cns.fr/externe/GenomeBrowser/Vitis/>).

The *NAC* conserved domain sequences of all the genes were extracted through the Pfam web database (<https://pfam.xfam.org>) (Finn et al., 2016). Subsequently, a BLASTp analysis was performed using the phytozome v13 database accessible at <https://phytozome-next.jgi.doe.gov/> to validate all *VvNAC* protein sequences employed in this approach (Goodstein et al., 2012).

The amino acid composition, isoelectric point, and molecular weight of the *NAC* proteins were determined using ExPASy; http://ca.expasy.org/tools/pi_tool.html (Ison et al., 2013). CELLO v.2.5 webtool; <http://cello.life.nctu.edu.tw> was utilized to estimate the subcellular localization of the proteins (Yu et al., 2004).

TMHMM v2.0 (<http://www.cbs.dtu.dk/services/TMHMM/>) was used to determine the transmembrane helices (TMHs) of the identified *VvNAC* family proteins (Krzyszowski et al., 2009).

TMHs are regions within a protein that span the lipid bilayer of a cell membrane. By analyzing the amino acid sequence of the *VvNAC* genes, TMHMM v2.0 uses computational algorithms and statistical models to identify potential TMHs within the protein. The prediction is based on specific features such as hydrophobicity and amphipathicity, which are characteristic of transmembrane regions.

The nucleotide substitution rates (K_a and K_s) and their ratios (K_a/K_s) for duplicated *VvNAC* genes were computed using the straightforward K_a/K_s calculator feature within the TBtools software, which can be found at <https://bio.tools/tbtools> (Chen et al., 2020). To facilitate this analysis, the coding sequence (CDS) sequences of the genes were sourced from the phytozome v13 database.

3.1.2. *Conserved motifs and cis-acting regulatory elements (CAREs) of VvNAC genes*

The protein motifs of *Vitis vinifera* *NAC* were analyzed using the web database program MEME 5.5.2, which can be accessed at <https://meme-suite.org/meme/tools/meme>. This program utilizes various statistical modeling techniques to identify prevalent patterns, 10 Motifs were selected based on the criteria of a minimum width of 6 pixels and a maximum width of 50 pixels (Bailey et al., 2009).

The cis-acting regulatory elements of the *VvNAC* genes were identified using the Plant CARE database, which is accessible at <http://bioinformatics.psb.ugent.be/webtools/plantcare/html/> (Lescot et al., 2002) (Appendix - Table 7.1).

3.1.3. Protein–protein interaction analysis

In this study, we utilized the web-based STRING (Search Tool for the Retrieval of Interacting Genes/Proteins) database version 11.0 <https://string-db.org> to predict and construct protein-protein interaction networks involving VvNAC proteins, with *Arabidopsis* homologous proteins as references (Szklarczyk et al., 2023).

To configure the STRING analysis, we adjusted specific parameters as follows: the network type was defined as the full STRING network, the meaning of network edges was set to evidence, the minimum required interaction score was established as a medium confidence parameter (0.4), and the maximum number of interactions displayed for each protein did not exceed 10 interactors.

3.1.4. In silico expression profiles of VvNAC genes and statistical analysis

The microarray dataset of grapevine NAC gene expression in response to *Botrytis cinerea* stress was obtained from the open science resource Expression Atlas, which can be accessed at <https://www.ebi.ac.uk/gxa/experiments/E-GEOD-67932/Results> (Blanco-Ulate et al., 2015) (Appendix - Table 7.4). Expression patterns of these NAC genes were represented as a heat map viewed in TBtools <https://bio.tools/tbtools>.

Additionally, by using [Bioinformatics.com.cn](https://www.bioinformatics.com.cn) a statistical analysis for gene expression through hierarchical clustering data was generated.

3.2. In vitro part

3.2.1. Construction of plant expression vectors

VvNAC36 gene construct

Based on previous work started in the institute, the VvNAC36 has been chosen to check how much accurate the *in silico* results might be to compare to the laboratory work. Thus, genomic DNA was extracted from young grapevine leaves using the CTAB (Cetyltrimethylammonium bromide) method by following these steps:

1. Sample collection: We begin by collecting plant material, such as leaves, young shoots, or other tissues of interest, ensuring that the material is fresh and free from contamination.
2. Homogenization: We grind the collected plant material into a fine powder, using either a mortar and pestle or specialized homogenization equipment. Our goal is to break open the cells and release their contents.
3. Cell lysis: We transfer the powdered plant material into a tube or container and add a CTAB-based extraction buffer. The extraction buffer typically contains CTAB, EDTA (Ethylene Diamine Tetraacetic Acid), and other reagents to disrupt cell membranes and stabilize DNA. We then heat the mixture to facilitate cell lysis.
4. Incubation: After adding the extraction buffer, we incubate the mixture at an elevated temperature, usually around 65°C, for a specified period. This step helps break down cell membranes and release DNA into the solution
5. Phenol-Chloroform extraction: Following incubation, we add an equal volume of phenol-chloroform-isoamyl alcohol (PCI) to the mixture. Phenol-chloroform helps to separate DNA from

proteins, lipids, and other cellular components. We mix the solution thoroughly by inverting the tube or using a vortex mixer.

6. Centrifugation: We then centrifuge the mixture at high speed. This separates it into three phases: a lower organic phase (containing proteins and lipids), an interphase, and an upper aqueous phase (containing DNA). Carefully, we transfer the upper aqueous phase to a new tube.

7. Precipitation: To precipitate DNA, we add cold isopropanol or ethanol to the aqueous phase. DNA is not soluble in these alcohols, causing it to precipitate out of solution. We gently invert the tube to encourage DNA precipitation.

8. Centrifugation: After precipitation, we centrifuge the tube again to pellet the DNA. We then remove the alcohol-containing supernatant, taking care not to disturb the DNA pellet.

9. Washing: We wash the DNA pellet with cold ethanol to remove any residual contaminants, which further purifies the DNA.

10. Drying: We allow the DNA pellet to air dry for a brief period, ensuring it's completely dry to avoid ethanol contamination in downstream applications.

11. Rehydration: To dissolve the DNA, we rehydrate the DNA pellet in an appropriate buffer, such as TE buffer (Tris-EDTA). This buffer helps protect the DNA from degradation.

12. Storage: Finally, we store the extracted DNA at -20°C or -80°C for long-term preservation or at 4°C for short-term use.

Moreover, the *VvNAC36* gene was then amplified by PCR using Phusion High Fidelity Taq polymerase with forward and reverse primers, which contained *HindIII* and *SacI* restriction sites, respectively (Appendix- [Table 7.6, 7.7](#)).

The PCR amplification was performed for 34 cycles under the following conditions: 98°C for 30 s, 94°C for 10 s, 60°C for 90 s, and 72°C for 5 min, followed by a final extension step at 72°C for 10 min. The purified PCR product was cloned into the pENTR/D-TOPO vector from Invitrogen, further digested, and subcloned at the *HindIII* and *SacI* restriction sites into the pGWB605 binary vector, resulting in the recombinant plasmid pGWB605::*VvNAC36* (Appendix- [Figure 7.1](#)).

The pGWB605 vector contained a CaMV35S promoter and the synthetic green fluorescent protein (sGFP) reporter gene, as shown in a schematic structure using Benchling software ([Figure 3.2.1](#)).

The control vector 35S-sGFP (containing green fluorescent protein alone) and the recombinant plasmid pGWB605::*VvNAC36* were introduced into *Agrobacterium tumefaciens* strain GV3101 by direct transformation.




Figure 3.2.1. Schematic structure of the vector pGWB605 using Benchling software.

3.2.2. Transient Expression

For the pre-culture, *Agrobacterium tumefaciens* (*A. tumefaciens*) GV3101 carrying the binary vector pGWB605::VvNAC36 and 35S-sGFP were grown overnight in 2–3 mL Luria-Bertani medium (LB medium) supplemented with 100 mg/L spectinomycin and 30 mg/L rifampicin at 28 °C (Gusain et al., 2021). Subsequently, 1 mL of the pre-cultures was inoculated into fresh 25 mL LB medium containing the same antibiotics for the main cultures.

On the following day, culture solutions were centrifuged at 5,000 rpm for 15 minutes at room temperature, followed by resuspension of the *A. tumefaciens* pellet in a buffer consisting of 1 M MgCl₂, 0.5 M 2-(N-morpholino)ethanesulfonic acid (MES) (pH 5.6), and 10 mM acetosyringone. The buffer volume was adjusted to achieve an OD 600 of approximately 1 (Hoshikawa et al., 2018). The buffer was then incubated at room temperature for 2–3 hours, as shown in Photo B (Figure 3.2.2).

Using a needleless 1-mL syringe, the cells were co-infiltrated at a 1:1 ratio with *A. tumefaciens* containing the p19 gene, a silencing suppressor (Lindbo, 2007), into the abaxial side of the leaves of 6-week-old plants represented by the photos marked as A, C, and D (Figure 3.2.2).

3. Materials and methods

Following infiltration, the plants were left undisturbed for 5 days before examination with the Leica TCS SP8 confocal microscope. This examination involved analyzing pieces of infiltrated leaves taken from the infected region at the National Agricultural Research and Innovation Centre (NARIC) in Hungary.




Figure. 3.2.2 Agroinfiltration of tobacco leaves. (A) 6-week-old *N. benthamiana* plant that is appropriate to be infiltrated. (B) *Agrobacterium tumefaciens* containing the binary vector in the infiltration buffer. (C) The suspension was slowly infiltrated into the leaves. (D) The colour of the infiltrated region changed to deeper green.

4. RESULTS

4.1. Identification and genomic distribution of NAC proteins

In this study, different databases were utilized to study and characterize the *Vitis vinifera's* NAC proteins. The genome of *V. vinifera* L. was found to have 74 NAC proteins. [Table 7.1](#) in appendix presents a comprehensive summary of the *VvNAC* genes, including their length, molecular weight (MW), isoelectric point (pI), and cellular location. The length of the proteins produced by NAC genes ranged from 128 amino acids (AA) in *VvNAC50* to 994 (AA) in *VvNAC54*. The molecular weight of these proteins varied from 14.76 to 110.67 kDa (*VvNAC50* and *VvNAC54*, respectively). The isoelectric point of the NAC proteins ranged from 4.1 in *VvNAC47* to 10.52 in *VvNAC73*.

4.2. TMHs and KaKs value of *VvNAC* genes

Notably, nine out of the 74 *VvNAC* genes identified encoded proteins with transmembrane helices (TMHs). It was found that eight proteins, namely *VvNAC15*, *VvNAC20*, *VvNAC25*, *VvNAC29*, *VvNAC48*, *VvNAC54*, *VvNAC67*, and *VvNAC71* contained only one transmembrane domain while *VvNAC72* contained 6 domains ([Table 4.2-A](#)).

Therefore, this information is valuable because the presence and location of transmembrane helices can provide insights into the protein's function and its association with cellular membranes. It suggests that these *VvNAC* proteins are likely to be membrane-associated or have specific interactions with the cell membrane.

Based on the analysis conducted using the KaKs calculator via TBtools, none of the 20 tandem repeat sequences exhibited Ka/Ks values greater than 1 ([Table 4.2-B](#)). This outcome implies the absence of positive selection in any of the tandem repeat sequences, which are typically associated with facilitating adaptive genetic variation and potentially influencing species evolution.

Instead, all genes displayed Ka/Ks values below 1, indicating the prevalence of negative selection during the course of evolution. This negative selection process serves to reduce the rate of changes in the amino acid profile. In summary, the findings indicate that the majority of *VvNAC* genes exhibit a slow evolutionary rate, characterized by the absence of positive selection.

4. Results

Table 4.2. (A) The predicted number of transmembrane helices (TMHs) of *VvNAC* genes using TMHMM v2.0.

“Number of predicted TMHs”: It refers to the actual count or number of TMHs that have been predicted in the protein sequence; **“Expected number TMHs AAs”**: It refers to the expected number of TMHs predicted throughout the entire length of the protein considering all the amino acids (AAs) in the protein; **“Expected number first 60 AAs”**: It refers to the expected number of TMHs predicted only within the first 60 amino acids of the protein.

Gene ID	Length Base pair (bp)	Number of predicted TMHs	Transmembrane sequences (position)	Expected number, TMHs AAs	Expected number, first 60 AAs
<i>VvNAC15</i>	572	1	541-563	21.57467	0.00038
<i>VvNAC20</i>	559	1	530-552	21.15784	0
<i>VvNAC25</i>	632	1	600-622	22.00932	0
<i>VvNAC29</i>	214	1	190-212	24.22788	0.00015
<i>VvNAC48</i>	560	1	526-548	22.59021	0.00015
<i>VvNAC54</i>	994	1	537-559	34.49748	0.00212
<i>VvNAC67</i>	375	1	341-363	22.82849	0.00112
<i>VvNAC71</i>	578	1	551-573	20.45545	0
<i>VvNAC72</i>	659	6	449-471 484-506 526-548 569-591 606-623 636-658	127.28941	0.00023

Table 4.2. (B) The Ka/Ks values of *Vitis vinifera* tandem repeat sequences.

Tandem Repeat Sequences	Ka	Ks	Ka/Ks
<i>VvNAC52/VvNAC54</i>	0.156069171	0.226033598	0.690468905
<i>VvNAC50/VvNAC53</i>	0.428950207	0.616323419	0.695982327
<i>VvNAC71/VvNAC25</i>	0.488769856	1.700995236	0.28734346
<i>VvNAC21/VvNAC20</i>	0.428894814	1.341508352	0.319710879
<i>VvNAC64/VvNAC69</i>	0.355003859	2.934806313	0.120963301
<i>VvNAC34/VvNAC37</i>	0.1981519	2.090261815	0.094797646
<i>VvNAC58/VvNAC47</i>	0.219426571	2.011036466	0.109111185
<i>VvNAC01/VvNAC07</i>	0.202803106	1.131456332	0.179240771
<i>VvNAC39/VvNAC08</i>	0.192497524	2.430682324	0.079194851
<i>VvNAC66/VvNAC43</i>	0.281571219	0.652713909	0.43138535
<i>VvNAC03/VvNAC18</i>	0.247258086	1.59634065	0.154890553

4. Results

<i>VvNAC45/VvNAC42</i>	0.351230457	1.161810834	0.302312947
<i>VvNAC11/VvNAC05</i>	0.240144639	2.824820383	0.085012357
<i>VvNAC65/VvNAC16</i>	0.466070377	3.195651915	0.145845164
<i>VvNAC22/VvNAC02</i>	0.131794832	1.188950664	0.110849706
<i>VvNAC49/VvNAC24</i>	0.175166293	1.760241308	0.099512659
<i>VvNAC29/VvNAC28</i>	0.240883303	0.661545539	0.364122028
<i>VvNAC30/VvNAC31</i>	0.045084736	0.074318177	0.606644812
<i>VvNAC10/VvNAC27</i>	0.161614425	1.03734359	0.155796428
<i>VvNAC04/VvNAC41</i>	0.299027751	2.406023034	0.124282996

4.3. Conserved motifs of *VvNAC* genes

The *VvNAC* genes contain a highly conserved DNA binding NAC domain at the N-terminal, consisting of 160 amino acid residues that are divided into five subdomains: A, B, C, D, and E (Figure 4.3-A). However, the C-terminal part is highly variable and lacks any known protein domains.

To further investigate the conserved motifs of NAC proteins from *Vitis vinifera* L., the MEME 5.5.2 web database program was utilized, resulting in the identification of one to ten motifs. A thorough investigation of the conserved motifs present in each *VvNAC* gene was also performed.

The outcomes of the investigation pertaining to the distribution of NAC protein motifs have elucidated significant patterns. Specifically, within the subset of 74 grape NAC proteins, a substantial portion comprising 57 proteins (equivalent to 77% of the total) has been identified as encompassing subdomains A to E. Furthermore, a distinct profile has emerged where six proteins exhibit the absence of a singular NAC sub-domain, namely A, B, or C. In addition, an analogous group of six proteins is characterized by the lack of two sub-domains, manifesting either as B and C or A and C.

A remarkable observation pertains to the presence of two sub-domains, A and B, within a sole protein, *VvNAC66*. Moreover, a noteworthy subset consists of three proteins, specifically *VvNAC35*, *VvNAC38*, and *VvNAC50*, each harboring only a solitary sub-domain, namely the C sub-domain. Notably, another protein, *VvNAC72*, distinctly carries the A sub-domain.

Moreover, the range of motifs identified within *Vitis* NAC proteins extended from one to thirteen. Remarkably, six *VvNAC* proteins namely, *VvNAC28*, *VvNAC30*, *VvNAC31*, *VvNAC32*, *VvNAC36*, and *VvNAC60* hosted eight motifs each. In parallel, four proteins, specifically *VvNAC09*, *VvNAC35*, *VvNAC38*, and *VvNAC66*, showcased the presence of three motifs, while *VvNAC50* featured two motifs.

Notably, motifs 1 and 7 exhibited a consistent occurrence among the *VvNAC* proteins. The analysis unveiled six motifs prevalent in 52.7% of the 74 *VvNAC* proteins, whereas seven motifs were identified in 9.4%, and five motifs presented in 20.2% of the *VvNAC* proteins. It's noteworthy to mention that *VvNAC54* was notable for harboring the highest count of motifs, specifically thirteen, while conversely, *VvNAC72* exhibited the lowest, featuring just a solitary motif (Figure 4.3-B) (Table 4.3.).




Figure 4.3 (A) Illustration of the predicted subdomains of *VvNAC* genes using MEME 5.5.2. The *VvNAC* genes contain a highly conserved DNA binding NAC domain at the N-terminal, consisting of 160 amino acid residues that are divided into five subdomains: A, B, C, D, and E.

4. Results




Figure 4.3 (B) Motif analysis of grape *VvNAC* genes using MEME 5.5.2. The distribution of the motifs within the *VvNAC* genes reveals that six motifs are present in 53% of the genes, while seven motifs are found in 9.4% of the genes and five motifs are present in 20.2%. Additionally, *VvNAC54* has the highest number of motifs (13), while *VvNAC72* has the lowest (1).

Table 4.3. Distribution of the identified motifs' number and subdomains of *Vitis vinifera* *NAC* genes via MEME 5.5.2.

Gene ID	Number of motifs	Sub-domains
<i>VvNAC01</i>	5	ACDE
<i>VvNAC02</i>	6	ABCDE
<i>VvNAC03</i>	6	ABCDE
<i>VvNAC04</i>	5	ABCDE
<i>VvNAC05</i>	6	ABCDE
<i>VvNAC06</i>	6	ABCDE
<i>VvNAC07</i>	5	ACDE

4. Results

<i>VvNAC08</i>	5	ABCDE
<i>VvNAC09</i>	3	BDE
<i>VvNAC10</i>	7	ABCDE
<i>VvNAC11</i>	6	ABCDE
<i>VvNAC12</i>	5	ADE
<i>VvNAC13</i>	6	ABCDE
<i>VvNAC14</i>	6	ABCDE
<i>VvNAC15</i>	7	ABCDE
<i>VvNAC16</i>	6	ABCDE
<i>VvNAC17</i>	6	ABCDE
<i>VvNAC18</i>	6	ABCDE
<i>VvNAC19</i>	6	ADE
<i>VvNAC20</i>	7	ABCDE
<i>VvNAC21</i>	6	ABCDE
<i>VvNAC22</i>	6	ABCDE
<i>VvNAC23</i>	6	ABCDE
<i>VvNAC24</i>	6	ABCDE
<i>VvNAC25</i>	7	ABCDE
<i>VvNAC26</i>	6	ABCDE
<i>VvNAC27</i>	6	ABCDE
<i>VvNAC28</i>	8	ABCDE
<i>VvNAC29</i>	5	ABCDE
<i>VvNAC30</i>	8	ABCDE
<i>VvNAC31</i>	8	ABCDE
<i>VvNAC32</i>	8	ABCDE
<i>VvNAC33</i>	6	ABCDE
<i>VvNAC34</i>	6	ADE
<i>VvNAC35</i>	3	C
<i>VvNAC36</i>	8	ABCDE
<i>VvNAC37</i>	5	ADE
<i>VvNAC38</i>	3	C
<i>VvNAC39</i>	5	ABCDE

4. Results

<i>VvNAC40</i>	6	ABCDE
<i>VvNAC41</i>	6	ABCDE
<i>VvNAC42</i>	6	ABCDE
<i>VvNAC43</i>	5	ABCDE
<i>VvNAC44</i>	6	ABCDE
<i>VvNAC45</i>	6	ABCDE
<i>VvNAC46</i>	6	ABCDE
<i>VvNAC47</i>	5	ABDE
<i>VvNAC48</i>	5	ABCDE
<i>VvNAC49</i>	6	ABCDE
<i>VvNAC50</i>	2	C
<i>VvNAC51</i>	6	ABCDE
<i>VvNAC52</i>	7	ABCDE
<i>VvNAC53</i>	6	ABCDE
<i>VvNAC54</i>	13	ABCDE
<i>VvNAC55</i>	6	ABCDE
<i>VvNAC56</i>	6	ABCDE
<i>VvNAC57</i>	6	ABCDE
<i>VvNAC58</i>	5	ABDE
<i>VvNAC59</i>	5	ABCDE
<i>VvNAC60</i>	8	ABCDE
<i>VvNAC61</i>	6	ABCDE
<i>VvNAC62</i>	5	BCDE
<i>VvNAC63</i>	6	ABCDE
<i>VvNAC64</i>	6	ABCDE
<i>VvNAC65</i>	6	ABCDE
<i>VvNAC66</i>	3	AB
<i>VvNAC67</i>	7	ABCDE
<i>VvNAC68</i>	6	ABCDE
<i>VvNAC69</i>	6	ABCDE
<i>VvNAC70</i>	6	ABCDE
<i>VvNAC71</i>	7	ABCDE

<i>VvNAC72</i>	1	A
<i>VvNAC73</i>	5	ABDE
<i>VvNAC74</i>	6	ABCDE

4.4. Protein–protein interaction of *VvNAC*

The predicted protein-protein interaction map revealed intricate interactions among proteins themselves as well as with several other proteins. To identify optimal matches, NAC protein sequences from *Vitis vinifera* L. were compared to *Arabidopsis thaliana* proteins, and the corresponding proteins were selected for subsequent network analysis.

The *VvNACs* protein interaction network was established based on *Arabidopsis* protein orthologs, and those *VvNAC* proteins exhibiting high similarity to *Arabidopsis* proteins were referred to as STRING proteins (Figures 4.4 and Appendix – Table 7.1, 7.2).

Specifically, *VvNAC26* and *VvNAC43* displayed homology with NAP, *VvNAC67* with NTL8, *VvNAC51*, *VvNAC52*, *VvNAC53*, *VvNAC54*, *VvNAC55*, *VvNAC69*, and *VvNAC71* with NTL9, *VvNAC64* with NAC028, and *VvNAC60* with NAC047. Notably, all of these proteins exhibited significant interactions with the NTL1 protein transporter activity, highlighting their functional relevance.

Furthermore, within a distinct cluster, *VvNAC47* and *VvNAC58* showed homology with XND1, *VvNAC63* with VND7, *VvNAC24* and *VvNAC49* with NST1, *VvNAC01*, *VvNAC07*, and *VvNAC73* with NAC083, and *VvNAC34* and *VvNAC37* with NAC073. These proteins displayed robust associations with various regulatory factors, including MYB46 (involved in regulating secondary wall biosynthesis in fibres and vessels), MYB83 (a key player in the NAC012/SND1-mediated transcriptional network governing secondary wall biosynthesis), MYB80 (critical for the timing of tapetal programmed cell death, a pivotal process in pollen development), MYB85 (a putative transcription factor), AtMYB103 (a putative MYB family transcription factor), and KNAT7 (a KNOTTED-like homeobox protein in *Arabidopsis thaliana* 7, implicated in secondary cell wall biosynthesis).

Moreover, *VvNAC21* and *VvNAC50* exhibited homology with NAC050, a DNA-binding transcription factor involved in multicellular organismal development and transcription regulation. These proteins demonstrated strong associations with JM14, a transcriptional repressor involved in DRM2-mediated maintenance of DNA methylation, as well as with AT1G72460, a leucine-rich repeat protein kinase family protein participating in the transmembrane receptor protein tyrosine kinase signalling pathway and protein amino acid phosphorylation.

Finally, *VvNAC08* and *VvNAC39*, homologous to ATAF1, acted as pivotal genes connecting the two clusters and exhibited strong interactions with *VvNAC44* (homologous to NAC032), which positively regulates age-dependent senescence, dark-induced leaf senescence, and stress-induced senescence.

Consequently, based on STRING database, the Gene Ontology (GO) predictions revealed various gene functions within the network, including the nucleus, DNA binding, organic cyclic compound binding, regulation of cellular metabolic processes, regulation of macromolecule metabolic

processes, regulation of transcription (DNA-templated), regulation of nucleobase-containing compound metabolism, cell cycle, and DNA repair.




Figure 4.4 Protein-protein interaction network of *Arabidopsis* NAC homologs predicted with STRING 11.5.

4.5. Promoter regions detection and analysis of cis-acting regulatory elements

To identify cis-acting regulatory elements (CAREs) in *Vitis vinifera*'s NAC genes, the promoter sequences were analyzed using the Plant Care database. Specifically, the sequences were examined up to 1500 base pairs (bp) upstream of the translation start site. CAREs are small, conserved patterns of nucleotides that are crucial components of genes, commonly found at the 5' ends.

In total, 74 *Vitis vinifera* CAREs were analyzed to explore their response to diverse stress types. The investigation revealed that the *Vitis vinifera* NAC genes contain more than 3000 CAREs. The identified CAREs include crucial elements that respond to both biotic and abiotic stressors, involve in hormonal regulation, assist in the development, and perform various functions in the promoter regions of VvNAC genes (Appendix - Table 7.3).

4.5.1. Stress responsive cis-acting regulatory elements

To begin, abiotic stress-related motifs, such as MYC, MYB, and MBS-MYB binding sites, LTR (low-temperature elements), STRE (stress-responsive elements), and DRE (dehydration-responsive elements), were identified. Additionally, various CAREs were discovered in response to biotic stress, including STRE, wound-responsive (WUN motif and WRE3), elicitor-responsive (W box), and TC-rich repeats (defense and stress).

Furthermore, the study demonstrated that all *Vitis NAC* promoters contained motifs specific to light stresses, such as the Sp1, AE-box, Box 4, Chs-CMA1a, GA motif, GT1 motif, GATA-motif, I-box, Box S, G box, Gap box, and Box II.

4.5.2. *Cis-acting regulatory elements in hormonal regulation.*

The cis-regulatory elements of the *VvNACs* were found to be involved in hormonal regulation, including multiple ligand-responsive elements such as the TGA-box (auxin responsive elements), ERE (ethylene responsive elements), GARE motifs (gibberellin responsive elements), P-box, ABRE (abscisic acid-responsive elements), AuxRR-core, TCA elements (salicylic acid-responsive elements), TATC-box, and TGA-element. These elements enhance the plants' stress resistance capabilities.

4.5.3. *Role of cis-acting regulatory elements in cellular development:*

Finally, while the number of motifs involved in cellular development is relatively fewer compared to other CAREs, the presence of several elements has been demonstrated. These include the CAT box (for meristem-specific activation), Motif I, GCN4 motif (for endosperm expression), AC-I and AC-II (for xylem-specific expression), O2-site (for zein metabolism regulation), F-box (for reproductive and vegetative development and growth, as well as cell defense and death), ARE (for anaerobic induction), the circadian motif, HD-Zip1 (for leaf development), AP-1 (for inducible flowering), and MBSI (for flavonoid biosynthesis gene regulation).

Given that most *VvNACs* exhibit three distinct types of stress-responsive CAREs, it is plausible to assume that these genes play a crucial role in enhancing tolerance to various biotic and abiotic stresses. Overall, these findings suggest that *Vitis vinifera's NAC* genes contain a diverse range of CAREs that respond to various stressors and are involved in hormonal regulation and cellular development.

4.5.4. *In silico analysis of VvNAC gene expression*

Biotic expression

The transcription profiling by high throughput sequencing of grapevine *NAC* genes at 3 different stages (early S1, middle S2, and late S3) of noble rot caused by *Botrytis cinerea* was extracted from the Expression Atlas database, as referenced in (Appendix – [Table 7.4](#)).

The analysis of the microarray data set showed that several *VvNAC* genes, including *VvNAC39*, *VvNAC44* and *VvNAC73* were highly regulated during stages S2 and S3. Additionally, a slight up-regulation was detected in *VvNAC08*, *VvNAC30*, *VvNAC31*, *VvNAC36*, *VvNAC41*, *VvNAC60*, *VvNAC61* and *VvNAC74*. Conversely, there was a significant down-regulation in *VvNAC13*, *VvNAC51*, *VvNAC52*, *VvNAC53* and *VvNAC54* ([Figure 4.5](#)).




Figure 4.5 Heat map of the microarray data representing the expression analysis of *VvNAC* genes in response to *Botrytis cinerea*. Based on Expression Atlases at three different stages of Noble Rot infection in ripe grapes; early (S1), middle (S2) and late (S3). The expression patterns of *VvNAC* genes were analyzed using the offline program TBtools.

4.5.5. Subcellular localization prediction

The subcellular localization of NAC proteins in *Vitis vinifera* L. was analyzed using the CELLO v.2.5 web database tool. Results revealed that most proteins (73%) were localized in the nucleus, while eleven proteins, namely *VvNAC08*, *VvNAC15*, *VvNAC20*, *VvNAC26*, *VvNAC30*, *VvNAC31*, *VvNAC32*, *VvNAC36*, *VvNAC37*, *VvNAC38*, and *VvNAC64*, were present in both the cytoplasm and nucleus.

Interestingly, *VvNAC72* was highly predicted to be located in the plastid. Moreover, *VvNAC39* and *VvNAC40* were linked to mitochondria, nuclei, and the cytoplasm, while *VvNAC05* was found in the nucleus and mitochondria. In addition, *VvNAC50* was located in the mitochondria and extracellular matrix, while *VvNAC73* was distributed in the cytoplasm and mitochondria. Furthermore, *VvNAC35* and *VvNAC51* were found in the cytoplasm, and *VvNAC52* was located in the cytoplasm and extracellular matrix, as illustrated in [Figure 4.6](#).

4. Results




Figure 4.6 Heat map of the subcellular-localization prediction of *VvNAC* genes through web database tool CELLO Life v.2.5 in different organelles. The results were analyzed using the offline program TBtools: Nucl (nucleus), cyto (cytoplasmic), extrac (extracellular matrix), mito (mitochondrial), chlor (chloroplast), plas (plastid), ER (endoplasmic reticulum), vac (vacuolar), lyso (lysosome), cytos (cytosol), pero (peroxisome), gol (Golgi apparatus).

4.5.6. Statistical Analysis

Through hierarchical clustering a statistical analysis for gene expression data was generated, by creating a tree-like structure, known as a dendrogram, the relationships between genes were visually represented.

The dendrogram formed 4 clusters of the *VvNAC* genes with similar expression profiles were grouped together in branches. Cluster 1 and cluster 2 contain low expressed genes while cluster 3 and 4 contain the high expressed genes as shown in [Figure 4.7](#).




Figure 4.7 The dendrogram of the *VvNAC* genes based on their expression profiles.

The Euclidean distance were used to measure the distance between the clusters to determine their similarities in a quantitative manner based on the Euclidean formula:

$$d_{uv} = \sqrt{(u_1 - v_1)^2 + (u_2 - v_2)^2 + (u_3 - v_3)^2}$$

As a result, showed in Appendix- [Table 7.5](#), the distance between the clusters 2 & 3 based on the single linkage was 4.2; the distance between the closest pair *VvNAC64*&*VvNAC52*.

The distance between the clusters 1 & 4 based on the single linkage was: 6; the distance between *VvNAC03*&*VvNAC73*.

The distance between the clusters 2 & 4 based on the single linkage was: 10.5; the distance between *VvNAC52*&*VvNAC73*.

While the distance between the clusters 3 & 4 based on the single linkage was: 2.2; the distance between *VvNAC30*&*VvNAC73*.

Therefore, the proximity of clusters 3 and 4 in terms of gene expression suggests that they may share similar regulatory mechanisms or pathways in response to *Botrytis cinerea*. Genes in these clusters may be involved in common biological processes. While the larger distance between clusters 1&4, 2&3 and 2&4 indicates dissimilarity and the genes in these clusters have distinct

expression patterns. This could be due to a different regulatory mechanism or involvement in different biological processes.

4.5.7. *In vitro* verifications: *VvNAC36* model

In *Vitis vinifera* L. a recent study has proved a remarkable regulation of the *VvNAC36* gene in *A. thaliana* in response to powdery mildew colonization. The up-regulation of *VvNAC36* 1.5-fold in the case of powdery mildew (PM) while no response was detected for salicylic acid (SA), has demonstrated the role of *VvNAC36* in response to biotic stress (Tóth et al., 2016).

Transient expression

Continuing Tóth's work, a simple and rapid transient transformation in tobacco leaf was performed to determine the localization of *VvNAC36* protein that might help in better understanding its role in plants.

In plants, yeast, and mammals, the GFP tag is widely used to mark proteins for localization studies. Confocal microscope was used to image GFP fusion proteins that were transiently expressed in leaves. The use of a constitutive cauliflower mosaic virus 35S promoter (CaMV 35S) enabled GFP to be seen in leaf tissues that normally have a lot of autofluorescence. The fusion proteins *VvNAC36::sGFP* and the control *35S::sGFP* were agroinfiltrated into leaves of 3 to 5-week-old *N. benthamiana* plants.

Our investigation of the subcellular localization of the *VvNAC36::sGFP* fusion protein in tobacco epidermal cells indicated that the protein was mainly present in the cytoplasmic area (Figure 4.8; A, B and C).

Furthermore, our analysis suggested that the *VvNAC36* protein might also be present in various organelles, as GFP was detected in different shapes (Figure 4.8; D, E and F). However, this observation could not be confirmed conclusively due to the limited availability of advanced biotechnology methods.

Finally, for the positive control *35S::sGFP*, the detection of the green fluorescent protein showed an accumulation in both the cytoplasm and nucleus of the tobacco leaf epidermal cells (Figure 4.8; G, H and I).




Figure 4.8 Subcellular localization of *VvNAC36* and the control in transiently transformed leaves of *Nicotiana benthamiana* visualized by using confocal microscope, Bars = 20.15 µm. The GFP fluorescence pattern of the fusion protein *VvNAC36::sGFP* was detected in the cytoplasmic area as demonstrated by white arrows in images A, B, and C. Images D, E, and F further depict the visualization of *VvNAC36::sGFP* expression in other regions as indicated by green arrows. For the positive control *35S::sGFP* (GFP alone), the GFP fluorescence was observed in the cytoplasm and nucleus, as indicated by blue arrows in images G, H, and I.

5. DISCUSSION AND LIMITATIONS

Botrytis cinerea is a dangerous plant pathogenic fungus with wide host ranges. This aggressive pathogen uses multiple weapons to invade and cause serious damages to the plants, such as grape (*Vitis vinifera* L.); a member of the *Vitaceae* family.

Being one of the largest among plant-specific TFs, the NAC protein family has a role in plant development, abiotic stress and defense responses. In many plant species NAC proteins have been functionally characterized, including e.g. *Vitis vinifera* L., *Arabidopsis thaliana*, *Oryza sativa*, *Zea mays*, *Triticum aestivum*, *Glycine max*, *Populus trichocarpa* (Fang et al., 2008; Pinheiro et al., 2009). However, the functions for majority of the NAC genes in grape remain unknown.

The current study employed an *in silico* approach to conduct a comprehensive genome wide detection of NAC domain transcription factors (TFs) within the *Vitis vinifera* L. genome, aiming to identify and characterize potential NAC proteins that exhibit resistance to *B. cinerea*. Furthermore, to validate the computational findings, an *in vitro* approach was implemented specifically for the *VvNAC36* gene through a transient expression assay.

In this study, a total of 74 NAC genes were analyzed from *Vitis vinifera* L.. The identified NAC genes varied greatly in protein length from 128 AA (*VvNAC50*) to 994 AA (*VvNAC54*).

Regarding subcellular localization prediction, it was discovered that 54 out of the 74 identified NAC genes in *Vitis* (accounting for 73% of the total) were potentially localized within the nucleus. In contrast, the remaining NAC genes displayed extracellular localization or were distributed across various organelles. Usually, transcription factors need to be localized to nucleus to execute their function, either independently or by interacting with other partners.

For instance, *ATAF1* is localized to nucleus (Lu et al., 2007), whereas *ONAC020* and *ONAC023* completely gets localize to nucleus after interacting with *ONAC26* (Mathew et al., 2016). Similarly, *NTL4* is targeted to the nucleus only upon heat stress after processing (Lee et al., 2014). There are numerous reports which shows the localization of NAC genes in different organelles other than nucleus, as in the case of *ONAC023*, which is localized to the cytoplasm (Mathew et al., 2016). Sometimes, TFs gets localized to nucleus after splicing of membrane bound TFs or upon proteolytic cleavage (Ng et al., 2013; Seo, 2014).

Furthermore, analyzing nucleotide substitution rates (Ka and Ks) provides valuable insights into evolutionary processes, genetic diversity, disease mechanisms, and the functional aspects of genomes. The analysis conducted in this study implies the absence of positive selection in any of the tandem repeat sequences, which are typically associated with facilitating adaptive genetic variation and potentially influencing species evolution.

However, all genes displayed Ka/Ks values below 1, indicating the prevalence of negative selection during the course of evolution. This negative selection process serves to reduce the rate of changes in the amino acid profile. In summary, the findings indicate that the majority of *VvNAC* genes exhibit a slow evolutionary rate, characterized by the absence of positive selection.

Thus, this might suggest that these *VvNAC* genes are performing important functions that haven't required rapid changes to their genetic makeup due to advantageous variations. This information can be valuable for understanding the genetic stability and adaptability of the plant species in question.

Additionally, the presence of diverse conserved motifs implies functional divergence among these VvNAC proteins. This divergence underscores their involvement in various critical processes, including meristem development, root development, flowering induction, embryo development, vascular-specific expression, hormone signaling, abiotic stress responses, and defense mechanisms.

The exploration of critical cis-acting regulatory elements (CAREs) highlighted their significance in governing responses to both abiotic and biotic stresses. Among these CAREs, notable examples include 228 Abscisic Acid Response Elements (ABRE), 145 stress-responsive elements (STRE), 17 Dehydration-Responsive Elements (DRE-core), 58 elicitor-responsive motifs (W-box), 78 wound-responsive elements (WRE3 and WUN motifs), and 21 TC-rich repeats, as shown in [Figure 5.9](#).

Besides these, several other promoter elements were also identified which have a role in various plant development processes. CAREs are necessary for stress-responsive transcriptional regulation (Mundy et al., 1990). The existence of different cis-acting regulatory elements indicates the transcription of several stress responsive genes via a variety of TFs. Moreover, the significance of the association between CAREs has already been documented for stress-responsive transcription (Narusaka et al., 2003).

In several reports, various cis-motifs as DNA-binding sites for the NAC TFs have been identified, which include NACRS (NAC-recognition sequence for drought response) (Tran et al., 2004), IDE2 motif (iron deficiency-responsive) (Ogo et al., 2008), SNBE (secondary wall NAC binding element) (Zhong et al., 2007), and calmodulin-binding (CBNAC) (Kim et al., 2007).

Moreover, the VvNACs protein-protein interaction network based on their homologous proteins from *Arabidopsis* provides a comprehensive and efficient way to explore molecular interactions, functional annotations, and potential roles of genes within biological systems.

Consequently, as indicated in [Figure 4.4](#), this study's results illuminated robust homology and strong connections between VvNAC proteins and established *Arabidopsis* counterparts, including *NAC083*, *ATAF1*, *NAC073*, *NAC050*, *NAC028*, *NAC047*, *NAP*, *NTL9*, *NTL8*, *MYB83*, *MYB80*, *XND1*, *VND7*, *NST1*, and *NTL1*.

This observation hints at analogous functionalities of VvNAC proteins. Particularly noteworthy was the remarkable homology shared among VvNAC08, VvNAC39, and *ATAF1*. This is exemplified by prior studies indicating that *ATAF1* plays a pivotal role as a transcriptional activator in response to salt stress, abscisic acid (ABA), and biotic stress (Wu et al., 2009; Liu et al., 2016).

Next, the expression profiles of VvNAC genes unveiled up-regulation in response to *B. cinerea* infection for specific genes, including VvNAC08, VvNAC30, VvNAC31, VvNAC36, VvNAC39, VvNAC41, VvNAC44, VvNAC60, VvNAC61, VvNAC73 and VvNAC74.

These genes were found to be enriched with biotic stress-related motifs, thus highlighting their potential involvement in biotic stress responses ([Figure 5.10](#)). As the photo showed, VvNAC39 and VvNAC44 with TC-rich repeats, STRE and W box; VvNAC73 with W box and STRE; VvNAC08 and VvNAC30 with WRE3, STRE and W box; VvNAC60 with STRE and WUN-motif; VvNAC31 with TC-rich repeats and W box; VvNAC41 with STRE, W box and WUN-motif; finally VvNAC61 and VvNAC74 with STRE and WRE3.

Thus, different novel *VvNAC* genes that respond to *B. cinerea* infection were detected using a combination of cis-element and microarray data analysis. As a result, these genes are excellent resources for *B. cinerea* stress in grapes in genetic engineering and molecular breeding.

It is important to mention here that the present study provides very detailed analysis of the identified *Vitis vinifera* NAC genes and their cis-regulatory elements as compared to the previous reports by (Wang et al. 2013; Le Henaff et al. 2013), respectively. We have carried out strict promoter motif analysis, protein-protein interactions, evolutionary rate analysis, and subcellular localization of the stress-related NAC proteins which lacked in previous reports and therefore, provides much deeper understanding of mechanisms involved in stresses and especially in biotic stress tolerance in *Vitis vinifera* L.




Figure 5.9 Various cis-acting regulatory elements related to stress, hormone and development process in *VvNAC* genes were identified using Plant Care database. The results were analyzed using the offline program TBtools. Among the identified CAREs: 228 Abscisic acid Response Elements (ABREs), 145 stress-responsive elements (STREs), 17 dehydration-responsive elements (DRE-core), 58 elicitor-responsive elements (W box), 78 wound-responsive elements (WRE3 and WUN motifs), and 21 TC-rich repeats. ABRE, DRE, MYB, MYC, TC-rich repeats, STRE, W Box, LTR, MBS, WUN-motif and WRE3 are related to stress; CAT-Box, HD-Zip1, O2 site, F-Box, RY-element, GCN4-motif, Circadian, CCGTCC-motif, Box 4, ARE MRE, G-Box and Mbs1 are related to development; AUXRR-core, CGTXA-motif, GARE-motif, P-Box, TCA- element, TGA-element, TATC-Box, ERE and SARE are related to hormone.

6. Conclusions and Recommendations




Figure 5.10 The predicted stress responsive *VvNAC* genes with their CAREs related to biotic stress. The results were illustrated using the offline program TBtools. Each gene contains at least two or more CAREs that respond to biotic stress.

5.1. *VvNAC36* protein

Based on CELLO Life, the localization of *VvNAC36* protein is more likely to be in the nucleus and cytoplasm with a percentage of 54.4% and 21.1% respectively.

In order to explain the *in silico* result, a transient expression has been done for *VvNAC36* gene. Thus, as a result in the case of the *VvNAC36* protein, the GFP fluorescence presented in the cytoplasmic area, as shown in the Figure 4.8., while, a localization in the nucleus and the cytoplasmic area showed in the case of the control.

Consequently, comparing to the size of the nucleus of the control, we confirm the absence of any nuclear localization for the gene *VvNAC36* and just the cytoplasmic one was approved. Despite the good results, but future studies should use more advanced techniques to deeply investigate and confirm the specific localization of this gene.

The expression analysis based on microarray data available has showed a slight up-regulation just in the early stage of infection of the ripe grape (S1), this might be explained by the presence of different CAREs responsible for biotic stress in the promoter of *VvNAC36* such as; STRE and W box as showed in the photo Figure 5.10.

The computational results showed a weak resistance to *B. cinerea* that occurred just in S1 and did not present in the middle or late stages of the infection. Thus, a stable transformation must be done in the future in order to check how *VvNAC36* might respond to *B. cinerea* in different stages.

5.2. Study limitations

The COVID-19 pandemic in 2020 posed significant challenges, hindering my ability to carry out experiments like stable transformation due to lockdown restrictions. In response, I turned to an *in silico* research approach for my theory-focused Ph.D. dissertation. This choice was driven by its efficiency, flexibility, and its potential to offer fresh theoretical insights into complex biological

processes. Importantly, it also aligns with ethical research practices and proves cost-effective, making it a practical and responsible solution to the limitations imposed by the pandemic.

6. CONCLUSIONS AND RECOMMENDATIONS

- This study aimed to highlight and interpret the functions of *VvNAC* genes in *Vitis vinifera* L., beyond their known roles in development and stress responses, using both *in vitro* and *in silico* approaches.
- Based on the grapevine genomic databases, we have characterized 74 *VvNAC* genes.
- As a result, the analyzing of the protein's lengths and number of motifs is just one aspect of a gene's potential to respond to biotic stress and it should not be the sole determinant of a gene's function.
- For example: the longest gene *VvNAC54* with a higher number of motifs (13) showed down-regulation in response to *B. cinerea*, while one of the shortest genes, the *VvNAC73* with 5 motifs showed high expression to the disease.
- While it's logical to assume that longer length and more motifs might make a gene more effective in responding to stress, several factors can influence the outcomes, as we have observed in the case of *VvNAC54* and *VvNAC73*.
- Continuing, with the Ka/Ks value, all genes displayed Ka/Ks values below 1. This suggests that the genes in this tandem repeat sequence are likely under strong purifying selection, and they are likely performing important functions without substantial genetic modifications.
- Moreover, in the subcellular localization analysis, it is noteworthy to mention *VvNAC72*. A gene with a specialty to be in the plastid, makes us curious and eager to investigate further.
- In addition, protein-protein interactions analysis is important because understanding these relationships, such in the case of *VvNAC08*, *VvNAC39* and *VvNAC44* can be valuable in biotechnological applications, such as crop improvement. It may lead to the development of strategies to enhance stress resistance or other desirable traits in grapevines based on the knowledge from *Arabidopsis thaliana*.
- The CAREs in the up-regulated genes promoters might not be unique just to *Vitis vinifera* *NAC genes* but they represent conserved motifs with potential involvement in biotic stress response.
- Based on the expression profiles analysis, we predicted 10 *VvNAC* genes for *B. cinerea* tolerance. Followed by a statistical analysis through hierarchical clustering for the gene's expression.
- Additional experimental validation, such as quantitative real-time PCR (qRT-PCR) analysis, is required to verify the expression pattern of the predicted *VvNACs* in response to *B. cinerea*.

- In the case of the *in vitro* experiment related to *VvNAC36*, the results of the transient expression encountered with the computational tools' results.
- The stable transformation must be done, and *in vitro* experiments should be performed to the predicted *VvNAC* genes and not just for *VvNAC36*.
- Finally, this study has increased our knowledge of grape *NAC* genes and provided insight into their functions in case of the up-regulated genes.
- Furthermore, our findings have built a robust framework for researchers to select candidates to engineer grape cultivars for enhanced tolerance against biotic stress.

6.1. New scientific results

1. An *in silico* approach was applied for detailed functional characterization of *VvNAC* transcription factor genes in response to *B. cinerea*.
2. We predicted 10 *VvNAC* genes (*VvNAC08*, *VvNAC30*, *VvNAC31*, *VvNAC36*, *VvNAC39*, *VvNAC41*, *VvNAC44*, *VvNAC61*, *VvNAC73*, and *VvNAC74*) for *B. cinerea* tolerance, that could be useful as reference for researchers in future grape breeding programs.
3. We have determined through a transient expression the subcellular localization of the *VvNAC36*, in the cytoplasmic area.
4. The feasibility study of *Nicotiana bentamiana* L. in transient transformation with the *VvNAC36* and the correlation between *in silico* and *in vitro* approaches were portrayed.

7. APPENDIXES

A1: Bibliography

1. Araus, J. L., Kefauver, S. C., Vergara-Díaz, O., Gracia-Romero, A., Rezzouk, F. Z., Segarra, J., Buchailot, M. L., Chang-Espino, M., Vatter, T., Sanchez-Bragado, R., Fernandez-Gallego, J. A., Serret, M. D., & Bort, J. (2022). Crop phenotyping in a context of global change: What to measure and how to do it. *Journal of integrative plant biology*, 64(2), 592–618. <https://doi.org/10.1111/jipb.13191>.
2. Atkin, O. K., & Macherel, D. (2009). The crucial role of plant mitochondria in orchestrating drought tolerance. *Annals of botany*, 103(4), 581–597. <https://doi.org/10.1093/aob/mcn094>.
3. Bailey, T. L., Boden, M., Buske, F. A., Frith, M., Grant, C. E., Clementi, L., Ren, J., Li, W. W., & Noble, W. S. (2009). MEME SUITE: tools for motif discovery and searching. *Nucleic acids research*, 37(Web Server issue), W202–W208. <https://doi.org/10.1093/nar/gkp335>.
4. Balazadeh, S., Siddiqui, H., Allu, A. D., Matallana-Ramirez, L. P., Caldana, C., Mehrnia, M., Zanol, M. I., Köhler, B., & Mueller-Roeber, B. (2010). A gene regulatory network controlled by the NAC transcription factor ANAC092/AtNAC2/ORE1 during salt-promoted senescence. *The Plant journal: for cell and molecular biology*, 62(2), 250–264. <https://doi.org/10.1111/j.1365-313X.2010.04151.x>.
5. Bettoni, J. C., Marković, Z., Bi, W., Volk, G. M., Matsumoto, T., & Wang, Q. C. (2021). Grapevine Shoot Tip Cryopreservation and Cryotherapy: Secure Storage of Disease-Free Plants. *Plants (Basel, Switzerland)*, 10(10), 2190. <https://doi.org/10.3390/plants10102190>.
6. Bigeard, J., Colcombet, J., & Hirt, H. (2015). Signaling mechanisms in pattern-triggered immunity (PTI). *Molecular plant*, 8(4), 521–539. <https://doi.org/10.1016/j.molp.2014.12.022>.
7. Bilas, R., Szafran, K., Hnatuszko-Konka, K., Kononowicz, A. K. (2016). Cis-regulatory elements used to control gene expression in plants. *Plant Cell Tissue Organ Cult. (PCTOC)* 127 (2), 269–287. doi: [10.1007/s11240-016-1057-7](https://doi.org/10.1007/s11240-016-1057-7).
8. Blanco-Ulate, B., Amrine, K. C. H., Collins, T. S., Rivero, R. M., Vicente, A. R., Morales-Cruz, A., Doyle, C. L., Ye, Z., Allen, G., Heymann, H., Ebeler, S. E., & Cantu, D., 2015. Transcription profiling by high throughput sequencing of berries at distinct stages of noble rot caused by *Botrytis cinerea*. Expression Atlas, microarray data sets, E-GEOD-67932. <https://www.ebi.ac.uk/gxa/experiments/E-GEOD-67932/Results>.
9. Bourne P. (2005). Will a biological database be different from a biological journal? *PLoS computational biology*, 1(3), 179–181. <https://doi.org/10.1371/journal.pcbi.0010034>.
10. Breeze, E., Harrison, E., McHattie, S., Hughes, L., Hickman, R., Hill, C., Kiddle, S., Kim, Y. S., Penfold, C. A., Jenkins, D., Zhang, C., Morris, K., Jenner, C., Jackson, S.,

- Thomas, B., Tabrett, A., Legaie, R., Moore, J. D., Wild, D. L., Ott, S., ... Buchanan-Wollaston, V. (2011). High-resolution temporal profiling of transcripts during Arabidopsis leaf senescence reveals a distinct chronology of processes and regulation. *The Plant cell*, 23(3), 873–894. <https://doi.org/10.1105/tpc.111.083345>.
11. Breia, R., Conde, A., Badim, H., Fortes, A. M., Gerós, H., & Granell, A. (2021). Plant SWEETs: from sugar transport to plant-pathogen interaction and more unexpected physiological roles. *Plant physiology*, 186(2), 836–852. <https://doi.org/10.1093/plphys/kiab127>.
 12. Bult, C. J., Blake, J. A., Smith, C. L., Kadin, J. A., Richardson, J. E., & Mouse Genome Database Group (2019). Mouse Genome Database (MGD) 2019. *Nucleic acids research*, 47(D1), D801–D806. <https://doi.org/10.1093/nar/gky1056>.
 13. Cai, F. F., Zhou, W. J., Wu, R., & Su, S. B. (2018). Systems biology approaches in the study of Chinese herbal formulae. *Chinese medicine*, 13, 65. <https://doi.org/10.1186/s13020-018-0221-x>.
 14. Cantu, D., Yang, B., Ruan, R., Li, K., Menzo, V., Fu, D., Chern, M., Ronald, P. C., Dubcovsky, J. (2013). Comparative analysis of protein-protein interactions in the defense response of rice and wheat. *BMC genomics*, 14, 166. <https://doi.org/10.1186/1471-2164-14-166>.
 15. Chen, C., Chen, H., Zhang, Y., Thomas, H. R., Frank, M. H., He, Y., & Xia, R. (2020). TBtools: An Integrative Toolkit Developed for Interactive Analyses of Big Biological Data. *Molecular plant*, 13(8), 1194–1202. <https://doi.org/10.1016/j.molp.2020.06.009>.
 16. Chen, Q., Wang, Q., Xiong, L., & Lou, Z. (2011). A structural view of the conserved domain of rice stress-responsive NAC1. *Protein & cell*, 2(1), 55–63. <https://doi.org/10.1007/s13238-011-1010-9>.
 17. Christianson, J. A., Dennis, E. S., Llewellyn, D. J., & Wilson, I. W. (2010). ATAF NAC transcription factors: regulators of plant stress signaling. *Plant signaling & behavior*, 5(4), 428–432. <https://doi.org/10.4161/psb.5.4.10847>.
 18. Collinge, M., & Boller, T. (2001). Differential induction of two potato genes, Stprx2 and StNAC, in response to infection by *Phytophthora infestans* and to wounding. *Plant molecular biology*, 46(5), 521–529. <https://doi.org/10.1023/a:1010639225091>.
 19. Cramer, G. R., Ergül, A., Grimplet, J., Tillett, R. L., Tattersall, E. A., Bohlman, M. C., Vincent, D., Sonderegger, J., Evans, J., Osborne, C., Quilici, D., Schlauch, K. A., Schooley, D. A., & Cushman, J. C. (2007). Water and salinity stress in grapevines: early and late changes in transcript and metabolite profiles. *Functional & integrative genomics*, 7(2), 111–134. <https://doi.org/10.1007/s10142-006-0039-y>.
 20. de Oliveira, G. L., de Souza, A. P., de Oliveira, F. A., Zucchi, M. I., de Souza, L. M., & Moura, M. F. (2020). Genetic structure and molecular diversity of Brazilian grapevine germplasm: Management and use in breeding programs. *PloS one*, 15(10), e0240665. <https://doi.org/10.1371/journal.pone.0240665>.

21. Denby, K. J., Kumar, P., & Kliebenstein, D. J. (2004). Identification of *Botrytis cinerea* susceptibility loci in *Arabidopsis thaliana*. *The Plant journal : for cell and molecular biology*, 38(3), 473–486. <https://doi.org/10.1111/j.0960-7412.2004.02059.x>.
22. Duval, M., Hsieh, T. F., Kim, S. Y., & Thomas, T. L. (2002). Molecular characterization of AtNAM: a member of the *Arabidopsis* NAC domain superfamily. *Plant molecular biology*, 50(2), 237–248. <https://doi.org/10.1023/a:1016028530943>.
23. Edwards, D., & Batley, J. (2010). Plant genome sequencing: applications for crop improvement. *Plant biotechnology journal*, 8(1), 2–9. <https://doi.org/10.1111/j.1467-7652.2009.00459.x>.
24. Elad Y., Williamson B., Tudzynski P., Delen N. (2004). “*Botrytis* spp. and diseases, they cause in agricultural systems-an introduction,” in *Botrytis: biology, pathology and control*. Eds. Elad Y., Williamson B., Tudzynski P., Delen N. (Dordrecht, the Netherlands: Kluwer;), 1–27. <https://link.springer.com/book/10.1007%2F978-1-4020-2626-3>.
25. Emanuelli, F., Lorenzi, S., Grzeskowiak, L., Catalano, V., Stefanini, M., Troglio, M., Myles, S., Martinez-Zapater, J. M., Zyprian, E., Moreira, F. M., & Grando, M. S. (2013). Genetic diversity and population structure assessed by SSR and SNP markers in a large germplasm collection of grape. *BMC plant biology*, 13, 39. <https://doi.org/10.1186/1471-2229-13-39>.
26. Fan, W., & Dong, X. (2002). In vivo interaction between NPR1 and transcription factor TGA2 leads to salicylic acid-mediated gene activation in *Arabidopsis*. *The Plant cell*, 14(6), 1377–1389. <https://doi.org/10.1105/tpc.001628>.
27. Fang Y, You J, Xie K, Xie W, Xiong L. (2008). Systematic sequence analysis and identification of tissue-specific or stress-responsive genes of NAC transcription factor family in rice. *Mol Gen Genomics*. 2008;280(6):547–63. <https://doi.org/10.1007/s00438-008-0386-6>.
28. Fang, L., Su, L., Sun, X., Li, X., Sun, M., Karungo, S. K., Fang, S., Chu, J., Li, S., & Xin, H. (2016). Expression of *Vitis amurensis* NAC26 in *Arabidopsis* enhances drought tolerance by modulating jasmonic acid synthesis. *Journal of experimental botany*, 67(9), 2829–2845. <https://doi.org/10.1093/jxb/erw122>.
29. Feechan, A., Anderson, C., Torregrosa, L., Jermakow, A., Mestre, P., Wiedemann-Merdinoglu, S., Merdinoglu, D., Walker, A. R., Cadle-Davidson, L., Reisch, B., Aubourg, S., Bentahar, N., Shrestha, B., Bouquet, A., Adam-Blondon, A. F., Thomas, M. R., & Dry, I. B. (2013). Genetic dissection of a TIR-NB-LRR locus from the wild North American grapevine species *Muscadinia rotundifolia* identifies paralogous genes conferring resistance to major fungal and oomycete pathogens in cultivated grapevine. *The Plant journal : for cell and molecular biology*, 76(4), 661–674. <https://doi.org/10.1111/tpj.12327>.
30. Finn, R. D., Coggill, P., Eberhardt, R. Y., Eddy, S. R., Mistry, J., Mitchell, A. L., Potter, S. C., Punta, M., Qureshi, M., Sangrador-Vegas, A., Salazar, G. A., Tate, J., & Bateman, A. (2016). The Pfam protein families database: towards a more sustainable future. *Nucleic acids research*, 44(D1), D279–D285. <https://doi.org/10.1093/nar/gkv1344>.

31. Fujita, M., Fujita, Y., Maruyama, K., Seki, M., Hiratsu, K., Ohme-Takagi, M., Tran, L. S., Yamaguchi-Shinozaki, K., & Shinozaki, K. (2004). A dehydration-induced NAC protein, RD26, is involved in a novel ABA-dependent stress-signaling pathway. *The Plant journal: for cell and molecular biology*, 39(6), 863–876. <https://doi.org/10.1111/j.1365-313X.2004.02171.x>.
32. Gao, J.-P., Chao, D.-Y. and Lin, H.-X. (2007). Understanding Abiotic Stress Tolerance Mechanisms: Recent Studies on Stress Response in Rice. *Journal of Integrative Plant Biology*, 49: 742-750. <https://doi.org/10.1111/j.1744-7909.2007.00495.x>.
33. Garretón, V., Carpinelli, J., Jordana, X., Holuigue, L. (2002). The as-1 promoter element is an oxidative stress-responsive element and salicylic acid activates it via oxidative species. *Plant physiology*, 130(3), 1516–1526. <https://doi.org/10.1104/pp.009886>.
34. Gonzalez-Meler, M. A., Rucks, J. S., & Aubanell, G. (2014). Mechanistic insights on the responses of plant and ecosystem gas exchange to global environmental change: lessons from Biosphere 2. *Plant science : an international journal of experimental plant biology*, 226, 14–21. <https://doi.org/10.1016/j.plantsci.2014.05.002>.
35. Goodstein, D. M., Shu, S., Howson, R., Neupane, R., Hayes, R. D., Fazo, J., Mitros, T., Dirks, W., Hellsten, U., Putnam, N., & Rokhsar, D. S. (2012). Phytozome: a comparative platform for green plant genomics. *Nucleic acids research*, 40(Database issue), D1178–D1186. <https://doi.org/10.1093/nar/gkr944>.
36. Greve, K., La Cour, T., Jensen, M. K., Poulsen, F. M., & Skriver, K. (2003). Interactions between plant RING-H2 and plant-specific NAC (NAM/ATAF1/2/CUC2) proteins: RING-H2 molecular specificity and cellular localization. *The Biochemical journal*, 371(Pt 1), 97–108. <https://doi.org/10.1042/BJ20021123>.
37. Gubler WD, Hashim JM, Smilanick JL, Leavitt GM (2013) Gray mold (*Botrytis cinerea*). In LJ Bettiga, ed, *Grape Pest Management*, Ed 3. University of California, Agriculture and Natural Resources, Richmond, CA, p 133–135.
38. Gujjar RS, Akhtar M and Singh M. (2014). Transcription factors in abiotic stress tolerance. *Indian J. Plant Physiol.* 19:306-316. [10.1007/s40502-014-0121-8](https://doi.org/10.1007/s40502-014-0121-8).
39. Gusain, P, Uniyal, DP, Joga, R. (2021). Conservation and sustainable use of medicinal plants. In: Egbuna, C, Mishra, AP, Goyal, MR, editors. *Preparation of phytopharmaceuticals for the management of disorders*. London: Academic Press; 2021. p. 409–27. <https://doi.org/10.1016/B978-0-12-820284-5.00026-5>.
40. Haile, Z. M., Pilati, S., Sonogo, P., Malacarne, G., Vrhovsek, U., Engelen, K., Tudzynski, P., Zottini, M., Baraldi, E., & Moser, C. (2017). Molecular analysis of the early interaction between the grapevine flower and *Botrytis cinerea* reveals that prompt activation of specific host pathways leads to fungus quiescence. *Plant, cell & environment*, 40(8), 1409–1428. <https://doi.org/10.1111/pce.12937>.
41. Hegedus, D., Yu, M., Baldwin, D., Gruber, M., Sharpe, A., Parkin, I., Whitwill, S., & Lydiate, D. (2003). Molecular characterization of *Brassica napus* NAC domain transcriptional activators induced in response to biotic and abiotic stress. *Plant molecular biology*, 53(3), 383–397. <https://doi.org/10.1023/b:plan.0000006944.61384.11>.

42. Hönig, M., Roeber, V. M., Schmülling, T., & Cortleven, A. (2023). Chemical priming of plant defense responses to pathogen attacks. *Frontiers in plant science*, 14, 1146577. <https://doi.org/10.3389/fpls.2023.1146577>.
43. Hornsey IS (2007) *Botrytis cinerea*. In the *Chemistry and Biology of Winemaking*. Royal Society of Chemistry, Cambridge, UK, pp 367–382
44. Hoshikawa, K., Fujita, S., Renhu, N., Ezura, K., Yamamoto, T., Nonaka, S., Ezura, H., & Miura, K. (2019). Efficient transient protein expression in tomato cultivars and wild species using agroinfiltration-mediated high expression system. *Plant cell reports*, 38(1), 75–84. <https://doi.org/10.1007/s00299-018-2350-1>.
45. Hruz, T., Laule, O., Szabo, G., Wessendorp, F., Bleuler, S., Oertle, L., Widmayer, P., Gruissem, W., & Zimmermann, P. (2008). Genevestigator v3: a reference expression database for the meta-analysis of transcriptomes. *Advances in bioinformatics*, 2008, 420747. <https://doi.org/10.1155/2008/420747>.
46. Hu, H., Dai, M., Yao, J., Xiao, B., Li, X., Zhang, Q., & Xiong, L. (2006). Overexpressing a NAM, ATAF, and CUC (NAC) transcription factor enhances drought resistance and salt tolerance in rice. *Proceedings of the National Academy of Sciences of the United States of America*, 103(35), 12987–12992. <https://doi.org/10.1073/pnas.0604882103>.
47. Huang, daW., Sherman, B. T., & Lempicki, R. A. (2009). Systematic and integrative analysis of large gene lists using DAVID bioinformatics resources. *Nature protocols*, 4(1), 44–57. <https://doi.org/10.1038/nprot.2008.211>.
48. Ibraheem, O., Botha, C. E., & Bradley, G. (2010). *In-silico* analysis of cis-acting regulatory elements in 5' regulatory regions of sucrose transporter gene families in rice (*Oryza sativa* Japonica) and *Arabidopsis thaliana*. *Computational biology and chemistry*, 34(5-6), 268–283. <https://doi.org/10.1016/j.compbiolchem.2010.09.003>.
49. Ijaz, Usman & Pervaiz, Tehreem & Ahmed, Temoor & Shahid, Muhammad & Noman, Muhammad & Nadeem, Majid & Azeem, Farrukh. (2020). Plant Cis-regulatory elements: methods of identification and applications. 8. 207-222. [10.35495/ajab.2019.08.352](https://doi.org/10.35495/ajab.2019.08.352).
50. Ison, J., Kalas, M., Jonassen, I., Bolser, D., Uludag, M., McWilliam, H., Malone, J., Lopez, R., Pettifer, S., & Rice, P. (2013). EDAM: an ontology of bioinformatics operations, types of data and identifiers, topics and formats. *Bioinformatics (Oxford, England)*, 29(10), 1325–1332. <https://doi.org/10.1093/bioinformatics/btt113>.
51. Jensen, M. K., Kjaersgaard, T., Nielsen, M. M., Galberg, P., Petersen, K., O'Shea, C., & Skriver, K. (2010). The *Arabidopsis thaliana* NAC transcription factor family: structure-function relationships and determinants of ANAC019 stress signalling. *The Biochemical journal*, 426(2), 183–196. <https://doi.org/10.1042/BJ20091234>.
52. Jeong, J. S., Kim, Y. S., Baek, K. H., Jung, H., Ha, S. H., Do Choi, Y., Kim, M., Reuzeau, C., & Kim, J. K. (2010). Root-specific expression of OsNAC10 improves drought tolerance and grain yield in rice under field drought conditions. *Plant physiology*, 153(1), 185–197. <https://doi.org/10.1104/pp.110.154773>.

53. Jiang, R. H., & Tyler, B. M. (2012). Mechanisms and evolution of virulence in oomycetes. *Annual review of phytopathology*, 50, 295–318. <https://doi.org/10.1146/annurev-phyto-081211-172912>.
54. Jiang, Y., & Deyholos, M. K. (2006). Comprehensive transcriptional profiling of NaCl-stressed *Arabidopsis* roots reveals novel classes of responsive genes. *BMC plant biology*, 6, 25. <https://doi.org/10.1186/1471-2229-6-25>.
55. Ju, Y. L., Min, Z., Yue, X. F., Zhang, Y. L., Zhang, J. X., Zhang, Z. Q., & Fang, Y. L. (2020). Overexpression of grapevine VvNAC08 enhances drought tolerance in transgenic *Arabidopsis*. *Plant physiology and biochemistry: PPB*, 151, 214–222. <https://doi.org/10.1016/j.plaphy.2020.03.028>.
56. Ju, Y. L., Yue, X. F., Min, Z., Wang, X. H., Fang, Y. L., & Zhang, J. X. (2020). VvNAC17, a novel stress-responsive grapevine (*Vitis vinifera* L.) NAC transcription factor, increases sensitivity to abscisic acid and enhances salinity, freezing, and drought tolerance in transgenic *Arabidopsis*. *Plant physiology and biochemistry: PPB*, 146, 98–111. <https://doi.org/10.1016/j.plaphy.2019.11.002>.
57. Kaur, G., & Pati, P. K. (2016). Analysis of cis-acting regulatory elements of Respiratory burst oxidase homolog (Rboh) gene families in *Arabidopsis* and rice provides clues for their diverse functions. *Computational biology and chemistry*, 62, 104–118. <https://doi.org/10.1016/j.compbiolchem.2016.04.002>.
58. Keller, M., Viret, O., & Cole, F. M. (2003). Botrytis cinerea Infection in Grape Flowers: Defense Reaction, Latency, and Disease Expression. *Phytopathology*, 93(3), 316–322. <https://doi.org/10.1094/PHYTO.2003.93.3.316>.
59. Kerchev, P., van der Meer, T., Sujeeth, N., Verlee, A., Stevens, C. V., Van Breusegem, F., & Gechev, T. (2020). Molecular priming as an approach to induce tolerance against abiotic and oxidative stresses in crop plants. *Biotechnology advances*, 40, 107503. <https://doi.org/10.1016/j.biotechadv.2019.107503>.
60. Khraiwesh, B., Zhu, J. K., & Zhu, J. (2012). Role of miRNAs and siRNAs in biotic and abiotic stress responses of plants. *Biochimica et biophysica acta*, 1819(2), 137–148. <https://doi.org/10.1016/j.bbagr.2011.05.001>.
61. Kim, H. S., Park, B. O., Yoo, J. H., Jung, M. S., Lee, S. M., Han, H. J., Kim, K. E., Kim, S. H., Lim, C. O., Yun, D. J., Lee, S. Y., & Chung, W. S. (2007). Identification of a calmodulin-binding NAC protein as a transcriptional repressor in *Arabidopsis*. *The Journal of biological chemistry*, 282(50), 36292–36302. <https://doi.org/10.1074/jbc.M705217200>.
62. Kim, Y. S., Kim, S. G., Park, J. E., Park, H. Y., Lim, M. H., Chua, N. H., & Park, C. M. (2006). A membrane-bound NAC transcription factor regulates cell division in *Arabidopsis*. *The Plant cell*, 18(11), 3132–3144. <https://doi.org/10.1105/tpc.106.043018>.
63. Kjaersgaard, T., Jensen, M. K., Christiansen, M. W., Gregersen, P., Kragelund, B. B., & Skriver, K. (2011). Senescence-associated barley NAC (NAM, ATAF1,2, CUC) transcription factor interacts with radical-induced cell death 1 through a disordered

- regulatory domain. *The Journal of biological chemistry*, 286(41), 35418–35429. <https://doi.org/10.1074/jbc.M111.247221>.
64. Kleinow, Tatjana & Himbert, Sarah & Krenz, Björn & Jeske, Holger & Koncz, Csaba. (2009). NAC domain transcription factor ATAF1 interacts with SNF1-related kinases and silencing of its subfamily causes severe developmental defects in Arabidopsis. *Plant Science*. 177. 360-370. <https://doi.org/10.1016/j.plantsci.2009.06.011>.
65. Ko, J. H., Yang, S. H., Park, A. H., Lerouxel, O., & Han, K. H. (2007). ANAC012, a member of the plant-specific NAC transcription factor family, negatively regulates xylary fiber development in Arabidopsis thaliana. *The Plant journal: for cell and molecular biology*, 50(6), 1035–1048. <https://doi.org/10.1111/j.1365-313X.2007.03109.x>.
66. Kole, C., Muthamilarasan, M., Henry, R., Edwards, D., Sharma, R., Abberton, M., Batley, J., Bentley, A., Blakeney, M., Bryant, J., Cai, H., Cakir, M., Cseke, L. J., Cockram, J., de Oliveira, A. C., De Pace, C., Dempewolf, H., Ellison, S., Gepts, P., Greenland, A., ... Prasad, M. (2015). Application of genomics-assisted breeding for generation of climate resilient crops: progress and prospects. *Frontiers in plant science*, 6, 563. <https://doi.org/10.3389/fpls.2015.00563>.
67. Koley, P., Brahmachari, S., Saha, A., Deb, C., Mondal, M., Das, N., Das, A., Lahiri, S., Das, M., Thakur, M., & Kundu, S. (2022). Phytohormone Priming of Tomato Plants Evoke Differential Behavior in Rhizoctonia solani During Infection, With Salicylate Priming Imparting Greater Tolerance Than Jasmonate. *Frontiers in plant science*, 12, 766095. <https://doi.org/10.3389/fpls.2021.766095>.
68. Koltai, H. and Volpin, H. (2003). Agricultural genomics: an approach to plant protection. *European journal of plant pathology* 109: 101-108.
69. Korf I. (2004). Gene finding in novel genomes. *BMC bioinformatics*, 5, 59. <https://doi.org/10.1186/1471-2105-5-59>.
70. Krzywinski, M., Schein, J., Birol, I., Connors, J., Gascoyne, R., Horsman, D., Jones, S. J., & Marra, M. A. (2009). Circos: an information aesthetic for comparative genomics. *Genome research*, 19(9), 1639–1645. <https://doi.org/10.1101/gr.092759.109>.
71. Kumari, S., & Ware, D. (2013). Genome-wide computational prediction and analysis of core promoter elements across plant monocots and dicots. *PloS one*, 8(10), e79011. <https://doi.org/10.1371/journal.pone.0079011>.
72. Kutach, A. K., Kadonaga, J. T. (2000). The downstream promoter element DPE appears to be as widely used as the TATA box in Drosophila core promoters. *Molecular and cellular biology*, 20(13), 4754–4764. <https://doi.org/10.1128/MCB.20.13.4754-4764.2000>.
73. Le Hénanff, G., Profizi, C., Courteaux, B., Rabenoelina, F., Gérard, C., Clément, C., Baillieux, F., Cordelier, S., & Dhondt-Cordelier, S. (2013). Grapevine NAC1 transcription factor as a convergent node in developmental processes, abiotic stresses, and necrotrophic/biotrophic pathogen tolerance. *Journal of experimental botany*, 64(16), 4877–4893. <https://doi.org/10.1093/jxb/ert277>.

74. Le, D. T., Nishiyama, R., Watanabe, Y., Mochida, K., Yamaguchi-Shinozaki, K., Shinozaki, K., & Tran, L. S. (2011). Genome-wide survey and expression analysis of the plant-specific NAC transcription factor family in soybean during development and dehydration stress. *DNA research: an international journal for rapid publication of reports on genes and genomes*, 18(4), 263–276. <https://doi.org/10.1093/dnares/dsr015>.
75. Lee S, Lee HJ, Huh SU, Paek KH, Ha JH, Park CM. (2014). The Arabidopsis NAC transcription factor NTL4 participates in a positive feedback loop that induces programmed cell death under heat stress conditions. *Plant Sci.* 2014; 227:76–83. <https://doi.org/10.1016/j.plantsci.2014.07.003>.
76. Lenka S and Bansal KC. (2019). Abiotic stress responsive *cis-acting regulatory elementss* (CREs) in rice (*Oryza sativa* L.) and other plants. *Plant J.* 13: 145-158.
77. Lescot, M., Déhais, P., Thijs, G., Marchal, K., Moreau, Y., Van de Peer, Y., Rouzé, P., & Rombauts, S. (2002). PlantCARE, a database of plant cis-acting regulatory elements and a portal to tools for in silico analysis of promoter sequences. *Nucleic acids research*, 30(1), 325–327. <https://doi.org/10.1093/nar/30.1.325>.
78. Levine M. (2010). Transcriptional enhancers in animal development and evolution. *Current biology: CB*, 20(17), R754–R763. <https://doi.org/10.1016/j.cub.2010.06.070>.
79. Li, L., He, Y., Zhang, Z., Shi, Y., Zhang, X., Xu, X., Wu, J. L., & Tang, S. (2021). OsNAC109 regulates senescence, growth and development by altering the expression of senescence- and phytohormone-associated genes in rice. *Plant molecular biology*, 105(6), 637–654. <https://doi.org/10.1007/s11103-021-01118-y>.
80. Lindbo J. A. (2007). High-efficiency protein expression in plants from agroinfection-compatible Tobacco mosaic virus expression vectors. *BMC biotechnology*, 7, 52. <https://doi.org/10.1186/1472-6750-7-52>.
81. Liu, Q., Yan, S., Huang, W., Yang, J., Dong, J., Zhang, S., Zhao, J., Yang, T., Mao, X., Zhu, X., & Liu, B. (2018). NAC transcription factor ONAC066 positively regulates disease resistance by suppressing the ABA signaling pathway in rice. *Plant molecular biology*, 98(4-5), 289–302. <https://doi.org/10.1007/s11103-018-0768-z>.
82. Liu, X., Hong, L., Li, X. Y., Yao, Y., Hu, B., & Li, L. (2011). Improved drought and salt tolerance in transgenic Arabidopsis overexpressing a NAC transcriptional factor from *Arachis hypogaea*. *Bioscience, biotechnology, and biochemistry*, 75(3), 443–450. <https://doi.org/10.1271/bbb.100614>.
83. Liu, Y., Sun, J., & Wu, Y. (2016). Arabidopsis ATAF1 enhances the tolerance to salt stress and ABA in transgenic rice. *Journal of plant research*, 129(5), 955–962. <https://doi.org/10.1007/s10265-016-0833-0>.
84. Lu, P. L., Chen, N. Z., An, R., Su, Z., Qi, B. S., Ren, F., Chen, J., & Wang, X. C. (2007). A novel drought-inducible gene, ATAF1, encodes a NAC family protein that negatively regulates the expression of stress-responsive genes in Arabidopsis. *Plant molecular biology*, 63(2), 289–305. <https://doi.org/10.1007/s11103-006-9089-8>.

85. Luscombe, N. M., Greenbaum, D., & Gerstein, M. (2001). What is bioinformatics? A proposed definition and overview of the field. *Methods of information in medicine*, 40(4), 346–358.
86. Masri, R., & Kiss, E. (2023). The role of NAC genes in response to biotic stresses in plants. *Physiological and Molecular Plant Pathology*, 102034. <https://doi.org/10.1016/j.pmpp.2023.102034>.
87. Mathew, I. E., Das, S., Mahto, A., & Agarwal, P. (2016). Three Rice NAC Transcription Factors Heteromerize and Are Associated with Seed Size. *Frontiers in plant science*, 7, 1638. <https://doi.org/10.3389/fpls.2016.01638>.
88. Mei, F., Chen, B., Li, F., Zhang, Y., Kang, Z., Wang, X., & Mao, H. (2021). Overexpression of the wheat NAC transcription factor TaSNAC4-3A gene confers drought tolerance in transgenic Arabidopsis. *Plant physiology and biochemistry: PPB*, 160, 37–50. <https://doi.org/10.1016/j.plaphy.2021.01.004>.
89. Morales-Valle, H., Silva, L. C., Paterson, R. R., Venâncio, A., & Lima, N. (2011). Effects of the origins of Botrytis cinerea on earthy aromas from grape broth media further inoculated with Penicillium expansum. *Food microbiology*, 28(5), 1048–1053. <https://doi.org/10.1016/j.fm.2011.02.005>.
90. Mou, Z., Fan, W., Dong, X. (2003). Inducers of plant systemic acquired resistance regulate NPR1 function through redox changes. *Cell*, 113(7), 935–944. [https://doi.org/10.1016/s0092-8674\(03\)00429-x](https://doi.org/10.1016/s0092-8674(03)00429-x).
91. Mount, D. W. (2001). *Bioinformatics: Sequence and genome analysis*. Cold Spring Harbor, N.Y: Cold Spring Harbor Laboratory Press.
92. Mundy J, Yamaguchi-Shinozaki K, Chua NH. (1990). Nuclear proteins bind conserved elements in the abscisic acid-responsive promoter of a rice rabgene. *Proc Natl Acad Sci U S A*. 1990;87(4):1406–10. <https://doi.org/10.1073/pnas.87.4.1406>.
93. Nakashima k, Takasaki H, Mizoi J, Shinozaki K, Yamaguchi-Shinozaki K. (2012). NAC transcription factors in plant abiotic stress responses, *Biochimica et Biophysica Acta (BBA) - Gene Regulatory Mechanisms*, Volume 1819, Issue 2, 2012, Pages 97-103, ISSN 1874-9399, <https://doi.org/10.1016/j.bbagrm.2011.10.00>.
94. Nakashima, K., Tran, L. S., Van Nguyen, D., Fujita, M., Maruyama, K., Todaka, D., Ito, Y., Hayashi, N., Shinozaki, K., & Yamaguchi-Shinozaki, K. (2007). Functional analysis of a NAC-type transcription factor OsNAC6 involved in abiotic and biotic stress-responsive gene expression in rice. *The Plant journal: for cell and molecular biology*, 51(4), 617–630. <https://doi.org/10.1111/j.1365-313X.2007.03168.x>
95. Narusaka Y, Nakashima K, Shinwari ZK, Sakuma Y, Furihata T, Abe H, Narusaka M, Shinozaki K, Yamaguchi-Shinozaki K. (2003). Interaction between two cis-acting elements, ABRE and DRE, in ABA dependent expression of Arabidopsis rd29 gene in response to dehydration and high-salinity stresses. *Plant J*. 2003;34(2):137–48. <https://doi.org/10.1046/j.1365-313X.2003.01708.x>.
96. Ng S, Ivanova A, Duncan O, Law SR, Van Aken O, De Clercq I, Wang Y, Carrie C, Xu L, Kmiec B, Walker H, Van Breusegem F, Whelan J, Giraud E. (2013). A membrane-

- bound NAC transcription factor, ANAC017, mediates mitochondrial retrograde signaling in Arabidopsis. *Plant Cell*. 2013;25(9):3450–71. <https://doi.org/10.1105/tpc.113.113985>.
97. Nuruzzaman, M., Manimekalai, R., Sharoni, A. M., Satoh, K., Kondoh, H., Ooka, H., & Kikuchi, S. (2010). Genome-wide analysis of NAC transcription factor family in rice. *Gene*, 465(1-2), 30–44. <https://doi.org/10.1016/j.gene.2010.06.008>.
98. Ogo Y, Kobayashi T, Itai RN, Nakanishi H, Kakei Y, Takahashi M, Toki S, Mori S, Nishizawa NK. (2008). A novel NAC transcription factor, IDEF2, that recognizes the iron deficiency responsive element 2 regulates the genes involved in iron homeostasis in plants. *J Biol Chem*. 2008;283(19):13407–17. <https://doi.org/10.1074/jbc.M708732200>.
99. Oh, S. K., Lee, S., Yu, S. H., & Choi, D. (2005). Expression of a novel NAC domain-containing transcription factor (CaNAC1) is preferentially associated with incompatible interactions between chili pepper and pathogens. *Planta*, 222(5), 876–887. <https://doi.org/10.1007/s00425-005-0030-1>.
100. Olsen, A. N., Ernst, H. A., Leggio, L. L., & Skriver, K. (2005). NAC transcription factors: structurally distinct, functionally diverse. *Trends in plant science*, 10(2), 79–87. <https://doi.org/10.1016/j.tplants.2004.12.010>.
101. Ooka, H., Satoh, K., Doi, K., Nagata, T., Otomo, Y., Murakami, K., Matsubara, K., Osato, N., Kawai, J., Carninci, P., Hayashizaki, Y., Suzuki, K., Kojima, K., Takahara, Y., Yamamoto, K., & Kikuchi, S. (2003). Comprehensive analysis of NAC family genes in *Oryza sativa* and *Arabidopsis thaliana*. *DNA research: an international journal for rapid publication of reports on genes and genomes*, 10(6), 239–247. <https://doi.org/10.1093/dnares/10.6.239>.
102. Pandey B, Prakash P, Verma PC and Srivastava R. (2019). Regulated Gene Expression by Synthetic Modulation of the Promoter Architecture in Plants. In book: Current Developments in Biotechnology and Bioengineering. pp. 235–255.
103. Pandey, S. P., & Somssich, I. E. (2009). The role of WRKY transcription factors in plant immunity. *Plant physiology*, 150(4), 1648–1655. <https://doi.org/10.1104/pp.109.138990>.
104. Papatheodorou, I., Moreno, P., Manning, J., Fuentes, A. M., George, N., Fexova, S., Fonseca, N. A., Füllgrabe, A., Green, M., Huang, N., Huerta, L., Iqbal, H., Jianu, M., Mohammed, S., Zhao, L., Jarnuczak, A. F., Jupp, S., Marioni, J., Meyer, K., Petryszak, R., ... Brazma, A. (2020). Expression Atlas update: from tissues to single cells. *Nucleic acids research*, 48(D1), D77–D83. <https://doi.org/10.1093/nar/gkz947>.
105. Pezet, R., Viret, O., Perret, C. and Tabacchi, R. (2003), Latency of *Botrytis cinerea* Pers.: Fr. and Biochemical Studies During Growth and Ripening of Two Grape Berry Cultivars, Respectively Susceptible and Resistant to Grey Mould. *Journal of Phytopathology*, 151: 208–214. <https://doi.org/10.1046/j.1439-0434.2003.00707.x>.
106. Pinheiro GL, Marques CS, Costa MD, Reis PA, Alves MS, Carvalho CM, Fietto GL, Fontes EP. (2009). Complete inventory of soybean NAC transcription factors: sequence conservation and expression analysis uncover their distinct roles in stress response. *Gene*. 2009;444(1-2):10–23. <https://doi.org/10.1016/j.gene.2009.05.012>.

107. Priest, H. D., Filichkin, S. A., & Mockler, T. C. (2009). Cis-acting regulatory elements in plant cell signaling. *Current opinion in plant biology*, 12(5), 643–649. <https://doi.org/10.1016/j.pbi.2009.07.016>.
108. Pruitt, R. N., Gust, A. A., & Nürnberger, T. (2021). Plant immunity unified. *Nature plants*, 7(4), 382–383. <https://doi.org/10.1038/s41477-021-00903-3>.
109. Prusky, D., Alkan, N., Mengiste, T., & Fluhr, R. (2013). Quiescent and necrotrophic lifestyle choice during postharvest disease development. *Annual review of phytopathology*, 51, 155–176. <https://doi.org/10.1146/annurev-phyto-082712-102349>.
110. Puranik, S., Sahu, P. P., Srivastava, P. S., & Prasad, M. (2012). NAC proteins: regulation and role in stress tolerance. *Trends in plant science*, 17(6), 369–381. <https://doi.org/10.1016/j.tplants.2012.02.004>.
111. Rao, Vaddi & Das, Sussant & Rao, Vaddi & Srinubabu, Gedela. (2008). Recent developments in life sciences research: Role of bioinformatics. *African Journal of Biotechnology* (ISSN: 1684-5315) Vol 7 Num 5. 7.
112. Raut S.A., Sathe S.R., Raut A. (2010). " Bioinformatics: Trends in gene expression analysis," 2010 International Conference on Bioinformatics and Biomedical Technology, 2010, pp. 97-100, [10.1109/ICBBT.2010.5479003](https://doi.org/10.1109/ICBBT.2010.5479003).
113. Ren, C., Liu, X., Zhang, Z., Wang, Y., Duan, W., Li, S., & Liang, Z. (2016). CRISPR/Cas9-mediated efficient targeted mutagenesis in Chardonnay (*Vitis vinifera* L.). *Scientific reports*, 6, 32289. <https://doi.org/10.1038/srep32289>.
114. Ren, T., Qu, F., & Morris, T. J. (2000). HRT gene function requires interaction between a NAC protein and viral capsid protein to confer resistance to turnip crinkle virus. *The Plant cell*, 12(10), 1917–1926. <https://doi.org/10.1105/tpc.12.10.1917>.
115. Rhaman, M. S., Imran, S., Rauf, F., Khatun, M., Baskin, C. C., Murata, Y., & Hasanuzzaman, M. (2020). Seed Priming with Phytohormones: An Effective Approach for the Mitigation of Abiotic Stress. *Plants* (Basel, Switzerland), 10(1), 37. <https://doi.org/10.3390/plants10010037>.
116. Rhee, S. Y., & Mutwil, M. (2014). Towards revealing the functions of all genes in plants. *Trends in plant science*, 19(4), 212–221. <https://doi.org/10.1016/j.tplants.2013.10.006>.
117. Rombauts, S., Florquin, K., Lescot, M., Marchal, K., Rouzé, P., & van de Peer, Y. (2003). Computational approaches to identify promoters and cis-regulatory elements in plant genomes. *Plant physiology*, 132(3), 1162–1176. <https://doi.org/10.1104/pp.102.017715>.
118. Rushton, P. J., Bokowiec, M. T., Han, S., Zhang, H., Brannock, J. F., Chen, X., Laudeman, T. W., & Timko, M. P. (2008). Tobacco transcription factors: novel insights into transcriptional regulation in the Solanaceae. *Plant physiology*, 147(1), 280–295. <https://doi.org/10.1104/pp.107.114041>.
119. Ryan, P. T., Ó'Maoláidigh, D. S., Drost, H. G., Kwaśniewska, K., Gabel, A., Grosse, I., Graciet, E., Quint, M., & Wellmer, F. (2015). Patterns of gene expression during

- Arabidopsis flower development from the time of initiation to maturation. *BMC genomics*, 16(1), 488. <https://doi.org/10.1186/s12864-015-1699-6>.
120. Sako, K., Nguyen, H. M., & Seki, M. (2021). Advances in Chemical Priming to Enhance Abiotic Stress Tolerance in Plants. *Plant & cell physiology*, 61(12), 1995–2003. <https://doi.org/10.1093/pcp/pcaa119>.
121. Schreiber, M., Stein, N., & Mascher, M. (2018). Genomic approaches for studying crop evolution. *Genome biology*, 19(1), 140. <https://doi.org/10.1186/s13059-018-1528-8>.
122. Seo P.J. (2014). Recent advances in plant membrane-bound transcription factor research: emphasis on intracellular movement. *J Integr Plant Biol.* 2014;56(4):334–42. <https://doi.org/10.1111/jipb.12139>.
123. Seo, P. J., Kim, S. G., & Park, C. M. (2008). Membrane-bound transcription factors in plants. *Trends in plant science*, 13(10), 550–556. <https://doi.org/10.1016/j.tplants.2008.06.008>.
124. Shao, H., Wang, H., & Tang, X. (2015). NAC transcription factors in plant multiple abiotic stress responses: progress and prospects. *Frontiers in plant science*, 6, 902. <https://doi.org/10.3389/fpls.2015.00902>.
125. Shen, Hui & Yin, Yanbin & Chen, Fang & Xu, Ying & Dixon, Richard. (2009). A Bioinformatic Analysis of NAC Genes for Plant Cell Wall Development in Relation to Lignocellulosic Bioenergy Production. *BioEnergy Research*. 2. 217-232. [10.1007/s12155-009-9047-9](https://doi.org/10.1007/s12155-009-9047-9).
126. Shinozaki, K., & Yamaguchi-Shinozaki, K. (2022). Functional genomics in plant abiotic stress responses and tolerance: From gene discovery to complex regulatory networks and their application in breeding. *Proceedings of the Japan Academy. Series B, Physical and biological sciences*, 98(8), 470–492. <https://doi.org/10.2183/pjab.98.024>.
127. Sipiczki M. (2006). Metschnikowia strains isolated from botrytized grapes antagonize fungal and bacterial growth by iron depletion. *Applied and environmental microbiology*, 72(10), 6716–6724. <https://doi.org/10.1128/AEM.01275-06>.
128. Song, S. Y., Chen, Y., Chen, J., Dai, X. Y., & Zhang, W. H. (2011). Physiological mechanisms underlying OsNAC5-dependent tolerance of rice plants to abiotic stress. *Planta*, 234(2), 331–345. <https://doi.org/10.1007/s00425-011-1403-2>.
129. Sperotto, R. A., Ricachenevsky, F. K., Duarte, G. L., Boff, T., Lopes, K. L., Sperb, E. R., Grusak, M. A., & Fett, J. P. (2009). Identification of up-regulated genes in flag leaves during rice grain filling and characterization of OsNAC5, a new ABA-dependent transcription factor. *Planta*, 230(5), 985–1002. <https://doi.org/10.1007/s00425-009-1000-9>.
130. Sun, L., Zhang, H., Li, D., Huang, L., Hong, Y., Ding, X. S., Nelson, R. S., Zhou, X., & Song, F. (2013). Functions of rice NAC transcriptional factors, ONAC122 and ONAC131, in defense responses against Magnaporthe grisea. *Plant molecular biology*, 81(1-2), 41–56. <https://doi.org/10.1007/s11103-012-9981-3>.

131. Sutton, J. C. 1998. Botrytis Fruit Rot (Gray Mold) and Blossom Blight. Pp. 28-31 in: Compendium of Strawberry Diseases, 2nd edition, Maas, J. L. (ed.). APS Press. St. Paul, MN. <https://doi.org/10.1080/07060669009501048>.
132. Szklarczyk, D., Gable, A. L., Lyon, D., Junge, A., Wyder, S., Huerta-Cepas, J., Simonovic, M., Doncheva, N. T., Morris, J. H., Bork, P., Jensen, L. J., & Mering, C. V. (2019). STRING v11: protein-protein association networks with increased coverage, supporting functional discovery in genome-wide experimental datasets. *Nucleic acids research*, 47(D1), D607–D613. <https://doi.org/10.1093/nar/gky1131>.
133. Szklarczyk, D., Kirsch, R., Koutrouli, M., Nastou, K., Mehryary, F., Hachilif, R., Gable, A. L., Fang, T., Doncheva, N. T., Pyysalo, S., Bork, P., Jensen, L. J., & von Mering, C. (2023). The STRING database in 2023: protein-protein association networks and functional enrichment analyses for any sequenced genome of interest. *Nucleic acids research*, 51(D1), D638–D646. <https://doi.org/10.1093/nar/gkac1000>.
134. Takasaki, H., Maruyama, K., Kidokoro, S., Ito, Y., Fujita, Y., Shinozaki, K., Yamaguchi-Shinozaki, K., & Nakashima, K. (2010). The abiotic stress-responsive NAC-type transcription factor OsNAC5 regulates stress-inducible genes and stress tolerance in rice. *Molecular genetics and genomics: MGG*, 284(3), 173–183. <https://doi.org/10.1007/s00438-010-0557-0>.
135. This, P., Lacombe, T., & Thomas, M. R. (2006). Historical origins and genetic diversity of wine grapes. *Trends in genetics : TIG*, 22(9), 511–519. <https://doi.org/10.1016/j.tig.2006.07.008>.
136. Toth, Z., Winterhagen, P., Kalapos, B., Su, Y., Kovacs, L., & Kiss, E. (2016). Expression of a Grapevine NAC Transcription Factor Gene Is Induced in Response to Powdery Mildew Colonization in Salicylic Acid-Independent Manner. *Scientific reports*, 6, 30825. <https://doi.org/10.1038/srep30825>.
137. Tran, L. S., Nakashima, K., Sakuma, Y., Simpson, S. D., Fujita, Y., Maruyama, K., Fujita, M., Seki, M., Shinozaki, K., & Yamaguchi-Shinozaki, K. (2004). Isolation and functional analysis of Arabidopsis stress-inducible NAC transcription factors that bind to a drought-responsive cis-element in the early responsive to dehydration stress 1 promoter. *The Plant cell*, 16(9), 2481–2498. <https://doi.org/10.1105/tpc.104.022699>.
138. Tran, L. S., Nishiyama, R., Yamaguchi-Shinozaki, K., & Shinozaki, K. (2010). Potential utilization of NAC transcription factors to enhance abiotic stress tolerance in plants by biotechnological approach. *GM crops*, 1(1), 32–39. <https://doi.org/10.4161/gmcr.1.1.10569>.
139. Tutar Y. (2014). miRNA and cancer; computational and experimental approaches. *Current pharmaceutical biotechnology*, 15(5), 429. <https://doi.org/10.2174/138920101505140828161335>.
140. Veloso, J., & van Kan, J. (2018). Many Shades of Grey in Botrytis-Host Plant Interactions. *Trends in plant science*, 23(7), 613–622. <https://doi.org/10.1016/j.tplants.2018.03.016>.

141. Wallace, I. M., Blackshields, G., & Higgins, D. G. (2005). Multiple sequence alignments. *Current opinion in structural biology*, 15(3), 261–266. <https://doi.org/10.1016/j.sbi.2005.04.002>.
142. Walls, R. L., Athreya, B., Cooper, L., Elser, J., Gandolfo, M. A., Jaiswal, P., Mungall, C. J., Preece, J., Rensing, S., Smith, B., & Stevenson, D. W. (2012). Ontologies as integrative tools for plant science. *American journal of botany*, 99(8), 1263–1275. <https://doi.org/10.3732/ajb.1200222>.
143. Wan, W. L., Fröhlich, K., Pruitt, R. N., Nürnberger, T., & Zhang, L. (2019). Plant cell surface immune receptor complex signaling. *Current opinion in plant biology*, 50, 18–28. <https://doi.org/10.1016/j.pbi.2019.02.001>.
144. Wang, N., Zheng, Y., Xin, H., Fang, L., & Li, S. (2013). Comprehensive analysis of NAC domain transcription factor gene family in *Vitis vinifera*. *Plant cell reports*, 32(1), 61–75. <https://doi.org/10.1007/s00299-012-1340-y>.
145. Wang, X., Basnayake, B. M., Zhang, H., Li, G., Li, W., Virk, N., Mengiste, T., & Song, F. (2009). The Arabidopsis ATAF1, a NAC transcription factor, is a negative regulator of defense responses against necrotrophic fungal and bacterial pathogens. *Molecular plant-microbe interactions: MPMI*, 22(10), 1227–1238. <https://doi.org/10.1094/MPMI-22-10-1227>.
146. Williamson, B., Tudzynski, B., Tudzynski, P., Van Kan, J.A.L. (2007), *Botrytis cinerea*: the cause of grey mould disease. *Molecular Plant Pathology*, 8: 561-580. <https://doi.org/10.1111/j.1364-3703.2007.00417.x>.
147. Wu, Y., Deng, Z., Lai, J., Zhang, Y., Yang, C., Yin, B., Zhao, Q., Zhang, L., Li, Y., Yang, C., & Xie, Q. (2009). Dual function of Arabidopsis ATAF1 in abiotic and biotic stress responses. *Cell research*, 19(11), 1279–1290. <https://doi.org/10.1038/cr.2009.108>.
148. Xiong, J. (2009). *Essential bioinformatics: introduction and biological databases*. Cambridge University press, USA. <http://www.cambridge.org/catalogue>.
149. Yang, C., Huang, Y., Lv, W., Zhang, Y., Bhat, J. A., Kong, J., Xing, H., Zhao, J., & Zhao, T. (2020). GmNAC8 acts as a positive regulator in soybean drought stress. *Plant science: an international journal of experimental plant biology*, 293, 110442. <https://doi.org/10.1016/j.plantsci.2020.110442>.
150. Yang, S. D., Seo, P. J., Yoon, H. K., & Park, C. M. (2011). The Arabidopsis NAC transcription factor VNI2 integrates abscisic acid signals into leaf senescence via the COR/RD genes. *The Plant cell*, 23(6), 2155–2168. <https://doi.org/10.1105/tpc.111.084913>.
151. Yildiz, I., Mantz, M., Hartmann, M., Zeier, T., Kessel, J., Thurow, C., Gatz, C., Petzsch, P., Köhrer, K., & Zeier, J. (2021). The mobile SAR signal N-hydroxypipicolinic acid induces NPR1-dependent transcriptional reprogramming and immune priming. *Plant physiology*, 186(3), 1679–1705. <https://doi.org/10.1093/plphys/kiab166>.
152. Yoshii, M., Shimizu, T., Yamazaki, M., Higashi, T., Miyao, A., Hirochika, H., & Omura, T. (2009). Disruption of a novel gene for a NAC-domain protein in rice confers

- resistance to Rice dwarf virus. *The Plant journal: for cell and molecular biology*, 57(4), 615–625. <https://doi.org/10.1111/j.1365-313X.2008.03712.x>.
153. Yoshii, M., Yamazaki, M., Rakwal, R., Kishi-Kaboshi, M., Miyao, A., & Hirochika, H. (2010). The NAC transcription factor RIM1 of rice is a new regulator of jasmonate signaling. *The Plant journal: for cell and molecular biology*, 61(5), 804–815. <https://doi.org/10.1111/j.1365-313X.2009.04107.x>.
154. You F. M. (2023). Plant Genomics-Advancing Our Understanding of Plants. *International journal of molecular sciences*, 24(14), 11528. <https://doi.org/10.3390/ijms241411528>.
155. Yu, C. S., Lin, C. J., & Hwang, J. K. (2004). Predicting subcellular localization of proteins for Gram-negative bacteria by support vector machines based on n-peptide compositions. *Protein science: a publication of the Protein Society*, 13(5), 1402–1406. <https://doi.org/10.1110/ps.03479604>.
156. Yuan, X., Wang, H., Cai, J. *et al.* NAC transcription factors in plant immunity. *Phytopathol Res* 1, 3 (2019). <https://doi.org/10.1186/s42483-018-0008-0>.
157. Zeller, G., Henz, S. R., Widmer, C. K., Sachsenberg, T., Rättsch, G., Weigel, D., & Laubinger, S. (2009). Stress-induced changes in the Arabidopsis thaliana transcriptome analyzed using whole-genome tiling arrays. *The Plant journal: for cell and molecular biology*, 58(6), 1068–1082. <https://doi.org/10.1111/j.1365-313X.2009.03835.x>.
158. Zhang, N., Yuan, S., Zhao, C., Park, R. F., Wen, X., Yang, W., Zhang, N., & Liu, D. (2021). TaNAC35 acts as a negative regulator for leaf rust resistance in a compatible interaction between common wheat and Puccinia triticina. *Molecular genetics and genomics: MGG*, 296(2), 279–287. <https://doi.org/10.1007/s00438-020-01746-x>.
159. Zheng, X., Chen, B., Lu, G., & Han, B. (2009). Overexpression of a NAC transcription factor enhances rice drought and salt tolerance. *Biochemical and biophysical research communications*, 379(4), 985–989. <https://doi.org/10.1016/j.bbrc.2008.12.163>.
160. Zhong R, Richardson EA, Ye ZH. (2007). Two NAC domain transcription factors, SND1 and NST1, function redundantly in regulation of secondary wall synthesis in fibers of Arabidopsis. *Planta*. 2007;225(6):1603–11. <https://doi.org/10.1007/s00425-007-0498-y>.
161. Zhong, R., Lee, C., & Ye, Z. H. (2010). Global analysis of direct targets of secondary wall NAC master switches in Arabidopsis. *Molecular plant*, 3(6), 1087–1103. <https://doi.org/10.1093/mp/ssq062>.
162. Zhu J. K. (2002). Salt and drought stress signal transduction in plants. *Annual review of plant biology*, 53, 247–273. <https://doi.org/10.1146/annurev.arplant.53.091401.143329>.

A2: Supplementary data

Table 7.1. Characterization and distribution of the VvNAC genes.

Species	Gene ID	Length	Transcript Name	MW (Da)	IP	Orthologues in <i>A. thaliana</i>
Grape	VvNAC01	251	VIT_01s0146g00280	28601.2	9.6551	ANAC083 (AT5G13180.1)
Grape	VvNAC02	320	VIT_02s0154g00020	37460.2	7.8983	ANAC007 (AT1G12260.1)
Grape	VvNAC03	331	VIT_00s0375g00040	36744.2	9.7936	ANAC025 (AT1G61110.1)
Grape	VvNAC04	279	VIT_04s0008g00150	32252.8	7.6617	ANAC036 (AT2G17040.1)
Grape	VvNAC05	284	VIT_17s0000g06400	32180.8	10.2287	ANAC100 (AT5G61430.1)
Grape	VvNAC06	327	VIT_17s0000g03660	37083.7	8.4093	ANAC058 (AT3G18400.1)
Grape	VvNAC07	209	VIT_17s0000g00770	23960.8	8.0627	ANAC083 (AT5G13180.1)
Grape	VvNAC08	263	VIT_18s0001g02300	30130.2	7.9471	ANAC002(AT1G01720.1)
Grape	VvNAC09	355	VIT_18s0001g11980	38637	5.4619	ANAC057(AT3G17730.1)
Grape	VvNAC10	353	VIT_14s0108g00860	39332.4	8.2468	ANAC094(AT5G39820.1)
Grape	VvNAC11	290	VIT_14s0108g01070	32887.3	7.8281	ANAC100(AT5G61430.1)
Grape	VvNAC12	430	VIT_01s0011g02990	48383.6	4.7046	ANAC008(AT1G25580.1)
Grape	VvNAC13	365	VIT_02s0012g01040	41109.5	4.5281	ANAC071(AT4G17980.1)
Grape	VvNAC14	320	VIT_18s0122g00800	36190	6.6689	ANAC031 (AT1G76420.1)
Grape	VvNAC15	572	VIT_18s0001g01820	63327.9	4.4857	ANAC017 (AT1G34190.1)
Grape	VvNAC16	346	VIT_19s0014g02200	38342.9	8.002	ANAC098 (AT5G53950.1)
Grape	VvNAC17	333	VIT_19s0014g03290	37190.8	8.8885	ANAC072 (AT4G27410.2)
Grape	VvNAC18	331	VIT_19s0014g03300	36791.5	8.888	ANAC056 (AT3G15510.1)
Grape	VvNAC19	454	VIT_11s0016g02880	51170.8	6.8955	ANAC075 (AT4G29230.1)
Grape	VvNAC20	559	VIT_13s0019g05240	62299.2	4.2277	ANAC078 (AT5G04410.1)
Grape	VvNAC21	385	VIT_13s0019g05230	43682.5	7.6375	ANAC050 (AT3G10480.1)
Grape	VvNAC22	345	VIT_16s0022g00690	40174.2	6.8679	ANAC007 (AT1G12260.1)
Grape	VvNAC23	274	VIT_04s0023g03110	32148.3	9.4087	ANAC037 (AT2G18060.1)
Grape	VvNAC72	659	VIT_02s0025g01870	74672.92	7.26	ANAC083 (AT5G13180.1)
Grape	VvNAC24	357	VIT_02s0025g02710	40855.6	7.1335	ANAC043 (AT2G46770.1)
Grape	VvNAC25	632	VIT_02s0025g03020	70493.1	4.811	ANAC014 (AT1G33060.1)
Grape	VvNAC26	282	VIT_01s0026g02710	32458.7	8.0663	ANAC029 (AT1G69490.1)
Grape	VvNAC27	385	VIT_01s0026g02360	43110.1	8.4525	ANAC009 (AT1G26870.1)
Grape	VvNAC28	295	VIT_19s0027g00910	33681	8.1248	ANAC042 (AT2G43000.1)
Grape	VvNAC29	214	VIT_19s0027g00880	25648.6	10.3268	ANAC042 (AT2G43000.1)
Grape	VvNAC30	277	VIT_19s0027g00870	32287.3	7.8995	ANAC042 (AT2G43000.1)
Grape	VvNAC31	277	VIT_19s0027g00860	32731.8	7.9703	ANAC042 (AT2G43000.1)
Grape	VvNAC32	284	VIT_19s0027g00840	32932	7.53	ANAC042 (AT2G43000.1)
Grape	VvNAC33	295	VIT_19s0027g00230	33364.1	6.767	ANAC022 (AT1G56010.2)
Grape	VvNAC34	303	VIT_12s0028g03050	34457.8	8.6499	ANAC073 (AT4G28500.1)
Grape	VvNAC36	298	VIT_12s0028g00860	34198.3	8.4111	ANAC042 (AT2G43000.1)
Grape	VvNAC35	480	VIT_12s0028g02670	53469.5	5.12	ANAC062 (AT3G49530.1)
Grape	VvNAC37	311	VIT_10s0003g00350	35466.8	6.8317	ANAC073 (AT4G28500.1)
Grape	VvNAC38	484	VIT_10s0003g00500	53537.14	5.58	ANAC028 (AT1G65910.1)
Grape	VvNAC39	265	VIT_07s0031g02610	30286.2	9.1243	ANAC002 (AT1G01720.1)

7. Appendices

Grape	VvNAC40	204	VIT_12s0035g02020	24138.6	10.067	ANAC034 (AT2G02450.1)
Grape	VvNAC41	279	VIT_03s0038g03410	31946.5	8.5729	ANAC036 (AT2G17040.1)
Grape	VvNAC42	376	VIT_06s0004g03080	42958.1	8.8622	ANAC025 (AT1G61110.1)
Grape	VvNAC43	364	VIT_06s0004g02340	41280.7	4.9285	ANAC029 (AT1G69490.1)
Grape	VvNAC44	297	VIT_06s0004g00020	34433.6	5.8409	ANAC032 (AT1G77450.1)
Grape	VvNAC45	349	VIT_08s0040g02110	40037	8.5158	ANAC025 (AT1G61110.1)
Grape	VvNAC46	225	VIT_18s0041g00700	26238.6	8.9381	ANAC090 (AT5G22380.1)
Grape	VvNAC47	198	VIT_04s0044g01500	22707.4	4.1292	ANAC104 (AT5G64530.1)
Grape	VvNAC48	560	VIT_04s0044g01220	62640.2	4.8558	ANAC103 (AT5G64060.1)
Grape	VvNAC49	352	VIT_15s0048g02660	40519.1	7.4725	ANAC043 (AT2G46770.1)
Grape	VvNAC50	128	VIT_15s0048g02340	14764	10.0079	ANAC050 (AT3G10480.1)
Grape	VvNAC51	202	VIT_15s0048g02320	22668.8	6.3591	ANAC116 (AT4G35580.1)
Grape	VvNAC52	218	VIT_15s0048g02310	24590.8	6.9826	ANAC116 (AT4G35580.1)
Grape	VvNAC53	455	VIT_15s0048g02300	51587.7	5.3227	ANAC116 (AT4G35580.1)
Grape	VvNAC54	994	VIT_15s0048g02280	110673	7.8715	ANAC116 (AT4G35580.2)
Grape	VvNAC55	335	VIT_15s0048g02270	37092.7	7.2815	ANAC116 (AT4G35580.1)
Grape	VvNAC56	281	VIT_07s0005g03610	31942	6.5797	ANAC074 (AT4G28530.1)
Grape	VvNAC57	238	VIT_17s0053g00740	27340.6	4.622	ANAC057 (AT3G17730.1)
Grape	VvNAC58	192	VIT_12s0055g00510	21978.4	4.9228	ANAC104 (AT5G64530.1)
Grape	VvNAC59	398	VIT_14s0066g00360	44158.1	6.2255	ANAC044 (AT3G01600.1)
Grape	VvNAC60	661	VIT_08s0007g07670	74009	7.0075	ANAC047 (AT3G04070.1)
Grape	VvNAC61	390	VIT_08s0007g07640	44064.2	8.248	ANAC087 (AT5G18270.1)
Grape	VvNAC62	217	VIT_08s0007g02940	24445.9	8.098	ANAC090 (AT5G22380.1)
Grape	VvNAC63	313	VIT_18s0072g01060	36310.7	6.3205	ANAC030 (AT1G71930.1)
Grape	VvNAC64	634	VIT_04s0079g00280	72544.7	5.2116	ANAC028 (AT1G65910.1)
Grape	VvNAC65	325	VIT_04s0008g02710	36575.4	8.6525	ANAC039 (AT2G24430.2)
Grape	VvNAC66	164	VIT_04s0008g06550	18677.4	8.1931	ANAC025 (AT1G61110.1)
Grape	VvNAC67	375	VIT_06s0080g00970	42903.5	6.3296	ANAC040 (AT2G27300.1)
Grape	VvNAC68	272	VIT_19s0085g01210	31198.3	8.0124	ANAC033 (AT1G79580.1)
Grape	VvNAC69	281	VIT_19s0085g00950	32077.6	5.2474	ANAC116 (AT4G35580.1)
Grape	VvNAC70	323	VIT_18s0089g01120	37007.3	8.9677	ANAC070 (AT4G10350.1)
Grape	VvNAC71	578	VIT_16s0098g00760	64685.1	4.6747	ANAC116 (AT4G35580.1)
Grape	VvNAC73	162	VIT_14s0068g01490	18617.7	10.5263	ANAC083 (AT5G13180.1)
Grape	VvNAC74	232	VIT_06s0080g00780	26616.9	7.6584	ANAC090 (AT5G22380.1)

7. Appendices

Table 7.2. Detailed information of protein-to-protein interaction network in VvNAC proteins based on known *Arabidopsis thaliana* proteins

Query Item	Preferred Name	String Id
<i>VvNAC61</i>	<i>ANAC087</i>	3702.AT5G18270.1
<i>VvNAC08 and VvNAC39</i>	<i>ATAF1</i>	3702.AT1G01720.1
<i>VvNAC03, VvNAC42, VvNAC45 and VvNAC66</i>	<i>NAC025</i>	3702.AT1G61110.1
<i>VvNAC64</i>	<i>NAC028</i>	3702.AT1G65910.1
<i>VvNAC44</i>	<i>NAC032</i>	3702.AT1G77450.1
<i>VvNAC04 and VvNAC41</i>	<i>NAC036</i>	3702.AT2G17040.1
<i>VvNAC65</i>	<i>NAC038</i>	3702.AT2G24430.2
<i>VvNAC28, VvNAC29, VvNAC30, VvNAC31, VvNAC32 and VvNAC36</i>	<i>NAC042</i>	3702.AT2G43000.1
<i>VvNAC59</i>	<i>NAC044</i>	3702.AT3G01600.1
<i>VvNAC60</i>	<i>NAC047</i>	3702.AT3G04070.1
<i>VvNAC21 and VvNAC50</i>	<i>NAC050</i>	3702.AT3G10480.3
<i>VvNAC09 and VvNAC57</i>	<i>NAC057</i>	3702.AT3G17730.1
<i>VvNAC70</i>	<i>NAC070</i>	3702.AT4G10350.1
<i>VvNAC34 and VvNAC37</i>	<i>NAC073</i>	3702.AT4G28500.1
<i>VvNAC56</i>	<i>NAC074</i>	3702.AT4G28530.1
<i>VvNAC01, VvNAC07, and VvNAC73</i>	<i>NAC083</i>	3702.AT5G13180.1
<i>VvNAC46 and VvNAC62</i>	<i>NAC090</i>	3702.AT5G22380.1
<i>VvNAC33</i>	<i>NAC1</i>	3702.AT1G56010.2
<i>VvNAC48</i>	<i>NAC103</i>	3702.AT5G64060.1
<i>VvNAC26, VvNAC43</i>	<i>NAP</i>	3702.AT1G69490.1
<i>VvNAC24 and VvNAC49</i>	<i>NST1</i>	3702.AT2G46770.1
<i>VvNAC67</i>	<i>NTL8</i>	3702.AT2G27300.1
<i>VvNAC51, VvNAC52, VvNAC53, VvNAC54, VvNAC55, VvNAC69, and VvNAC71</i>	<i>NTL9</i>	3702.AT4G35580.2
<i>VvNAC63</i>	<i>VND7</i>	3702.AT1G71930.1
<i>VvNAC47 and VvNAC58</i>	<i>XND1</i>	3702.AT5G64530.1

Annotation
Involved in chlorophyll catabolic processes, such as NYC1, SGR1, SGR2 and PAO
Transcript level increases in response to wounding and abscisic acid.
May be associated with anther development and pollen production.
Involved in multicellular organismal development, regulation of transcription.
Positively regulates age- dependent senescence, dark-induced leaf senescence and stress-induced senescence.
Involved in leaf and inflorescence stem morphogenesis.
Involved in multicellular organismal development, regulation of transcription.
Negative regulator of leaf senescence and enhances tolerance to various abiotic stresses through the regulation of DREB2A
Involved in multicellular organismal development, regulation of transcription.
Involved in ethylene biosynthesis. Mediates waterlogging- induced hyponastic leaf movement, and cell expansion.

7. Appendices

Involves in multicellular organismal development, regulation of transcription.
Involves in multicellular organismal development, regulation of transcription.
Regulates root cap maturation, in a partially redundant fashion.
Plays a regulatory role in the development of secondary cell wall fibers.
Involves in multicellular organismal development, regulation of transcription.
Negatively regulates the xylem vessel formation and mediates signaling crosstalk between salt stress response and leaf aging process.
Involves in multicellular organismal development, regulation of transcription.
involved in shoot apical meristem formation and auxin-mediated lateral root formation.
Involves in multicellular organismal development, regulation of transcription.
Increases levels of the senescence-inducing hormone abscisic acid (ABA). Involved in the control of dehydration in senescing leaves.
Transcription activator of genes involved in biosynthesis of secondary walls.
Regulates gibberellic acid-mediated salt- responsive repression of seed germination and flowering.
Functions synergistically with SNI1 as negative regulator of pathogen induced PR1 expression.
Involves in xylem formation in roots and shoots.
Negatively regulates secondary cell wall fibre synthesis and programmed cell death.

Table 7.3. The description of cis-acting regulatory elements of the *VvNAC* genes associated with plant stress, hormone, and development process.

Element name	Function
TGA-element	auxin-responsive element
TGA-box	part of an auxin-responsive element
TATC-box	<i>cis</i> -acting element involved in gibberellin-responsiveness
P-box	gibberellin-responsive element
ABRE	<i>cis</i> -acting element involved in the abscisic acid responsiveness
TCA-element	<i>cis</i> -acting element involved in salicylic acid responsiveness
CGTCA-motif	<i>cis</i> -acting regulatory element involved in the MeJA-responsiveness
TGACG-motif	<i>cis</i> -acting regulatory element involved in the MeJA-responsiveness
G-box	<i>cis</i> -acting regulatory element involved in light responsiveness
GATA-motif	part of a light responsive element
GT1-motif	light responsive element
circadian	<i>cis</i> -acting regulatory element involved in circadian control
LTR	<i>cis</i> -acting element involved in low-temperature responsiveness
ARE	<i>cis</i> -acting regulatory element essential for the anaerobic induction
MBS	MYB binding site involved in drought-inducibility
Box 4	part of a conserved DNA module involved in light responsiveness
TCT-motif	part of a light responsive element
TCCC-motif	part of a light responsive element
TC-rich repeats	<i>cis</i> -acting element involved in defense and stress responsiveness
DRE core	dehydration-responsive element
W box	elicitor responsiveness (disease resistance)
WRE3	wound response element
O2-site	<i>cis</i> -acting regulatory element involved in zein metabolism regulation
F-box	plant vegetative and reproduction growth and development; cell death and defense
as-1	root specific expression
AC-I	vascular-specific expression
STRE	Stress Response Element
HD-Zip1	<i>cis</i> -regulatory element for the leaf development
MBSI	involved in flavonoid biosynthesis
MYC	drought responsiveness
MYB	drought responsiveness
ERE	ethylene-responsive element
GARE-motif	<i>cis</i> -acting element involved in gibberellin-responsiveness
AuxRR-core	<i>cis</i> -acting regulatory element involved in auxin responsiveness

7. Appendices

Table 7.4. Microarrays data profiles related to *Vitis Vinifera* NAC genes in different stages of noble rot caused by *Botrytis cinerea*.

Gene ID	S1	S1; <i>B. cinerea</i> vs control. pValue	S2	S2; <i>B. cinerea</i> vs control. pValue	S3	S3; <i>B. cinerea</i> vs control. pValue
VvNAC01	0.2	0.535123329	1.4	4.05E-13	1	8.31E-07
VvNAC03	0.5	8.97E-04	1.1	1.69E-15	1.8	1.81E-36
VvNAC05	0.2	0.054213407	0.5	0.01939994		-
VvNAC07	0.2	0.784384669	1.1	0.119409044	0.2	0.808673942
VvNAC08	0.9	7.02E-08	3	2.76E-55	2.9	7.35E-108
VvNAC09	-	-	-0.3	0.073799196	-0.1	0.823009682
VvNAC11	-	-	-0.7	4.31E-06	0.1	0.591375038
VvNAC12	0.1	0.579271484	-1	5.20E-06	-0.2	0.495032535
VvNAC13	-0.3	0.371289174	-2.2	1.37E-24	-1.4	7.28E-12
VvNAC15	0.1	0.790590977	0.2	0.070755021	0.5	5.39E-06
VvNAC16	0.5	0.556117243	-	-	-	-
VvNAC18	-	-	0.8	1.86E-10	0.5	2.62E-05
VvNAC19	0.3	0.596523954	-0.7	0.265140926	-0.1	0.885888324
VvNAC20	-	-	0.4	0.003475854	0.3	0.002903393
VvNAC21	0.1	0.866239969	0.1	0.84133058	-0.4	0.537836971
VvNAC25	-0.2	0.552255046	0.5	3.73E-04	0.2	0.215992073
VvNAC26	0.1	0.85741699	-0.9	0.007091799	-1.1	0.003552427
VvNAC30	2.4	1.38E-05	3.7	2.31E-04	3.8	2.71E-04
VvNAC31	2.2	1.16E-04	3.3	0.001521837	2.5	0.036107373
VvNAC32	0.4	0.665733821	1.2	0.36443176	1.6	0.21718526
VvNAC33	0.2	0.753921202	1.4	2.41E-06	1.2	7.67E-04
VvNAC34	0.2	0.86602089	-	-	0.3	0.775124288
VvNAC35	0.1	0.738531823	-0.1	0.768116371	-0.1	0.594757038
VvNAC36	2.8	3.17E-26	1.1	0.015461522	0.9	0.115624957
VvNAC37	0.3	0.389041352	-	-	0.5	0.02987791
VvNAC38		-	-	-	0.1	0.451716644
VvNAC39	2.8	2.59E-34	5.6	0	5.5	0
VvNAC40	0.1	0.971570803	-	-	-	-
VvNAC41	1.2	0.06726654	2.7	0.021384867	-	-
VvNAC42	0.2	0.842798049	0.6	0.444459485	0.6	0.524252737
VvNAC44	2.7	5.97E-08	5	3.16E-22	6	7.89E-49
VvNAC45	0.3	0.818243519	-1.2	0.322247125	-0.1	0.947742753
VvNAC47	0.3	0.804206544	-0.2	0.873140439	-	-
VvNAC48	-	-	0.7	2.53E-09	0.4	2.29E-05
VvNAC51	-	-	-2.1	0.004527507	-2	0.016130879
VvNAC52	-0.3	0.705051056	-1.4	0.17591224	-2.1	0.053624394
VvNAC53	-0.7	0.007015424	-2.6	4.56E-15	-2.1	2.51E-12
VvNAC54	-0.2	0.780162316	-2	4.88E-07	-2.4	1.44E-06
VvNAC55	-0.6	0.007133217	-2.2	6.75E-19	-1.5	9.42E-09
VvNAC56	0.4	0.658823723	-	-	-	-
VvNAC59	-0.2	0.546888979	-0.7	0.025259741	-0.4	0.181143476

7. Appendices

VvNAC60	1.5	1.19E-16	3.6	4.03E-73	3	2.74E-81
VvNAC61	0.9	1.98E-14	2.9	1.74E-60	2.1	1.36E-34
VvNAC64	1.6	1.54E-08	1.1	0.019968952	0.7	0.169537728
VvNAC67	0.4	0.381666354	0.4	0.341189888	-0.1	0.874983883
VvNAC71	-0.5	0.009982447	-0.8	9.35E-05	-0.2	0.205413888
VvNAC72	0.2	0.840394162	0.2	0.881535544	-	-
VvNAC73	3	3.26E-17	5.2	3.54E-49	5.4	4.20E-59
VvNAC74	1.7	0.002941632	3.5	6.48E-06	2.2	0.045673339

Table 7.5. Calculation of the Euclidean distance of the clusters. In order to assess the dissimilarity between two clusters, a pairwise comparison of all VvNAC genes within each cluster was conducted, wherein the sum of differences in expression across various infection stages was computed.

Clusters	Dif b/t which VvNAC genes	D 1² (S1)	D 2² (S2)	D 3² (S3)	Sum of difs	ED
Clusters 2&3	<i>NAC31 & NAC13</i>	6.25	30.25	15.21	51.71	7.190967
	<i>NAC31 & NAC51</i>	4.84	29.16	20.25	54.25	7.36546
	<i>NAC31 & NAC52</i>	6.25	22.09	21.16	49.5	7.035624
	<i>NAC31 & NAC53</i>	8.41	34.81	21.16	64.38	8.023715
	<i>NAC31 & NAC54</i>	5.76	28.09	24.01	57.86	7.606576
	<i>NAC31 & NAC55</i>	7.84	30.25	16	54.09	7.35459
	<i>NAC30 & NAC13</i>	7.29	34.81	27.04	69.14	8.315047
	<i>NAC30 & NAC51</i>	5.76	33.64	33.64	73.04	8.546344
	<i>NAC30 & NAC52</i>	7.29	26.01	34.81	68.11	8.252878
	<i>NAC30 & NAC53</i>	9.61	39.69	34.81	84.11	9.17115
	<i>NAC30 & NAC54</i>	6.76	32.49	38.44	77.69	8.814193
	<i>NAC30 & NAC55</i>	9	34.81	28.09	71.9	8.479387
	<i>NAC61 & NAC13</i>	1.44	26.01	12.25	39.7	6.300794
	<i>NAC61 & NAC51</i>	0.81	25	16.81	42.62	6.528399
	<i>NAC61 & NAC52</i>	1.44	18.49	17.64	37.57	6.129437
	<i>NAC61 & NAC53</i>	2.56	30.25	17.64	50.45	7.102816
	<i>NAC61 & NAC54</i>	1.21	24.01	20.25	45.47	6.743145
	<i>NAC61 & NAC55</i>	2.25	26.01	12.96	41.22	6.42028
	<i>NAC08 & NAC13</i>	1.44	27.04	18.49	46.97	6.853466
	<i>NAC08 & NAC51</i>	0.81	26.01	24.01	50.83	7.129516
	<i>NAC08 & NAC52</i>	1.44	19.36	25	45.8	6.76757
	<i>NAC08 & NAC53</i>	2.56	31.36	25	58.92	7.675936
	<i>NAC08 & NAC54</i>	1.21	25	28.09	54.3	7.368853
	<i>NAC08 & NAC55</i>	2.25	27.04	19.36	48.65	6.974955
	<i>NAC64 & NAC13</i>	3.61	10.89	4.41	18.91	4.348563
	<i>NAC64 & NAC51</i>	2.56	10.24	7.29	20.09	4.482187
<i>NAC64 & NAC52</i>	3.61	6.25	7.84	17.7	4.207137	
<i>NAC64 & NAC53</i>	5.29	13.69	7.84	26.82	5.178803	
<i>NAC64 & NAC54</i>	3.24	9.61	9.61	22.46	4.739198	

7. Appendices

	<i>NAC64 & NAC55</i>	4.84	10.89	4.84	20.57	4.535416
	<i>NAC41 & NAC13</i>	2.25	24.01	1.96	28.22	5.31225
	<i>NAC41 & NAC51</i>	1.44	23.04	4	28.48	5.336666
	<i>NAC41 & NAC52</i>	2.25	16.81	4.41	23.47	4.844585
	<i>NAC41 & NAC53</i>	3.61	28.09	4.41	36.11	6.00916
	<i>NAC41&NAC54</i>	1.96	22.09	5.76	29.81	5.459853
	<i>NAC41&NAC55</i>	3.24	24.01	2.25	29.5	5.43139
	<i>NAC36&NAC13</i>	9.61	10.89	5.29	25.79	5.078386
	<i>NAC36&NAC51</i>	7.84	10.24	8.41	26.49	5.146844
	<i>NAC36&NAC52</i>	9.61	6.25	9	24.86	4.98598
	<i>NAC36&NAC53</i>	12.25	13.69	9	34.94	5.911007
	<i>NAC36&NAC54</i>	9	9.61	10.89	29.5	5.43139
	<i>NAC36&NAC55</i>	11.56	10.89	5.76	28.21	5.311309
	<i>NAC74&NAC13</i>	4	32.49	12.96	49.45	7.032069
	<i>NAC74&NAC51</i>	2.89	31.36	17.64	51.89	7.203471
	<i>NAC74&NAC52</i>	4	24.01	18.49	46.5	6.819091
	<i>NAC74&NAC53</i>	5.76	37.21	18.49	61.46	7.839643
	<i>NAC74&NAC54</i>	3.61	30.25	21.16	55.02	7.417547
	<i>NAC74&NAC55</i>	5.29	32.49	13.69	51.47	7.17426
	<i>NAC61&NAC13</i>	1.44	26.01	12.25	39.7	6.300794
	<i>NAC61&NAC51</i>	0.81	25	16.81	42.62	6.528399
	<i>NAC61&NAC52</i>	1.44	18.49	17.64	37.57	6.129437
	<i>NAC61&NAC53</i>	2.56	30.25	17.64	50.45	7.102816
	<i>NAC61&NAC54</i>	1.21	24.01	20.25	45.47	6.743145
	<i>NAC61&NAC55</i>	2.25	26.01	12.96	41.22	6.42028
Clusters 3&4	<i>NAC36&NAC73</i>	0.04	16.81	20.25	37.1	6.090977
	<i>NAC36&NAC39</i>	0	20.25	21.16	41.41	6.43506
	<i>NAC36&NAC44</i>	0.01	15.21	26.01	41.23	6.421059
	<i>NAC41&NAC73</i>	3.24	6.25	29.16	38.65	6.216912
	<i>NAC41&NAC39</i>	2.56	8.41	30.25	41.22	6.42028
	<i>NAC41&NAC44</i>	2.25	5.29	36	43.54	6.598485
	<i>NAC64&NAC73</i>	1.96	16.81	22.09	40.86	6.392183
	<i>NAC64&NAC39</i>	1.44	20.25	23.04	44.73	6.688049
	<i>NAC64&NAC44</i>	1.21	15.21	28.09	44.51	6.671582
	<i>NAC08&NAC73</i>	4.41	4.84	6.25	15.5	3.937004
	<i>NAC08&NAC39</i>	3.61	6.76	6.76	17.13	4.13884
	<i>NAC08&NAC44</i>	3.24	4	9.61	16.85	4.104875
	<i>NAC61&NAC73</i>	4.41	5.29	10.89	20.59	4.537621
	<i>NAC61&NAC39</i>	3.61	7.29	11.56	22.46	4.739198
	<i>NAC61&NAC44</i>	3.24	4.41	15.21	22.86	4.781213
	<i>NAC30&NAC73</i>	0.36	2.25	2.56	5.17	2.273763
	<i>NAC30&NAC39</i>	0.16	3.61	2.89	6.66	2.580698
	<i>NAC30&NAC44</i>	0.09	1.69	4.84	6.62	2.572936
	<i>NAC31&NAC73</i>	0.64	3.61	8.41	12.66	3.558089
	<i>NAC31&NAC39</i>	0.36	5.29	9	14.65	3.827532
	<i>NAC31&NAC44</i>	0.25	2.89	12.25	15.39	3.923009

7. Appendices

	<i>NAC60&NAC73</i>	2.25	2.56	5.76	10.57	3.251154
	<i>NAC60&NAC39</i>	1.69	4	6.25	11.94	3.455431
	<i>NAC60&NAC44</i>	1.44	1.96	9	12.4	3.521363
	<i>NAC74&NAC73</i>	1.69	2.89	10.24	14.82	3.849675
	<i>NAC74&NAC39</i>	1.21	4.41	10.89	16.51	4.06325
	<i>NAC74&NAC44</i>	1	2.25	14.44	17.69	4.205948
Clusters 1&4	<i>NAC12&NAC73</i>	8.41	38.44	31.36	78.21	8.843642
	<i>NAC12&NAC39</i>	7.29	43.56	32.49	83.34	9.129074
	<i>NAC12&NAC44</i>	6.76	36	38.44	81.2	9.011104
	<i>NAC45&NAC73</i>	7.29	40.96	30.25	78.5	8.860023
	<i>NAC45&NAC39</i>	6.25	46.24	31.36	83.85	9.156965
	<i>NAC45&NAC44</i>	5.76	38.44	37.21	81.41	9.022749
	<i>NAC09&NAC73</i>	9	30.25	30.25	69.5	8.336666
	<i>NAC09&NAC39</i>	7.84	34.81	31.36	74.01	8.602906
	<i>NAC09&NAC44</i>	7.29	28.09	37.21	72.59	8.519977
	<i>NAC11&NAC73</i>	9	34.81	28.09	71.9	8.479387
	<i>NAC11&NAC39</i>	7.84	39.69	29.16	76.69	8.757283
	<i>NAC11&NAC44</i>	7.29	32.49	34.81	74.59	8.63655
	<i>NAC19&NAC73</i>	7.29	34.81	30.25	72.35	8.50588
	<i>NAC19&NAC39</i>	6.25	39.69	31.36	77.3	8.792042
	<i>NAC19&NAC44</i>	5.76	32.49	37.21	75.46	8.686772
	<i>NAC47&NAC73</i>	7.29	29.16	29.16	65.61	8.1
	<i>NAC47&NAC39</i>	6.25	33.64	30.25	70.14	8.374963
	<i>NAC47&NAC44</i>	5.76	27.04	36	68.8	8.294577
	<i>NAC26&NAC73</i>	8.41	37.21	42.25	87.87	9.3739
	<i>NAC26&NAC39</i>	7.29	42.25	43.56	93.1	9.648834
	<i>NAC26&NAC44</i>	6.76	34.81	50.41	91.98	9.59062
	<i>NAC59&NAC73</i>	10.24	34.81	33.64	78.69	8.870738
	<i>NAC59&NAC39</i>	9	39.69	34.81	83.5	9.137833
	<i>NAC59&NAC44</i>	8.41	32.49	40.96	81.86	9.047652
	<i>NAC71&NAC73</i>	12.25	36	31.36	79.61	8.922444
	<i>NAC71&NAC39</i>	10.89	40.96	32.49	84.34	9.183681
	<i>NAC71&NAC44</i>	10.24	33.64	38.44	82.32	9.073037
	<i>NAC01&NAC73</i>	7.84	14.44	19.36	41.64	6.452906
	<i>NAC01&NAC39</i>	6.76	17.64	20.25	44.65	6.682066
	<i>NAC01&NAC44</i>	6.25	12.96	25	44.21	6.64906
	<i>NAC33&NAC73</i>	7.84	14.44	17.64	39.92	6.318228
	<i>NAC33&NAC39</i>	6.76	17.64	18.49	42.89	6.549046
	<i>NAC33&NAC44</i>	6.25	12.96	23.04	42.25	6.50
<i>NAC03&NAC73</i>	6.25	16.81	12.96	36.02	6.001666	
<i>NAC03&NAC39</i>	5.29	20.25	13.69	39.23	6.263386	
<i>NAC03&NAC44</i>	4.84	15.21	17.64	37.69	6.139218	
<i>NAC32&NAC73</i>	6.76	16	14.44	37.2	6.09918	
<i>NAC32&NAC39</i>	5.76	19.36	15.21	40.33	6.350591	
<i>NAC32&NAC44</i>	5.29	14.44	19.36	39.09	6.2522	
<i>NAC07&NAC73</i>	7.84	16.81	27.04	51.69	7.189576	

7. Appendices

	<i>NAC07&NAC39</i>	6.76	20.25	28.09	55.1	7.422937
	<i>NAC07&NAC44</i>	6.25	15.21	33.64	55.1	7.422937
	<i>NAC42&NAC73</i>	7.84	21.16	23.04	52.04	7.213876
	<i>NAC42&NAC39</i>	6.76	25	24.01	55.77	7.467931
	<i>NAC42&NAC44</i>	6.25	19.36	29.16	54.77	7.400676
	<i>NAC15&NAC73</i>	8.41	25	24.01	57.42	7.577599
	<i>NAC15&NAC39</i>	7.29	29.16	25	61.45	7.839005
	<i>NAC15&NAC44</i>	6.76	23.04	30.25	60.05	7.749194
	<i>NAC20&NAC73</i>	9	23.04	26.01	58.05	7.619055
	<i>NAC20&NAC39</i>	7.84	27.04	27.04	61.92	7.868926
	<i>NAC20&NAC44</i>	7.29	21.16	32.49	60.94	7.806408
	<i>NAC25&NAC73</i>	10.24	22.09	27.04	59.37	7.705193
	<i>NAC25&NAC39</i>	9	26.01	28.09	63.1	7.943551
	<i>NAC25&NAC44</i>	8.41	20.25	33.64	62.3	7.893035
	<i>NAC18&NAC73</i>	9	19.36	24.01	52.37	7.236712
	<i>NAC18&NAC39</i>	7.84	23.04	25	55.88	7.475293
	<i>NAC18&NAC44</i>	7.29	17.64	30.25	55.18	7.428324
	<i>NAC48&NAC73</i>	9	20.25	25	54.25	7.36546
	<i>NAC48&NAC39</i>	7.84	24.01	26.01	57.86	7.606576
	<i>NAC48&NAC44</i>	7.29	18.49	31.36	57.14	7.5591
	<i>NAC34&NAC73</i>	7.84	27.04	26.01	60.89	7.803204
	<i>NAC34&NAC39</i>	6.76	31.36	27.04	65.16	8.072174
	<i>NAC34&NAC44</i>	6.25	25	32.49	63.74	7.983733
	<i>NAC37&NAC73</i>	7.29	27.04	24.01	58.34	7.638063
	<i>NAC37&NAC39</i>	6.25	31.36	25	62.61	7.912648
	<i>NAC37&NAC44</i>	5.76	25	30.25	61.01	7.81089
	<i>NAC21&NAC73</i>	8.41	26.01	33.64	68.06	8.249848
	<i>NAC21&NAC39</i>	7.29	30.25	34.81	72.35	8.50588
	<i>NAC21&NAC44</i>	6.76	24.01	40.96	71.73	8.469357
	<i>NAC05&NAC73</i>	7.84	22.09	29.16	59.09	7.687002
	<i>NAC05&NAC39</i>	6.76	26.01	30.25	63.02	7.938514
	<i>NAC05&NAC44</i>	6.25	20.25	36	62.5	7.905694
	<i>NAC67&NAC73</i>	6.76	23.04	30.25	60.05	7.749194
	<i>NAC67&NAC39</i>	5.76	27.04	31.36	64.16	8.009994
	<i>NAC67&NAC44</i>	5.29	21.16	37.21	63.66	7.978722
Clusters 2&4	<i>NAC13&NAC73</i>	10.89	54.76	46.24	111.89	10.57781
	<i>NAC13&NAC39</i>	9.61	60.84	47.61	118.06	10.86554
	<i>NAC13&NAC44</i>	9	51.84	54.76	115.6	10.75174
	<i>NAC51&NAC73</i>	9	53.29	54.76	117.05	10.81896
	<i>NAC51&NAC39</i>	7.84	59.29	56.25	123.38	11.10766
	<i>NAC51&NAC44</i>	7.29	50.41	64	121.7	11.03177
	<i>NAC52&NAC73</i>	10.89	43.56	56.25	110.7	10.52141
	<i>NAC52&NAC39</i>	9.61	49	57.76	116.37	10.78749
	<i>NAC52&NAC44</i>	9	40.96	65.61	115.57	10.75035
	<i>NAC53&NAC73</i>	13.69	60.84	56.25	130.78	11.43591
	<i>NAC53&NAC39</i>	12.25	67.24	57.76	137.25	11.71537

7. Appendices

<i>NAC53&NAC44</i>	11.56	57.76	65.61	134.93	11.61594
<i>NAC54&NAC73</i>	10.24	51.84	60.84	122.92	11.08693
<i>NAC54&NAC39</i>	9	57.76	62.41	129.17	11.3653
<i>NAC54&NAC44</i>	8.41	49	70.56	127.97	11.31238
<i>NAC55&NAC73</i>	12.96	54.76	47.61	115.33	10.73918
<i>NAC55&NAC39</i>	11.56	60.84	49	121.4	11.01817
<i>NAC55&NAC44</i>	10.89	51.84	56.25	118.98	10.9078

Table 7.6. Gene information used for the subcellular localization adapted from Genoscope database.

Species	Locus Name	Gene ID	Length	Transcript Name	Orthologues in <i>A. thaliana</i>
Grape	GSVIVT01020 834001	<i>VvNAC36</i>	298	VIT_12s0028g00860	ANAC042 (AT2G43000.1)

Table 7.7. *VvNAC36* primers used in this study for transient expression vector construction.

Primer Name	Primer Sequences (5'-3')	Annealing Temperature	Amplification Length (Bp)
Forward: For_NAC042_5 gene	CACCACCTAAAGCTTGGACAAACA	60 °C	3000
Reverse: NAC pg_R	TCTACAGTCATAAACATGAGCTCG		




Figure 7.1 T-DNA region map of the construct pGWB605::*VvNAC36*

8. ACKNOWLEDGMENTS

- Firstly, I would like to express my deepest gratitude to my fiancée for the help and support through all the years and my family for their unwavering support throughout my PhD journey, and I am forever grateful.
- I would like to extend my gratitude to my supervisor, Prof. Dr. Erzsébet Kiss, for her support throughout my PhD study despite all the circumstances.
- I would like to extend my sincere gratitude to Dr. Veres Anikó and Dr. Tibor Nagy for their time, feedback, insightful comments, and guidance during the review of my dissertation. Your expertise and dedication have improved the quality of this work. Thank you for your contributions.
- Furthermore, I extend my appreciation to Prof. Dr. Lajos Helyes, the Head of the Plant Sciences Doctoral School at MATE, Prof. Dr. Zoltán Nagy, the Head of the Biological Sciences Doctoral School and Dr. János Balogh, the secretary at MATE Doctoral School of Biological Sciences, for their assistance and guidance.
- I would also like to acknowledge the valuable assistance provided by Zsuzsanna Tassy, International Coordinator, and the dedicated staff, including Mónika Törökné Hajdú and Edit Simáné Dolányi, at the Office of Doctoral and Habilitation Council of MATE, for their support during my studies.
- Lastly, I am grateful to the Stipendium Hungaricum, the scholarship program of the Hungarian government, for their financial support of this research. I would like to express my thanks to Csilla Kánai, Head of the International Relations Centre (IRC), for her cooperation and assistance.